

## REVIEW ARTICLE

## A Review on Antibacterial Phytochemical Constitutions Present in *Aerva lanata* and their Mode of Action Against Bacterial Biofilm

R. Rajasekaran, Yishak Gebrekidan

Department of Biology, College of Science, Eritrea Institute of Technology, Mai Nefhi, Asmara, Eritrea, North East Africa

Received: 17-12-2017; Revised: 18-01-2018 Accepted: 10-02-2018

### ABSTRACT

Antibacterial phytochemicals have unexplored chemical structures with high therapeutic potential, additionally; phytochemicals have several advantages, including green status, different mechanisms of action from antibiotics which could help to overcome the chemotherapeutic agent resistance problem and also ability to inhibit the growth of planktonic cell and biofilm. These phytochemicals are unmatched structural diversity, and it also has no target specific. In this study, an overview of the main classes of antibacterial phytochemicals present in *Aerva lanata* and their mode of action against bacterial biofilm is presented. A revision about the bacterial biofilm characteristics, biofilm formation, mechanism involved against antimicrobial agents, phytochemicals properties, and their targets to eradicate biofilm, anti-biofilm properties of various phytochemicals found in *A. lanata* is also done. The phytochemicals such as polyphenolics interfere with the adhesion potential, quorum sensing (QS) controlled, swarming motility and biofilm formation of *Escherichia coli*, and *Pseudomonas aeruginosa*. Catechin and tannic acid also present in *A. lanata* were able to promote a significant reduction in biofilm formation by *P. aeruginosa*, and it able to block biofilm formation by *E. coli* and *Pseudomonas putida*. Antibacterial phytochemicals isolated from the different plant part of *A. lanata* inhibited and reduced cell-surface adhesion, methicillin-resistant bacterial biofilm formation, inhibit bacterial motility, QS, and controls biofilms of *E. coli*, *P. aeruginosa*, and *Staphylococcus aureus*. Phenolic acids increased the susceptibility of dual species biofilms. Peptides react against bacterial biofilm by the process of cell membrane permeabilization, intracellular targets, inhibiting nucleic acids and protein synthesis, and cell wall adhesion of Gram-negative and Gram-positive bacteria.

**Keywords:** *Aerva lanata*, anti-biofilm activity, antibacterial phytochemical.

### INTRODUCTION

Medicinal plants are of great importance to the health of individuals and communities. The medicinal value of these plants lies in some chemical substances that produce a definite physiological action on the human body. The most important of these bioactive constituents of plants are alkaloids, tannins, flavonoids, and phenolic compounds.<sup>[1]</sup> Infectious diseases are the leading cause of death worldwide. Antibiotic resistance has become a global concern.<sup>[2]</sup> The clinical efficacy of many existing antibiotics is being threatened by the emergence of multidrug-resistant pathogens.<sup>[3]</sup>

Many infectious diseases have been known to be treated with herbal remedies throughout the history of mankind. Natural products, either as pure compounds or as standardized plant extracts, provide unlimited opportunities for new drug leads because of the unmatched availability of chemical diversity. There is a continuous and urgent need to discover new antimicrobial compounds with diverse chemical structures and novel mechanisms of action for new and reemerging infectious diseases.<sup>[4]</sup> Therefore, researchers are increasingly turning their attention to folk medicine, looking for new leads to develop better drugs against microbial infections.<sup>[5]</sup> The increasing failure of chemotherapeutics and antibiotic resistance exhibited by pathogenic microbial infectious agents has led to the screening of several medicinal plants, for their potential antimicrobial activity.<sup>[6,7]</sup>

#### \*Corresponding Author:

R. Rajasekaran,

Email: [rajasekharan.r@gmail.com](mailto:rajasekharan.r@gmail.com)

Secondary plant metabolites (phytochemicals), previously with unknown pharmacological activities, have been extensively investigated as a source of medicinal agents.<sup>[8]</sup> Thus, it is anticipated that phytochemicals with adequate antibacterial efficacy will be used for the treatment of bacterial infections.<sup>[9]</sup>

Advantage of using medicinal plants is that they do not cause any side effects when compared with synthetic drugs, because medicinal plants have high content of antioxidant compounds. This gives protective effects against diseases without reducing their therapeutic efficacy. Nowadays, herbal drugs have become world important objects, with both medicinal and economic implications.

## BACTERIAL BIOFILM DEFINITION AND CHARACTERISTICS

Biofilms are not easily defined as they vary greatly in structure and composition from one environmental niche to another. Microbial biofilms are extremely complex microbial ecosystems consisting of microorganisms attached to a surface and embedded in an organic polymer matrix of microbial origin. As well as microbial components, non-cellular materials such as mineral crystals, corrosion particles, clay or silt particles, or blood components, may also be found in the biofilm matrix. Biofilms, particularly in water systems, can be highly complex, while others such as those on medical devices, may be simpler, and composed of single, coccoid, or rod-shaped organisms.<sup>[10]</sup> Given these differences, it does not seem plausible to suggest that a true “biofilm model” can be defined that is applicable to every ecological, industrial and medical situation. Therefore, the definition of a biofilm has to be kept general and thus may be redefined as “microbial cells immobilized in a matrix of extracellular polymers acting as an independent functioning ecosystem, homeostatically regulated.”<sup>[11]</sup>

Surface polysaccharides also are used by bacteria as a means of adherence. In many natural settings, bacteria from multilayer communities called biofilms. Biofilms usually contain more than one species of bacteria. The first layer of the biofilm builds to a pili or some other attachment mechanism, then succeeding layers adhere to the first layer, using a polysaccharide slime to cement them together. Bacterial biofilm causes

a variety of problems, the national institute of health estimated that over 80% of microbial infections that occur in the human body involve biofilms. The most common diseases associated with biofilm formation are such as urinary tract infection (UTI) (*Escherichia coli* and *Klebsiella pneumonia*), burn wound (*Acinetobacter baumannii* and *Staphylococcus aureus*), dental plaque (*Streptococcus mutans*), respiratory infection (*Bordetella pertussis*), otitis media (*Haemophilus influenza*), and cystic fibrosis lung infection (*Pseudomonas aeruginosa*) are almost impossible to eliminate with antibiotics, necessitating the surgical removal of the implants. Bacteria in a biofilm are probably less susceptible to antibiotics than free-swimming (planktonic) bacteria because the antibiotic does not diffuse readily through the polysaccharide layer.<sup>[12]</sup>

## HISTORICAL PERSPECTIVES OF BIOFILM IN NATURE

In the 17<sup>th</sup> century, a dry-goods merchant named Antonie van Leeuwenhoek first observed “animalcules” swarming on living and dead matter. Leeuwenhoek’s curiosity and inventiveness were remarkable; he discovered these “animalcules” in the tartar on his own teeth and even after meticulous cleansing, the remaining opaque deposits isolated between his teeth were still “as thick as if it were batter.” These deposits contained a mat of various forms of “animalcules” that we now know were the bacteria of dental plaque. It is reasonable to suggest that this early study of dental plaque was the first documented evidence of the existence of microbial biofilms. Today, we, generally, define such biofilms as microbial communities adhered to a substratum and encased within an extracellular polymeric substance (EPS) produced by the microbial cells themselves. After van Leeuwenhoek’s early work, it was not until 1940 that the so-called “bottle effect” in marine microorganisms was first observed.<sup>[13]</sup> This showed that the growth of bacteria was substantially increased when they were attached to a surface. Further advancements in our knowledge of biofilms were made by Zobell in 1943 when he noted that bacteria on surfaces were greater in number compared with the surrounding seawater. From his studies, Zobell also postulated that the adhesion of bacteria consisted of a two-

stage process of reversible and then irreversible adhesion.

Despite the above studies being the first documented ones on biofilms, the extensive physical and chemical analysis of bacterial biofilms did not begin until the late 1960s and early 1970s, when a few investigators recognized the prevalence of bacterial biofilms<sup>[14-16]</sup> used scanning and transmission electron microscopy to examine biofilms on trickling filters in a wastewater treatment plant.<sup>[14]</sup> From this work, it was shown that biofilms were composed of a variety of different microorganisms and revealed that the matrix material or EPS was primarily composed of polysaccharides. The investigation of biofilms around this time was greatly aided by the use of electron microscopy, which provided information, not only on biofilm structure but also on the presence of EPS. In 1973, Characklis who investigated microbial slimes in industrial water revealed that biofilms were both tenacious and highly resistant to the antimicrobial effects of chlorine.<sup>[15]</sup> The true analysis of biofilms was not recognized until 1978. Many bacteria spent the majority of their existence within surface-attached, sessile communities. Work on dental plaque and sessile communities in mountain streams enabled<sup>[16]</sup> to hypothesize the mechanisms by which microorganism's adhered to living and non-living materials and derived benefit from this ecologic niche.

Since the 1970s, the study of biofilms in industrial, ecological, and medical settings has followed similar paths. Initial biofilm studies, generally, concentrated on composition, especially of the polymer matrix or "glycocalyx" that was thought to conserve and concentrate the digestive enzymes released by the bacteria, thus increasing the metabolic efficiency of the cells. Research<sup>[17]</sup> indicated that this glycocalyx also acted as anionic exchange matrix, trapping nutrients that were then transported into cells by highly efficient permeases. In 1981,<sup>[18]</sup> glycocalyx was characterized as a hydrated polyanionic polysaccharide matrix produced by polymerases affixed to the lipopolysaccharide component of the bacterial cell wall. In aqueous environments, biofilm production of glycocalyx is prevalent with organic and inorganic nutrients being concentrated at the solid/liquid interface. In addition, the glycocalyx provides a physical/chemical barrier

that offers partial protection against antibacterial agents.

Since biofilms form under diverse conditions, and may be composed of single or multiple species, the structures of various biofilms will necessarily have distinct features. Nevertheless, biophysical, structural, and chemical studies have led to a useful basic concept of a "biofilm model."<sup>[19]</sup> In this model, microorganisms form microcolonies surrounded by copious amounts of exopolysaccharide. Between the microcolonies are water-filled channels, and it has been proposed that these promote the influx of nutrients and the efflux of waste products.

### STAGES IN THE FORMATION OF BIOFILMS

The process of biofilm formation is complex, but generally recognized as consisting of five stages:<sup>[20]</sup>

1. Development of a surface conditioning film
2. Movement of microorganisms into close proximity with the surface
3. Adhesion (reversible and irreversible adhesion of microbes to the conditioned surface)
4. Growth and division of the organisms with the colonization of the surface, microcolony formation, and biofilm formation; phenotype and genotype changes
5. Biofilm cell detachment/dispersal.

### EPS AND THE GLYCOCALYX

EPS and glycocalyx are terms used to describe the polysaccharide produced by bacterial cells. EPS refers to one of the major components of biofilms, and glycocalyx refers to the polysaccharide matrix surrounding individual cells. EPS has an important role in biofilm structure and function and has a complex physical and chemical nature. Its functions are mostly protective in nature and this is one of the benefits for bacteria in the sessile state. Because the glycocalyx is the outermost component of bacterial cells, this layer mediates virtually all bacterial associations with surfaces and other cells: It dictates location, juxtaposition, and the eventual success in the ecosystem.<sup>[16]</sup> The physical properties of the biofilm are largely determined by the EPS, while the physiological properties

are determined by the bacterial cells. There are majorly eight processes in the development of biofilms.<sup>[21]</sup> These can be condensed to three main processes: The attachment of cells to a surface (colonization), growth of the attached cells into a mature biofilm, and the detachment of single cells (erosion) or large pieces (sloughing).

## BIOFILM STRUCTURE

Biofilm structure is the spatial arrangement of bacteria, cell clusters, EPS, and particulates. Since the structure can influence transport resistance, it is a significant determinant in the activity of the biofilm. Various conceptual and mathematical models have been proposed to describe the structure and function of biofilms.<sup>[21-23]</sup> Mathematical models describing transport, conversion, cell growth, and biofilm development are based on conceptual models. Biofilms and mats are matrices of cells and extracellular polymers (EPS). The EPS is produced by the cells and consists of polysaccharides, polyuronic acids, proteins, nucleic acids, and lipids.<sup>[24-26]</sup>

## MECHANISMS OF ANTIMICROBIAL RESISTANCE

The bacteria enclosed within the biofilm are extremely resistant to antibiotic treatments. Such resistance can be explained by different hypotheses, not necessarily limited to the following ones. First, the EPS secreted by biofilm bacteria, acts as a physical/chemical barrier, thus preventing penetration by antibodies or many antibiotics.<sup>[19]</sup> Second, embedded biofilm bacteria are generally not actively engaged in cell division and are smaller in size and less permeable to antibiotics. Third, the antibiotic degrading enzyme  $\beta$ -lactamase produced by several bacterial species can inhibit the activity of  $\beta$ -lactam ring structured antibiotics such as penicillins and cephamycins.  $\beta$ -lactamase may also be immobilized and accumulate in the EPS matrix so that the incoming antibiotic molecules can be inactivated effectively.<sup>[27]</sup> Fourth, up to 40% of the cell-wall protein composition of bacteria in biofilms is altered from that of its planktonic brethren. The membranes of biofilm bacteria might be better equipped to pump out antibiotics before they can cause damage, or

even antibiotics targets may disappear. Fifth, the antimicrobial agent is deactivated with the help of antimicrobial oxidants such as hypochlorite and  $H_2O_2$ . It is, however, known that the biofilm matrix does not form a completely impenetrable barrier to antimicrobial agents.<sup>[28]</sup>

There are six different characteristic mechanisms of bacterial resistance to antimicrobials.<sup>[29]</sup> They are (1) modification of the target site; (2) acquisition of alternative metabolic pathways to those inhibited by the drug; (3) alteration of permeability of the bacterial cell wall/membrane that restrict antibacterial agent access to target sites; (4) enzymatic modification or degradation of the antimicrobial agent; (5) over-expression of the drug target; and (6) active efflux pumps that extrude the antibiotic from the cell.

## TARGETS INVOLVED TO ERADICATE BIOFILM

Biofilm formation is regulated by combination of several mechanisms that are intrinsically related, such as adhesion, EPS synthesis, bacterial motility, and cell-to-cell interaction or QS.<sup>[30,31]</sup> Therefore, these cellular processes can be possible targets for the discovery of new drugs. This led to an increased interest in the search of natural products that have been proven to be able to restrict the capability of bacteria to adhere, communicate, and form complex biofilms.<sup>[32]</sup>

Phytochemicals may represent a natural antimicrobial strategy with considerable impact not only against free-living bacteria (planktonic) but also on bacterial biofilm formation.<sup>[33]</sup> Nevertheless, studies on biofilm prevention and control with phytochemicals are scarce. Diverse researchers already identified new strategies for biofilm control.<sup>[30,33,34]</sup> The use of phytochemicals in biofilm prevention and control is a relevant strategy. There are evidences that phytochemicals can interfere with diverse biofilm formation processes<sup>[35]</sup> such as inhibition of adhesion, degradation of EPS, interference with EPS production, motility inhibition adhesion, and QS inhibition. Other possible targets also possible, e.g., inhibition of cell viability of biofilms and reduction of metabolic activity of biofilm cells, Inhibition of biofilm formation, interference with virulence factors production, inhibition of nucleic acid synthesis, etc., are summarized in Table 1.

**Table 1:** Phytochemical targets involved to eradicate biofilm

S.no	Targets	Reference[s]
1	Microbial membrane associated with –OH group	[36,37]
2	Inhibition of adhesion	[38,39]
3	Motility inhibition (swimming and swarming), inhibition of EPS production	[40-42]
4	Antiquorum sensing or QSI	[43-49]
5	Inhibition of nucleic acid synthesis	[50]
6	Inhibition of enzyme involved in the radical generation, Destabilize, and permeabilize the cytoplasmic membrane	[50], [51-54]
7	Increase the membrane fluidity and permeability, disturb the membrane-embedded proteins, inhibit respiration and alter ion transport processes in both Gram-positive and Gram-negative bacteria	[55-57], [57-60]
8	Bind to sulfhydryl groups of external proteins of cell membrane	[61-63]
9	Disrupt the cell membrane; inhibit the nucleic acids and protein synthesis	[64-67]
10	Inhibits antibiotic efflux pumps	[68]
11	Intercalate with DNA, RNA polymerase, DNA gyrase, and topoisomerase IV	[69,70]
12	Attenuate the level of penicillin-binding protein	[71]
13	Inhibition of cell viability of biofilms	[72]
14	Interference with QS and inhibition of biofilm formation. Decrease of EPS production, reduction of metabolic activity of biofilm cells	[73-79]
15	Interference with virulence factors production	[80,81]
16	Reduction of viable bacterial cells counts of multispecies biofilms	[82]
17	Degradation of cell wall, disruption of cytoplasmic membrane, damage of membrane proteins, leakage of intracellular contents, coagulation of cytoplasm, and depletion of proton	[36,37,51,83,84]
18	Interference with motility, adhesion, biofilm formation, and QSI	[85-87]
19	Inhibition of enzymes $\beta$ -lactamases	[88]

QSI: QS inhibition, EPS: Extracellular polymeric substance

## IMPORTANCE AND PROPERTIES OF ANTIBACTERIAL PHYTOCHEMICALS

Phytochemicals have demonstrated distinctive properties.<sup>[33]</sup> Unlike synthetic molecules, phytochemical products display an unmatched structural diversity with complex and novel multilayer mechanisms of action. The synthetic chemotherapeutic agent are react against microbial growth in a specific manner (single target) such as some molecules breakdown the bacterial cell wall, some molecules alternation in membrane function etc, so these all are target specific. All the chemotherapeutic agent, targets and their functions are controlled by specific genes present in the pathogenic organism. Some currently used antibiotics act also through multiple modes of action (multiple molecular targets and/or targets encoded by multiple genes.<sup>[89]</sup> Therefore, compounds that inhibit bacterial growth by different mechanisms than the presently used by conventional antibiotics can provide an interesting approach to control drug-resistant infections. Moreover, contrarily to the previously considered strategy “one drug, one target, one disease,” it is now extensively recognized that the use of a single molecule able to operate simultaneously

in various targets is more advantageous for the treatment of complex infectious diseases.<sup>[90]</sup> In fact, there are no evidences on the emergence of resistance to phytochemicals.

The antibacterial mechanism of action of phytochemicals is not completely understood.<sup>[33]</sup> Hence, more studies are needed to know their exact antimicrobial targets. Degradation of the cell wall, disruption of cytoplasmic membrane, damage of membrane proteins, leakage of intracellular contents, coagulation of cytoplasm, and depletion of proton have been currently reported as the mechanisms responsible for cell death, caused by some of these compounds<sup>[36,37,51,83,84]</sup> Useful phytochemicals with antimicrobial activity can be divided into several classes that include: Phenolics and polyphenolics, terpenoids, and other essential oils constituents, alkaloids, lectins, peptides, and polyacetylenes among much other.<sup>[91,92]</sup>

Phytoconstituents employed by plants to protect them against pathogenic insects, bacteria, fungi, or protozoa have found applications in human medicine.<sup>[93]</sup> Some phytochemicals such as phenolic acids act essentially by helping in the reduction of particular adherence of organisms to the cells lining the bladder, and the teeth, which ultimately lowers the incidence of UTI and the

usual dental caries. Plants can also exert either bacteriostatic or bactericidal activity of microbes.

### **AERVA LANATA PLANT DESCRIPTION**

*A. lanata* belongs to the family Amaranthaceae. This family consists of about 169 genera and 2300 species. *A. lanata* (L) Juss Ex. Schult commonly called Polphala of Amaranthaceae is a perennial shrub which is seen commonly in different waste parts of India, Eritrea and another part of the world. Most of the plants of this family are herbs, erect/with climbing branches. Leaves are opposite or alternate, ex-stipulate. Flowers are usually hermaphrodite, small usually in terminal simple or paniculate spikes, cymes or cluster; bracts hyaline never leafy, bracteoles. Fruits are membranous utricle, irregularly rupturing capsule, rarely berry. Seeds inverted or erect, orbicular or kidney shaped.<sup>[94]</sup> *A. lanata* grows wild on the mountain slopes, fields, and bare patches of ground up to an altitude, 900 m in the hills, and a native of Asia, Africa, and Australia.<sup>[94-97]</sup>

### **PHYTOCHEMICAL CONSTITUTIONS IN *A. LANATA***

Phytochemical studies have been carried out by several workers with the report of different kinds of bioactive compounds such as alkaloids, flavonoids, oxalate, phytic acid, phytin phosphorus, phenolic compounds, phytosterols, carbohydrates, proteins, amino acids, quinines, anthraquinones, catechins, coumarins, phenols, quinones, saponins, steroids, glycosides, tannins, and xanthoproteins.<sup>[98,99]</sup> The phytochemical constituents present in the plant include alkaloids (ervine, methylervine, ervoside, aervine, methylaervine, aervoside, ervolanine, and aervolanine) and flavonoids (kaempferol, quercetin, isorhamnetin, persinol, and persinosides A and B).<sup>[100]</sup> The whole plant of *A. lanata* contains  $\beta$ -sitosterol,  $\alpha$ -amyirin, betulin, hentriacontane, sitosteryl palmitate, D-glucoside, glycosides, kaempferol-3-galactoside and kaempferol-3-rhamnoglucoside, starch, and free: Sugars (fructose, galactose, rhamnose, and sucrose).<sup>[95,101]</sup>

The antinutrient levels also revealed the presence of tannic acid, saponin, alkaloids, flavonoids, and oxalate. The phytic acid and phytin phosphorus

were also in low amount.<sup>[102-105]</sup> *A. lanata* also contains miscellaneous phytoconstituents such as methyl grevillate, lupeol, lupeol acetate benzoic acid,  $\beta$ -sitosteryl acetate, and tannic acid.<sup>[102]</sup> Different studies revealed the presence of 30 different types of steroids, 21 different types of saponins, 27 varieties of terpenoids, and 24 types of tannins in the methanolic extract of root, stem, leaves, and seeds of *A. lanata*<sup>[106-108]</sup> using high-performance thin-layer chromatography. Fourier-transform infrared (FT-IR) spectral analysis<sup>[109]</sup> of plant parts such as flowers, leaves, stem, and roots of *A. lanata* showed the presence of characteristic functional groups of alcohols, phenols, alkanes, carboxylic acids, aldehydes, alkenes, nitro compounds, alcohols, carboxylic acids, esters, aliphatic amines, and alkyl halides compounds derivatives which were responsible for medicinal properties of this plants.

The presence of different kinds of bioactive compounds particularly alkaloids such as canthin-6-one and beta-carboline, aervine (10-hydroxycanthin-6-one), methylaervine (10-methoxycanthin-6-one), aervoside (10- $\beta$ -Dglucopyranosyloxycanthin-6-one), and aervolanine (6-methoxy- $\beta$ -carboline-1-yl) propionic acid from leaves of *A. lanata*.<sup>[110-116]</sup> The alkaloids, phenolic compounds, phytosterols, carbohydrates, proteins, amino acids, flavonoids, and quinones were identified in different solvents extracts.<sup>[117]</sup>

A new type diterpene was identified<sup>[118]</sup> in methanol extract of dried seeds of the *A. lanata*. Qualitative phytochemical analysis<sup>[119]</sup> of the methanol and ethanol extracts prepared from *A. lanata* whole plant revealed the presence of alkaloids, anthraquinones, catechins, coumarins, flavonoids, phenols, quinones, saponins, steroids, sugar, glycosides, tannins, and xanthoproteins. The FT-IR spectrum confirmed the presence of alkyl group, methyl group, alcohol group, ethers, esters, carboxylic acid, and anhydrides. Phytochemical analysis carried out for leaf extracts of the plant showed the presence of several secondary metabolites. The methanol extract of leaf showed the highest phenolic and flavonoid content.<sup>[120]</sup> The isolation of above-mentioned bioactive compounds from the different research and review report concluded the whole plant of *A. lanata* would be useful to find out the novel drugs.

## ANTIBACTERIAL PHYTOCHEMICALS PRESENT IN *A. LANATA* AND THEIR MODE OF ACTION AGAINST BACTERIAL GROWTH

Natural products from various plants have a potential control microbial growth in diverse situations and the specific case of disease treatment, and numerous studies have aimed to describe the chemical composition of different plant antimicrobials and the mechanisms involved in microbial growth inhibition.

Phytochemicals are non-proteinaceous secondary metabolites or non-nutritive bioactive compounds as protective agents against external pathogenic stress.<sup>[121]</sup> Some of the phytochemicals interfere with the phospholipids bilayer of the cell membrane which has as a consequence a permeability increase and loss of cellular constituents, damage of the enzymes involved in the production of cellular energy and synthesis of structural components, and destruction or inactivation of genetic material. In general, the mechanism of action is considered to be the disturbance of the cytoplasmic membrane, disrupting the proton motive force, electron flow, active transport, and coagulation of cell contents<sup>[122]</sup>. Phytochemicals induce membrane permeabilization by forming pores or disrupting membrane integrity and induced lysis of cell wall. In general, phytochemicals have diverse antimicrobial mechanisms including damaging cell wall and cytoplasmic membrane as shown in Table 1.

Antimicrobial activity results of the same plant part tested most of the time varied from researcher to researcher. This is possible because concentration of plant constituents of the same plant organ can vary from one geographical location to another depending on the age of the plant, differences in topographical factors, the nutrient concentrations of the soil, extraction method as well as method used for antimicrobial study. It is, therefore, important that scientific protocols be clearly identified and adequately followed and reported. The different phytochemical constitutions present in *A. lanata* and their mode of antibacterial actions will be presented below.

### Alkaloids

Numerous plant families are known to produce alkaloids and have been reported that several of

them possess high antimicrobial activity and could, therefore, be a good alternative for actual drugs.<sup>[37]</sup> Number of reports revealed extracts from different parts of *A. lanata* containing alkaloids<sup>[98-105,110-117,119]</sup> showed antimicrobial activity against different multidrug-resistant bacteria, namely, *E. coli*, *P. aeruginosa*, *P. mirabilis*, *S. aureus*, *B. subtilis*, *Raoultella planticola*, *Enterobacter aerogenes*, *Agrobacterium tumefaciens*, and *K. pneumonia*.<sup>[123]</sup> Their mechanism of action can be attributed to their ability to increase membrane permeability and to intercalate with DNA. RNA polymerase, DNA gyrase, and topoisomerase IV are also possible targets.<sup>[69,70,124]</sup>

### Anthraquinones

These are derivatives of phenolic and glycosidic compound phytochemical present in *A. lanata*.<sup>[119]</sup> They are solely derived from anthracene giving variable oxidized derivatives such as anthrones and anthranols.<sup>[124,125]</sup>

### Catechin

Catechins are one of the subclasses/components of flavonoids present in different plants including *A. lanata*<sup>[119]</sup> have been identified as potent antimicrobial agents and were suggested as a therapeutic possibility.<sup>[126]</sup> It forms complexes with the bacterial cell wall of intestinal microorganisms.<sup>[127]</sup>

### Flavonoids

Flavonoids are one of the biggest classes of secondary metabolites found in various types of edible plants including *A. lanata*.<sup>[98-100,102-105,117,119,120]</sup> They have been identified as potent antimicrobial agents and were suggested as a therapeutic possibility.<sup>[126]</sup> Their activity is arguably due to the ability to form a complex with extracellular proteins, which then binds to the bacterial cell wall, increasing their permeability of the inner membrane of *E. coli* and also dissipation of membrane potential.<sup>[128]</sup> Moreover, their targets are the membranes with -OH groups.<sup>[36,37]</sup> Interference with metabolism and inhibition of nucleic acid synthesis was also reported as possible mechanisms of action.<sup>[50]</sup> Quinones also found in *A. lanata*,<sup>[124]</sup> it is a flavonoids derivative, the mechanism of bioactivity also similar as

flavonoids. Some of the flavonoid subclasses (quercetin and robinetin) found in various types of edible plants. This flavonoid can inhibit DNA gyrase, inactivate specific bacterial enzymes, inhibit antibiotic efflux in methicillin-resistant *S. aureus* (MRSA),<sup>[68]</sup> and inhibit the synthesis of nucleic acids of both Gram-negative and Gram-positive bacteria.<sup>[50]</sup> There are no evidences of these flavonoid subclasses found in *A. lanata*. More work is needed to be done to establish this assumption.

### Phenols

Phenolic compounds constitute one of the most diverse groups of phytochemicals, being widely distributed in plants and protecting them from microbial infections. They have antioxidant properties but are also potent anti-infective.<sup>[37,129]</sup> Phytochemical studies have been carried out by several workers with the report various solvent extract of *A. lanata* contain phenol.<sup>[109,117,119,120,124]</sup> The antimicrobial activity of plant phenols has been extensively studied against human pathogens, to characterize and develop new healthy food ingredients, medical compounds, and pharmaceuticals.<sup>[37,129]</sup> The phenolic derivatives differ in the patterns of the hydroxylations and methoxylations of their aromatic rings. The common hydroxycinnamic acids are ferulic acid and vanillic acid found in *A. lanata*. Their antimicrobial activity can be due to their ability to destabilize and permeabilize the cytoplasmic membrane, inhibition of enzymes involved in the radical generation and also the inhibition of the synthesis of nucleic acids of bacteria.<sup>[50-54]</sup> Antibacterial activity was also obtained ferulic acids against lactic acid bacteria (*Lactobacillus plantarum*, *Lactobacillus hammesii*, and *Listeria monocytogenes*), *E. coli*, *P. aeruginosa*, and *S. aureus* (including MRSA).<sup>[31,130-134]</sup> Moreover, it was observed synergistic effects between these compounds and the antibiotic streptomycin.<sup>[134]</sup>

### Steroids and glycosides

Plant steroids (or steroid glycosides) also referred to as “cardiac glycosides” are one of the most naturally occurring plant phytoconstituents that have found therapeutic applications as arrow poisons or cardiac drugs.<sup>[126]</sup> The phytochemical also reported in *A. lanata*<sup>[95,98,99,101,106-108,117,119,136-138]</sup> with antibacterial actions have been reported as

anti-inflammatory or cytotoxic activity against different harmful bacteria.<sup>[135]</sup>

### Tannins

Tannins are found in almost every plant part including *A. lanata*<sup>[107,109,139]</sup> nutritional and biological properties of tannins have been described.<sup>[140]</sup> In addition, antibacterial actions of tannins have been reported as bacteriostatic and bactericidal against different harmful bacteria, including *E. coli*, *Pseudomonas* spp., *Salmonella* spp., *Staphylococcus* spp., and *Streptococcus* spp.<sup>[141-143]</sup> Their mode of action is apparently related to their ability to inactivate microbial adhesins, enzymes, membrane proteins, and formation of complexes with cell wall. Furthermore, they can form complex with polysaccharide, which is suggested to be the main reason for their inhibitory effects on bacteria.<sup>[144,145]</sup>

### Terpenoids

Terpenoids are the largest group of natural compounds reported in *A. lanata*.<sup>[119,120,124,136,137,146,147]</sup> These bioactive products have a lot of biological properties, including antioxidant and antimicrobial activities. Due to their recognized antimicrobial potential, terpenoids have been the subject of several studies along the years.<sup>[148]</sup> The mechanism of action of terpenoids is not fully understood, but it is speculated that involves membrane disruption by the lipophilic compounds and their activity depend largely of the structure of the compound, as recently demonstrated by some authors.<sup>[55-57]</sup> This antibacterial action can result in the increase of membrane fluidity/permeability, disruption of membrane-embedded proteins, and change of ion transport processes in both Gram-positive and Gram-negative bacteria.<sup>[55-57]</sup> The triterpenes include steroids, sterols, and cardiac glycosides with anti-inflammatory, sedative, insecticidal, or cytotoxic activity.<sup>[135]</sup>

### Coumarin

Few researchers reported coumarin is phytochemical compound present in *A. lanata*.<sup>[98,99,119]</sup> It is a phenolic compound with fused benzene and pyrone groups<sup>[51,36,83]</sup> had broad-spectrum antibacterial activities against



*S. aureus*, *E. coli*, *S. typhimurium*, *Salmonella enteritidis*, *A. hydrophila*, *Yersinia* sp., *Shigella* sp., and *Vibrio parahaemolyticus*.<sup>[37]</sup>

### Amino acids and peptides

Short-length peptides (between 15 and 30 amino acids) with microbicidal activity are commonly named as antimicrobial peptides.<sup>[65]</sup> These biologically active molecules are an important component of the innate immune system of wide variety of organisms.<sup>[149,150]</sup> They comprise several protein groups with different features, as regard to the total charge of the molecule and the content of disulfide bonds.<sup>[66]</sup> Peptides with antimicrobial properties are present in all organs of a variety of plant species including *A. lanata*<sup>[117]</sup> constitutively or in response to microbial infections.<sup>[151-153]</sup> Antimicrobial peptides are effective against a wide range of microorganisms, namely, Gram-negative and Gram-positive bacteria, including multidrug-resistant strains,<sup>[67,154]</sup> Their mechanism of action is believed be the damage or destabilization of the microbial cell membranes by formation of ion channels, transmembrane pores, or extensive membrane rupture.<sup>[64,65]</sup>

### PHYTOCHEMICALS PRESENT IN *A. LANATA* AND THEIR MODE OF ACTION AGAINST BACTERIAL BIOFILM

Antibiotic resistance is a significant public health problem that is worsened when microorganisms are in biofilms.<sup>[155-157]</sup> The main weapons used to control harmful biofilms have been the antimicrobial products; nonetheless, there are no antimicrobials with ensured efficacy.<sup>[34]</sup> Natural products that have been proven to be able to restrict the capability of bacteria to adhere, communicate, and form complex biofilms.<sup>[32]</sup> There are evidences that phytochemicals can interfere with diverse biofilm formation processes.<sup>[30,33]</sup> Biofilm formation is regulated mainly by adhesion, EPS synthesis, bacterial motility, and QS.<sup>[30,84]</sup> Not all of the phytochemicals present in *A. lanata* involved to control the biofilm formation and growth. However, very few of the phytochemicals and their derivatives present in *A. lanata* have anti-biofilm properties. The phytochemicals present in *A. lanata* have an anti-biofilm properties are discussed below. The anti-biofilm phytochemicals are not target specific because the phytochemicals

are multi target compounds and it interfere or inhibit the biofilm formation and growth in different mechanism compare to synthetic drug. Polyphenolics demonstrated the ability to interfere with the adhesion potential of *Streptococcus*.<sup>[158-160]</sup> Polyphenol, ellagic acid, and glycosylated derivatives inhibited biofilm formation of *S. aureus*.<sup>[161]</sup> Pure ellagic acid also displayed anti-biofilm properties against *S. aureus* and *E. coli*.<sup>[161,162]</sup> Polyphenol, catechin, and tannic acid were able to promote a significant reduction in biofilm formation by *P. aeruginosa*.<sup>[163]</sup> Polyphenol inhibited QS controlled, swarming motility and biofilm formation of *E. coli* K-12 and *P. aeruginosa* PAO1 in a concentration-dependent manner.<sup>[164]</sup> Bioactive fraction isolated from leaves of *A. lanata* contains alkaloids, tannins, saponins, steroids, glycosides, and flavonoids inhibited MRSA biofilm formation. Anti-biofilm activities of phytochemicals reduce cell-surface adhesion and attenuate the level of penicillin-binding protein. Ferulic acids present in *A. lanata*<sup>[50-54]</sup> to inhibit bacterial motility, adhesion and to prevent and control biofilms of *E. coli*, *P. aeruginosa*, *S. aureus*, and *L. monocytogenes*. Catechin and tannic acid also present in *A. lanata*<sup>[165]</sup> were able to block N-acyl homoserine lactones synthesis<sup>[73]</sup> and biofilm formation<sup>[74]</sup> of *E. coli* and *Pseudomonas putida*. Moreover, (-)-epigallocatechin gallate present in *A. lanata* inhibited biofilm formation of *Staphylococcus* spp. by reduction of EPS production.<sup>[75]</sup> Ferulic acid is a phenolic derivatives hydroxycinnamic acid phytochemical present in *A. lanata* were able to inhibit biofilm formation of *S. aureus*.<sup>[166]</sup> Ferulic inhibit swimming motility and QS<sup>[167]</sup> of *Bacillus cereus* and *Pseudomonas fluorescens*. In addition, the development of biofilms in the presence of these phenolic acids increased the susceptibility of dual-species biofilms (*B. cereus* and *P. fluorescens*) to a second exposure to the chemicals. Several researchers reported *A. lanata* contain phenolic acids it increased the susceptibility of dual-species biofilms.<sup>[168]</sup> The antimicrobial peptides present in *A. lanata* react against bacterial biofilm by the process of cell membrane permeabilization, antimicrobial peptides can also act on intracellular targets, inhibiting nucleic acids and protein synthesis, and enzymatic activity.<sup>[64,67]</sup> These peptides have the ability to inhibit biofilm growth by competitive inhibition of adhesion of microbial proteins to host polysaccharide receptors has also been observed.<sup>[51]</sup> Due to their cationic and hydrophobic features,

antibacterial phytochemicals interact primarily with negatively charged components of the bacterial envelope, such as lipopolysaccharides of the outer membrane of Gram-negative bacteria or lipoteichoic acids present on the cell wall of both Gram-negative and Gram-positive bacteria.<sup>[64,65,151,169]</sup>

## CONCLUSION

Most of the synthetic and microbial-derived chemotherapeutic (Antibiotics) agents are failed to control the growth of the microbial life involved in the biofilm. Bacterial members involved to form biofilm are multidrug-resistant microbes; it causes a variety of problems including human diseases such as a dental plaque and urinary tract infection. The bacterial biofilm has variety of special mechanism to neutralize, tolerate and escape the antimicrobials. The biochemical features are also completely differing from planktonic cell. Therefore, new compounds are required to eradicate the biofilm. Plant secondary metabolites (phytochemicals) play a key role in plants defense and have evolve to inhibit the growth of biofilm bacteria and planktonic cell; these are simply known as antibacterial phytochemicals. Compare to the antimicrobials antibacterial phytochemicals are structurally differ, and their mode of action and targets also completely differ from chemotherapeutic agents. The *A. lanata* is widely used the herb. The pharmacological studies conform and support the therapeutic utility of the plant in various disorders mainly in diseases of the urinary system. These plants contain numerous phytochemicals among that few have an anti-biofilm activity. However, more work should be needed to isolate, identified and purified the main classes/subclasses of anti-biofilm phytochemicals present in *A. lanata*.

## REFERENCES

1. Hill AF. Economic Botany. A Textbook of useful Plants and Plant Products. 2<sup>nd</sup> ed. New York: McGraw-Hill Book Company Inc.; 1952.
2. Westh H, Zinn CS, Rosdahl VT. An international multicenter study of antimicrobial consumption and resistance in *Staphylococcus aureus* isolates from 15 hospitals in 14 countries. *Microb Drug Resist* 2004;10:169-76.
3. Bandow JE1, Brötz H, Leichert LI, Labischinski H, Hecker M. Proteomic approach to understanding antibiotic action. *Antimicrob Agents Chemother* 2003;47:948-55.
4. Rojas R1, Bustamante B, Bauer J, Fernández I, Albán J, Lock O. Antimicrobial activity of selected Peruvian medicinal plants. *J Ethnopharmacol* 2003;88:199-204.
5. Benkeblia N. Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*). *Lebensm Wiss u-Technol* 2004;37:263-8.
6. Colombo ML, Bosisio E. Pharmacological activities of *Chelidonium majus* L. (*Papaveraceae*). *Pharmacol Res* 1996;33:127-34.
7. Iwu MW, Duncan AR, Okunji CO. New antimicrobials of plant origin. In: Janick J, editor. *Perspectives on New Crops and New Uses*. Alexandria, VA: ASHS Press; 1999. p. 457-62.
8. Krishnaraju AV, Rao TV, Sundararaju D, Vanisree M, Tsay H, Gottumukkala V, et al. Assessment of bioactivity of Indian medicinal plants using Brine shrimp (*Artemia salina*) lethality assay. *Int J Appl Sci Eng* 2005;2:125-34.
9. Balandrin MF, Klocke JA, Wurtele ES, Bollinger WH. Natural plant chemicals: Sources of industrial and medicinal materials. *Science* 1985;228:1154-60.
10. Percival SL, Malic S, Cruz H, Williams DW. Introduction to biofilms. In: Percival SL, Knottenbelt DC, Cochrane CA, editors. *Biofilms and Veterinary Medicine*, Springer Series on Biofilms 6. Berlin Heidelberg: Springer-Verlag; 2011. p. 41-68.
11. Percival SL, Walker J, Hunter P. *Microbiological Aspects of Biofilms and Drinking Water*. New York: CRC Press; 2000.
12. Salyers AA, Whitt DD. Structural features of prokaryotes. In: *Microbiology, Diversity, Disease, and the Environment*. Urbana-Champaign: University of Illinois, Fitzgerald Science Press; 2001. p. 55.
13. Heukelekian H, Heller A. Relation between food concentration and surface for bacterial growth. *J Bacteriol* 1940;40:547-58.
14. Jones HC, Roth IL, Saunders WM. Electron microscopic study of a slime layer. *J Bacteriol* 1969;99:316-25.
15. Characklis WG. Attached microbial growths-II. Frictional resistance due to microbial slimes. *Water Res* 1973;7:1249-58.
16. Costerton JW, Geesey GG, Cheng KJ. How bacteria stick. *Sci Am* 1978;238:86-95.
17. Costerton JW, Geesey GG. In: Costerton JW, Colwell RR, editors. *Native Aquatic Bacteria: Enumeration, Activity, and Ecology*. Philadelphia, PA: ASTM Press; 1979. p. 7-1.
18. Costerton JW, Irvin RT, Cheng KJ. The bacterial glycocalyx in nature and disease. *Annu Rev Microbiol* 1981;35:299-324.
19. Costerton JW, Lewandowski Z, Caldwell DE, Korber DR, Lappin-Scott HM. Microbial biofilms. *Annu Rev Microbiol* 1995;49:711-45.
20. Palmer RJ Jr., White DC. Developmental biology of biofilms: Implications for treatment and control. *Trends Microbiol* 1997;5:435-40.
21. Characklis WG. Microbial fouling. In: Characklis WG, Marshall KC, editors. *Biofilms*. New York: John Wiley and Sons; 1990. p. 523-84.
22. Wanner O, Gujer W. A multispecies biofilm model. *Biotechnol Bioeng* 1986;28:314-28.

23. Rittmann BE, Manem JA. Development and experimental evaluation of a steady-state, multispecies biofilm model. *Biotechnol Bioeng* 1992;39:914-22.
24. Decho AW. Microbial exopolymer secretions in ocean environments: Their role(s) in food webs and marine processes. *Oceanogr Mar Biol Ann Rev* 1990;28:73-153.
25. Decho AW, Lopez GR. Exopolymer microenvironments of microbial flora: Multiple and interactive effects on trophic relationships. *Limnol Oceanogr* 1993;38:1633-45.
26. Schmidt JE, Ahring BK. Extracellular polymers in granular sludge from different upflow anaerobic sludge blanket (UASB) reactors. *Appl Microbiol Biotechnol* 1994;42:457-62.
27. Anderl JN, Zahller J, Roe F, Stewart PS. Role of nutrient limitation and stationary-phase existence in *Klebsiella pneumoniae* biofilm resistance to ampicillin and ciprofloxacin. *Antimicrob Agents Chemother* 2003;47:1251-6.
28. Percival SL, Bowler PG. Biofilms and their potential role in wound healing. *Wounds* 2004;16:234-40.
29. Coates A, Hu Y, Bax R, Page C. The future challenges facing the development of new antimicrobial drugs. *Nat Rev Drug Discov* 2002;1:895-910.
30. Landini P, Antoniani D, Burgess JG, Nijland R. Molecular mechanisms of compounds affecting bacterial biofilm formation and dispersal. *Appl Microbiol Biotechnol* 2010;86:813-23.
31. Borges A, Serra S, Cristina Abreu A, Saavedra MJ, Salgado A, Simões M, *et al.* Evaluation of the effects of selected phytochemicals on quorum sensing inhibition and *in vitro* cytotoxicity. *Biofouling* 2014;30:183-95.
32. Borges A, Simões LC, Saavedra MJ, Simões M. The action of selected isothiocyanates on bacterial biofilm prevention and control. *Int Biodeterior Biodegradation* 2014;86:25-33.
33. Simões M, Bennett RN, Rosa EA. Understanding antimicrobial activities of phytochemicals against multidrug resistant bacteria and biofilms. *Nat Prod Rep* 2009;26:746-57.
34. Simões M. Antimicrobial strategies effective against infectious bacterial biofilms. *Curr Med Chem* 2011;18:2129-45.
35. Borges A, Saavedra MJ, Simões M. Insights on antimicrobial resistance, biofilms and the use of phytochemicals as new antimicrobial agents. *Curr Med Chem* 2015;22:2590-614.
36. Samy RP, Gopalakrishnakone P. Therapeutic potential of plants as anti-microbials for drug discovery. *Evid based complement. Alternat Med* 2010;7:283-94.
37. Saleem M, Nazir M, Ali MS, Hussain H, Lee YS, Riaz N, *et al.* Antimicrobial natural products: An update on future antibiotic drug candidates. *Nat Prod Rep* 2010;27:238-54.
38. Rukayadi Y, Hwang JK. Effect of coating the wells of a polystyrene microtiter plate with xanthorrhizol on the biofilm formation of *Streptococcus mutans*. *J Basic Microbiol* 2006;46:410-5.
39. Lee KH, Kim BS, Keum KS, Yu HH, Kim YH, Chang BS, *et al.* Essential oil of curcuma longa inhibits *Streptococcus mutans* biofilm formation. *J Food Sci* 2011;76:H226-30.
40. Shi W, Sun H. Type IV pilus-dependent motility and its possible role in bacterial pathogenesis. *Infect Immun* 2002;70:1-4.
41. Packiavathy IA, Agilandeeswari P, Musthafa KS, Pandian SK, Veera A, Ravi A. Antibiofilm and quorum sensing inhibitory potential of *Cuminum cyminum* and its secondary metabolite methyl eugenol against gram negative bacterial pathogens. *Food Res Int* 2012;45:85-92.
42. Duan Q, Zhou M, Zhu L, Zhu G. Flagella and bacterial pathogenicity. *J Basic Microbiol* 2013;53:1-8.
43. Bjarnsholt T, Jensen PØ, Rasmussen TB, Christophersen L, Calum H, Hentzer M, *et al.* Garlic blocks quorum sensing and promotes rapid clearing of pulmonary *Pseudomonas aeruginosa* infections. *Microbiology* 2005;151:3873-80.
44. Persson T, Hansen TH, Rasmussen TB, Skindersø ME, Givskov M, Nielsen J, *et al.* Rational design and synthesis of new quorum-sensing inhibitors derived from acylated homoserine lactones and natural products from garlic. *Org Biomol Chem* 2005;3:253-62.
45. Slusarenko A, Patel A, Portz D. Control of plant diseases by natural products: Allicin from garlic as a case study. *Eur J Plant Pathol* 2008;121:313-22.
46. Khan MS, Zahin M, Hasan S, Husain FM, Ahmad I. Inhibition of quorum sensing regulated bacterial functions by plant essential oils with special reference to clove oil. *Lett Appl Microbiol* 2009;49:354-60.
47. Jakobsen TH, Bragason SK, Phipps RK, Christensen LD, van Gennip M, Alhede M, *et al.* Food as a source for quorum sensing inhibitors: Iberin from horseradish revealed as a quorum sensing inhibitor of *Pseudomonas aeruginosa*. *Appl Environ Microbiol* 2012;78:2410-21.
48. Jakobsen TH, van Gennip M, Phipps RK, Shanmugham MS, Christensen LD, Alhede M, *et al.* Ajoene, a sulfur-rich molecule from garlic, inhibits genes controlled by quorum sensing. *Antimicrob Agents Chemother* 2012;56:2314-25.
49. Cady NC, McKean KA, Behnke J, Kubec R, Mosier AP, Kasper SH, *et al.* Inhibition of biofilm formation, quorum sensing and infection in *Pseudomonas aeruginosa* by natural products-inspired organosulfur compounds. *PLoS One* 2012;7:38492.
50. Cushnie TP, Lamb AJ. Antimicrobial activity of flavonoids. *Int J Antimicrob Agents* 2005;26:343-56.
51. Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999;12:564-82.
52. Puupponen-Pimiä R, Nohynek L, Alakomi HL, Oksman-Caldentey KM. Bioactive berry compounds—novel tools against human pathogens. *Appl Microbiol Biotechnol* 2005;67:8-18.
53. Muthuswamy S, Rupasinghe HP. Fruit phenolics as natural antimicrobial agents: Selective antimicrobial activity of catechin, chlorogenic acid and phloridzin. *J Food Agric Environ* 2007;5:81-5.
54. Maddox CE, Laur LM, Tian L. Antibacterial activity of phenolic compounds against the phytopathogen *Xylella fastidiosa*. *Curr Microbiol* 2010;60:53-8.
55. Trombetta D, Castelli F, Sarpietro MG, Venuti V, Cristani M, Daniele C, *et al.* Mechanisms of

- antibacterial action of three monoterpenes. *Antimicrob Agents Chemother* 2005;49:2474-8.
56. Daisy P, Mathew S, Suveena S, Rayan NA. A novel terpenoid from *Elephantopus scaber* antibacterial activity on *Staphylococcus aureus*: A substantiate computational approach. *Int J Biomed Sci* 2008;4:196-203.
57. Brehm-Stecher BF, Johnson EA. Sensitization of *Staphylococcus aureus* and *Escherichia coli* to antibiotics by the sesquiterpenoids nerolidol, farnesol, bisabolol, and apritone. *Antimicrob Agents Chemother* 2003;47:3357-60.
58. González-Lamothe R, Mitchell G, Gattuso M, Diarra MS, Malouin F, Bouarab K. Plant antimicrobial agents and their effects on plant and human pathogens. *Int J Mol Sci* 2009;10:3400-19.
59. Kubo I, Muroi H, Himejima M. Antibacterial activity of totarol and its potentiation. *J Nat Prod* 1992;55:1436-40.
60. Horiuchi K, Shiota S, Hatano T, Yoshida T, Kuroda T, Tsuchiya T, *et al.* Antimicrobial activity of oleanolic acid from *Salvia officinalis* and related compounds on vancomycin-resistant enterococci (VRE). *Biol Pharm Bull* 2007;30:1147-9.
61. Delaquis PJ, Mazza G. Antimicrobial properties of isothiocyanates in food preservation. *Food Technol* 1995;49:73-84.
62. Luciano FB, Hosseinian FS, Beta T, Holley RA. Effect of free-SH containing compounds on allyl isothiocyanate antimicrobial activity against *Escherichia coli* O157:H7. *J Food Sci* 2008;73:M214-20.
63. Aires A, Mota VR, Saavedra MJ, Monteiro AA, Simões M, Rosa EA, *et al.* Initial *in vitro* evaluations of the antibacterial activities of glucosinolate enzymatic hydrolysis products against plant pathogenic bacteria. *J Appl Microbiol* 2009;106:2096-105.
64. Kim AB. Antimicrobial peptides: Pore formers or metabolic inhibitors in bacteria? *Nat Rev Microbiol* 2005;3:238-50.
65. Wiesner J, Vilcinskas A. Antimicrobial peptides: The ancient arm of the human immune system. *Virulence* 2010;1:440-64.
66. Barbosa Pelegrini P, Del Sarto RP, Silva ON, Franco OL, Grossi-de-Sa MF. Antibacterial peptides from plants: What they are and how they probably work. *Biochem Res Int* 2011;2011:250349.
67. Seo MD, Won HS, Kim JH, Mishig-Ochir T, Lee BJ. Antimicrobial peptides for therapeutic applications: A review. *Molecules* 2012;17:12276-86.
68. Gibbons S, Moser E, Kaatz GW. Catechin gallates inhibit multidrug resistance (MDR) in *Staphylococcus aureus*. *Planta Med* 2004;70:1240-2.
69. Yi ZB, Yan Y, Liang YZ, Bao Z. Evaluation of the antimicrobial mode of berberine by LC/ESI-MS combined with principal component analysis. *J Pharm Biomed Anal* 2007;44:301-4.
70. Iwasa K, Moriyasu M, Yamori T, Turuo T, Lee DU, Wiegrebe W, *et al.* *In vitro* cytotoxicity of the protoberberine-type alkaloids. *J Nat Prod* 2001;64:896-8.
71. Santiago C, Lim KH, Loh HS, Ting KN. Inhibitory effect of *Duabanga grandiflora* on MRSA biofilm formation via prevention of cell-surface attachment and PBP2a production. *Molecules* 2015;20:4473-82.
72. Kuźma Ł, Rózsalski M, Walencka E, Rózsalska B, Wysockińska H. Antimicrobial activity of diterpenoids from hairy roots of *Salvia sclarea* L.: Salvipisone as a potential anti-biofilm agent active against antibiotic resistant staphylococci. *Phytomedicine* 2007;14:31-5.
73. Huber BI, Eberl L, Feucht W, Polster J. Influence of polyphenols on bacterial biofilm formation and quorum-sensing. *Z Naturforsch C* 2003;58:879-84.
74. Negi PS. Plant extracts for the control of bacterial growth: Efficacy, stability and safety issues for food application. *Int J Food Microbiol* 2012;156:7-17.
75. Blanco AR, Sudano-Roccaro A, Spoto GC, Nostro A, Rusciano D. Epigallocatechin gallate inhibits biofilm formation by ocular staphylococcal isolates. *Antimicrob Agents Chemother* 2005;49:4339-43.
76. Soe WM, Giridharan M, Lin RT, Sakharkar MK. Effect of combinations of antibiotics and gallates on biofilm formation in *Staphylococcus aureus*. *Lett Drug Des Discov* 2010;7:160-4.
77. Cobrado L, Azevedo MM, Silva-Dias A, Ramos JP, Pina-Vaz C, Rodrigues AG, *et al.* Cerium, chitosan and *hamamelitannin* as novel biofilm inhibitors? *J Antimicrob Chemother* 2012;67:1159-62.
78. Cobrado L, Silva-Dias A, Azevedo MM, Pina-Vaz C, Rodrigues AG. *In vivo* antibiofilm effect of cerium, chitosan and *hamamelitannin* against usual agents of catheter-related bloodstream infections. *J Antimicrob Chemother* 2013;68:126-30.
79. Kiran MD, Giacometti A, Cirioni O, Balaban N. Suppression of biofilm related, device-associated infections by staphylococcal quorum sensing inhibitors. *Int J Artif Organs* 2008;31:761-70.
80. Packiavathy IA, Sasikumar P, Pandian SK, Veera Ravi A. Prevention of quorum-sensing-mediated biofilm development and virulence factors production in *Vibrio* spp. By curcumin. *Appl Microbiol Biotechnol* 2013;97:10177-87.
81. Packiavathy IA, Priya S, Pandian SK, Ravi AV. Inhibition of biofilm development of uropathogens by curcumin-an anti-quorum sensing agent from *curcuma longa*. *Food Chem* 2014;148:453-60.
82. Xie Q, Johnson BR, Wenckus CS, Fayad MI, Wu CD. Efficacy of berberine, an antimicrobial plant alkaloid, as an endodontic irrigant against a mixed-culture biofilm in an *in vitro* tooth model. *J Endod* 2012;38:1114-7.
83. Savoia D. Plant-derived antimicrobial compounds: Alternatives to antibiotics. *Future Microbiol* 2012;7:979-90.
84. Borges A, Ferreira C, Saavedra MJ, Simões M. Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria. *Microb Drug Resist* 2013;19:256-65.
85. Niu C, Gilbert ES. Colorimetric method for identifying plant essential oil components that affect biofilm formation and structure. *Appl Environ Microbiol* 2004;70:6951-6.
86. Niu C, Afre S, Gilbert ES. Subinhibitory concentrations of cinnamaldehyde interfere with quorum sensing. *Lett Appl Microbiol* 2006;43:489-94.

87. Brackman G, Defoirdt T, Miyamoto C, Bossier P, Van Calenbergh S, Nelis H, *et al.* Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. By decreasing the DNA-binding activity of the quorum sensing response regulator luxR. *BMC Microbiol* 2008;8:149.
88. Sendamangalam V, Choi OK, Kim D, Seo Y. The antibiofouling effect of polyphenols against *Streptococcus mutans*. *Biofouling* 2011;27:13-9.
89. Silver LL. Challenges of antibacterial discovery. *Clin Microbiol Rev* 2011;24:71-109.
90. Jayaraman P, Sakharkar KR, Daniel LC, Siddiqi MI, Dhillion SK, Sakharkar MK, *et al.* Hybrid-drug design targeting *Pseudomonas aeruginosa* dihydropteroate synthase and dihydrofolate reductase. *Front Biosci (Elite Ed)* 2013;5:864-82.
91. Newman DJ, Cragg GM, Snader KM. The influence of natural products upon drug discovery. *Nat Prod Rep* 2000;17:215-34.
92. Stavri M, Piddock LJ, Gibbons S. Bacterial efflux pump inhibitors from natural sources. *J Antimicrob Chemother* 2007;59:1247-60.
93. Nascimento GG, Locatelli J, Freitas PC, Silva GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic resistant bacteria. *Braz J Microbiol* 2000;31:247-56.
94. Kirtikar KR, Basu BD. *Indian Medicinal Plants*. 2<sup>nd</sup> ed., Vol. 3. Dehradun: International Book Distributors; 2005. p. 12.
95. Gupta AK, Neeraj T, Sharma M. *Quality Standards of Indian Medicinal Plants*. Vol. 3. New Delhi: ICMR; 2005. p. 9-19.
96. Yemane B, Berhane Y, Reddy KS. Ethnobotanical study of medicinal plants in sub region Logo Anseba, Region Gash Barka, Eritrea. *J Pharm Biol Sci* 2016;11:63-73.
97. Yemane B, Adebrihan M, Reddy KS. Traditional medicinal plants used BY tigrigna ethnic group in central region of Eritrea. *J Pharm Biol Sci* 2017;12:40-6.
98. Zapesochnaya GG, Kurkin VA, Okhanov VV, Perzykh LN, Miroshnilov AI. Structure of the alkaloids of *Aerva lanata*. *Chem Nat Compd* 1991a;27:725-8.
99. Zapesochnaya GG, Pervykh LN, Kurkin VA. A study of the herb *Aerva lanata*. alkaloids. *Chem Nat Compd* 1991b;27:336-40.
100. Zapesochnaya G, Kurkin V, Okhanov V, Miroshnikov A. Canthin-6-one and  $\beta$ -carboline alkaloids from *Aerva lanata*. *Planta Med* 1992;58:192-6.
101. Gupta AK, Tandon N. *Reviews of Indian Medicinal Plants*. Vol. 1. New Delhi: ICMR; 2004. p. 338-44.
102. Omoyeni OA, Adeyeye EI. Chemical composition, calcium, zinc and phytate interrelationships in *Aerva lanata* (L) Juss. ex schult leaves. *Orient J Chem* 2009;25:485-8.
103. Saleh NA, Mansour RM, Markham KR. An acylated isorhamnetin glycoside from *Aerva javanica*. *Phytochemistry* 1990;29:1344-5.
104. Pervykh LN, Karasartov BS, Zapesochnaya GG. A study of the herb *Aerva lanata* IV. Flavonoid glycosides. *Chem Nat Compd* 1992;28:509-10.
105. Ahmed E, Imran M, Malik A, Ashraf M. Antioxidant activity with flavonoidal constituents from *Aerva persica*. *Arch Pharm Res* 2006;29:343-7.
106. Yamunadevi M, Wesely EG, Johnson M. Chromatographic finger print analysis of steroids in *Aerva lanata* L by HPTLC technique. *Asian Pac J Trop Biomed* 2011;1:428-33.
107. Yamunadevi M, Wesely EG, Johnson M. Phytochemical studies on the Terpenoids of medicinally important plant *Aerva lanata* L. using HPTLC. *Asian Pac J Trop Biomed* 2011;1:220-5.
108. Yamunadevi M, Wesely EG, Johnson M. Chromatographic finger print studies on Saponins of *Aerva lanata* (L.) juss. Ex schultes by using HPTLC. *Int J Curr Pharm Res* 2012;4:52-7.
109. Yamunadevi M, Wesely EG, Johnson M. FTIR spectroscopic studies on *Aerva lanata* (L.) Juss. Ex schult. *Asian J Pharm Clin Res* 2012;5:82-6.
110. Manokaran S, Jaswanth A, Sengottuvelu S, Nandhakumar J, Duraisamy R, Karthikeyan D, *et al.* Hepatoprotective activity of *Aerva lanata* Linn. Against paracetamol Induced hepatotoxicity in rats. *Res J Pharm Tech* 2008;1:398-400.
111. Deshmukh T, Yadav BV, Badole SL, Bodhankar SL, Dhaneshwar SR. Antihyperglycaemic activity of alcoholic extract of *Aerva lanata* [L.] A. L. juss. Ex J. A. schultes leaves in alloxan induced diabetic mice. *J Appl Biomed* 2008;6:81-7.
112. Krishnan GA, Rai VK, Nandy BC, Meena KC, Dey S, Tyagi PK, *et al.* Hypoglycemic and antihyperlipidaemic effect of ethanolic extract of aerial parts of *Aerva lanata* Linn. In normal and alloxan induced diabetic rats. *Int J Pharm Sci Drug Res* 2009;1:191-4.
113. Anantha D, Kumar TI, Kumar MS, Reddy AM, Mukharjee NS, Rao AL. *In vitro* anti helmentic activity of aqueous and alcoholic extracts of *Aerva lanata* seeds and leaves. *J Pharm Sci Res* 2010;2:317-21.
114. Muthukumaran P, Shanmuganathan P, Malathi C. Antioxidative and Antimicrobial study of *Aerva lanata*. *Asian J Biochem Pharm Res* 2011;1:265-71.
115. Ragavendran P, Sophia D, Raj CA, Starlin T, Gopalakrishnan VK. Phytochemical screening, antioxidant activity of *Aerva lanata* (L)—an *in vitro* study. *Asian J Pharm Clin Res* 2012;5:77-81.
116. Arthi I, Subburaju T, Lakshminarayanan B, Rajkumar LA. *In vitro* evaluation of antioxidant activity of aqueous extract of *Aerva lanata* (L). *J Pharm Res* 2012;5:66-8.
117. Gujjeti RP, Mamidala E. Phytochemical screening and thin layer chromatographic studies of *Aerva lanata* root extract. *Int J Innov Res Sci Eng Technol* 2013;2:5725-30.
118. Chauhan VS, Swapna M. Isolation, characterization and cytotoxic activity of diterpenoid and flavonoids of *Aerva lanata*. *Int J Pharm Chem Sci* 2014;3:766-9.
119. Murugan M, Mohan VR. Phytochemical, FT-IR and antibacterial activity of whole plant extract of *Aerva lanata* (L.) Juss. Ex. Schult. *J Med Plants Stud* 2014;2:51-7.
120. Abinaya M, Gayathri VM, Khanna G. Evaluation of *in vitro* biological and ant biofilm activities of various leaf extracts of *Aerva lanata*. *Int J Pharm Sci Rev Res* 2016;38:65-72.
121. Lee H, Lee DG. Mode of action of bioactive

- phytochemicals, plant secondary metabolites, possessing antimicrobial properties. In: Méndez-Vilas A, editor. *The Battle Against Microbial Pathogens: Basic Science, Technological Advances and Educational Programs*. Spain: FORMATEX; 2015. p. 185-92.
122. Kotzekidou P, Giannakidis P, Boulamatsis A. Antimicrobial activity of some plant extracts and essential oils against food borne pathogens *in vitro* and on the fate of inoculated pathogens in chocolate. *LWT* 2008;4:119-27.
123. Singh G, Kumar P. Evaluation of antimicrobial activity of alkaloids of *Terminalia chebula* Retz. Against some multidrug-resistant microorganisms. *Int J Green Pharm* 2012;6:57-62.
124. Farn R. *Nature's Chemicals*. Oxford: Oxford University Press; 2010. p. 74-5.
125. Maurya R, Singh G, Yadav PP. Antiosteoporotic agents from natural sources. In: Atta-ur-Rahman FR, editor. *Studies in Natural Products Chemistry*. Vol. 35. Elsevier; 2008. p. 517-545.
126. Cazarolli LH, Zanatta L, Alberton EH, Figueiredo MS, Folador P, Damazio RG, *et al.* Flavonoids: Prospective drug candidates. *Mini Rev Med Chem* 2008;8:1429-40.
127. Friedman M, Henika PR, Levin CE, Mandrell RE, Kozukue N. Antimicrobial activities of tea catechins and theaflavins and tea extracts against *Bacillus cereus*. *J Food Prot* 2006;69:354-61.
128. Mirzoeva OK, Grishanin RN, Calder PC. Antimicrobial action of propolis and some of its components: The effects on growth, membrane potential and motility of bacteria. *Microbiol Res* 1997;152:239-46.
129. Soobrattee MA, Neergehen VS, Luximon-Ramma A, Aruoma OI, Bahorun T. Phenolics as potential antioxidant therapeutic agents: Mechanism and actions. *Mutat Res* 2005;579:200-13.
130. Ellnain-Wojtaszek M, Mirska I. Phenolic acids from *Ginkgo biloba* L. Part III. antimicrobial activity of phenolic acids. *Acta Pol Pharm* 1998;55:81-4.
131. Kubo I, Xiao P, Fujita K. Anti-MRSA activity of alkyl gallates. *Bioorg Med Chem Lett* 2002;12:113-6.
132. Kubo I, Fujita K, Nihei K. Molecular design of multifunctional antibacterial agents against methicillin resistant *Staphylococcus aureus* (MRSA). *Bioorg Med Chem* 2003;11:4255-62.
133. Taguri T, Tanaka T, Kouno I. Antibacterial spectrum of plant polyphenols and extracts depending upon hydroxyphenyl structure. *Biol Pharm Bull* 2006;29:2226-35.
134. Saavedra MJ, Borges A, Dias C, Aires A, Bennett RN, Rosa ES, *et al.* Antimicrobial activity of phenolics and glucosinolate hydrolysis products and their synergy with streptomycin against pathogenic bacteria. *Med Chem* 2010;6:174-83.
135. Martinez MJ, Lazaro RM, del Olmo LM, Benito PB. Anti-infectious activity in the *Anthemideae* tribe. In: Atta-ur-Rahman FR, editor. *Studies in Natural Products Chemistry*. Vol. 35. Elsevier; 2008. p. 445-516.
136. Gajalakshmi S, Vijayalakshmi S, Rajeswari VD. Pharmacological activities of *Aerva lanata*: A perspective review. *Int Res J Pharm* 2012;3:28-30.
137. Chowdhury D, Sayeed A, Islam A, Shah Alam Bhuiyan M, Astaq Mohal Khan GR. Antimicrobial activity and cytotoxicity of *Aerva lanata*. *Fitoterapia* 2002;73:92-4.
138. Kumar KN, Prabhu SN, Ravishankar B, Sahana, Yashovarma B. Chemical analysis and *in vitro* evaluation of antiurollthlatic activity of *Aerva lanata* (Linn.) juss. Ex schult. roots. *J Pharmacog Phytochem* 2015;3:1-7.
139. Yamuna DM, Wesely EG, Johnson M. Chromatographic studies on the tannins of *Aerva lanata* (L.) Juss. Ex schultes. *IOSR J Pharm* 2012;2:41-51.
140. Chung KT, Wong TY, Wei CI, Huang YW, Lin Y. Tannins and human health: A review. *Crit Rev Food Sci Nutr* 1998;38:421-64.
141. Chung KT, Lu Z, Chou MW. Mechanism of inhibition of tannic acid and related compounds on the growth of intestinal bacteria. *Food Chem Toxicol* 1998;36:1053-60.
142. Akiyama H, Fujii K, Yamasaki O, Oono T, Iwatsuki K. Antibacterial action of several tannins against *Staphylococcus aureus*. *J Antimicrob Chemother* 2001;48:487-91.
143. Haniffa MA, Kavitha K. Antibacterial activity of medical herbs against the fish pathogen *Aeromonas hydrophila*. *J Agric Technol* 2012;8:205-11.
144. Bossi A, Rinalducci S, Zolla L, Antonioli P, Righetti PG, Zapparoli G, *et al.* Effect of tannic acid on *Lactobacillus hilgardii* analysed by a proteomic approach. *J Appl Microbiol* 2007;102:787-95.
145. Engels C, Schieber A, Gänzle MG. Inhibitory spectra and modes of antimicrobial action of gallotannins from mango kernels (*Mangifera indica* L.). *Appl Environ Microbiol* 2011;77:2215-23.
146. Devarai HK, Koushik OS, Babu PS, Karthikeyan R. *In vitro* anti tubercular activity of leaves of *Aerva lanata* L. *Int Biol Biomed J* 2017;3:209-12.
147. Elangovan K, Sathasivam A, Rajasekaran R. Phytochemical screening and Antibacterial activity of *Aerva lanata* L. *Int J Sci Res* 2015;4:175-6.
148. Teixeira B, Marques A, Ramos C, Neng NR, Nogueira JM, Saraiva JA, *et al.* Chemical composition and antibacterial and antioxidant properties of commercial essential oils. *Ind Crops Prod* 2013;43:587-95.
149. Baltzer SA, Brown MH. Antimicrobial peptides: Promising alternatives to conventional antibiotics. *J Mol Microbiol Biotechnol* 2011;20:228-35.
150. Pushpanathan M, Gunasekaran P, Rajendhran J. Antimicrobial peptides: Versatile biological properties. *Int J Pept* 2013;2013:1-15.
151. Jenssen H, Hamill P, Hancock RE. Peptide antimicrobial agents. *Clin Microbiol Rev* 2006;19:491-511.
152. Broekaert WF, Cammue BP, De Bolle MF, Thevissen K, De Samblanx GW, Osborn RW, *et al.* Antimicrobial peptides from plants. *Crit Rev Plant Sci* 1997;16:297-323.
153. Stotz HU, Waller F, Wang K. In: Hiemstra PS, Zaat SA, editors. *Antimicrobial Peptides and Innate Immunity*. Basel: Springer Basel; 2013. p. 29-51.
154. Reddy KV, Yedery RD, Aranha C. Antimicrobial peptides: Premises and promises. *Int J Antimicrob*

- Agents 2004;24:536-47.
155. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: A common cause of persistent infections. *Science* 1999;284:1318-22.
156. Mah TF, O'Toole GA. Mechanisms of biofilm resistance to antimicrobial agents. *Trends Microbiol* 2001;9:34-9.
157. Anderson GG, O'Toole GA. Innate and induced resistance mechanisms of bacterial biofilms. *Curr Top Microbiol Immunol* 2008;322:85-105.
158. Daglia M, Tarsi R, Papetti A, Grisoli P, Dacarro C, Pruzzo C, *et al.* Antiadhesive effect of green and roasted coffee on *Streptococcus mutans*' adhesive properties on saliva-coated hydroxyapatite beads. *J Agric Food Chem* 2002;50:1225-9.
159. Xiao J, Zuo Y, Liu Y, Li J, Hao Y, Zhou X. Effects of *Nidus vespa* eextract and chemical fractions on glucosyltransferases, adherence and biofilm formation of *Streptococcus mutans*. *Arch Oral Biol* 2007;52:869-75.
160. Furiga A, Lonvaud-Funel A, Dorignac G, Badet C. *In vitro* anti-bacterial and anti-adherence effects of natural polyphenolic compounds on oral bacteria. *J Appl Microbiol* 2008;105:1470-6.
161. Quave CL, Estévez-Carmona M, Compadre CM, Hobby G, Hendrickson H, Beenken KE, *et al.* Ellagic acid derivatives from *Rubus ulmifolius* inhibit *Staphylococcus aureus* biofilm formation and improve response to antibiotics. *PLoS One* 2012;7:e28737.
162. Hancock V, Dahl M, Vejborg RM, Klemm P. Dietary plant components ellagic acid and tannic acid inhibit *Escherichia coli* biofilm formation. *J Med Microbiol* 2010;59:496-8.
163. Jagani S, Chelikani R, Kim DS. Effects of phenol and natural phenolic compounds on biofilm formation by *Pseudomonas aeruginosa*. *Biofouling* 2009;25:321-4.
164. Zhang J, Rui X, Wang L, Guan Y, Sun X, Dong M. Polyphenolic extract from *Rosa rugosa* tea inhibits bacterial quorum sensing and biofilm formation. *Food Control* 2014;42:125-31.
165. Borges A, Saavedra MJ, Simões M. The activity of ferulic and gallic acids in biofilm prevention and control of pathogenic bacteria. *Biofouling* 2012;28:755-67.
166. Ergün BC, Coban T, Onurdag FK, Banoglu E. Synthesis, antioxidant and antimicrobial evaluation of simple aromatic esters of ferulic acid. *Arch Pharm Res* 2011;34:1251-61.
167. Lemos M, Borges A, Teodósio J, Araújo P, Mergulhão F, Melo L, *et al.* The effects of ferulic and salicylic acids on *Bacillus cereus* and *Pseudomonas fluorescens* single and dual species biofilms. *Int Biodeterior Biodegradation* 2014;86:42-51.
168. Yang L, Rybtke MT, Jakobsen TH, Hentzer M, Bjarnsholt T, Givskov M, *et al.* Computer-aided identification of recognized drugs as *Pseudomonas aeruginosa* quorum-sensing inhibitors. *Antimicrob Agents Chemother* 2009;53:2432-43.
169. Schmidtchen A, Malmsten M. Peptide interactions with bacterial lipopolysaccharides. *Curr Opin Colloid Interface Sci* 2013;18:381-92.