



International Journal of Research in Pharmacology & Pharmacotherapeutics



ISSN Print: 2278-2648
ISSN Online: 2278-2656

IJRPP |Vol.4 | Issue 4 | Oct – Dec - 2015
Journal Home page: www.ijrpp.com

Research article

Open Access

Evaluation of alzheimer's disease of *mimusops elengi linn.* in the experimental model of rats

Bhukya Ramesh*¹, Santhosh Pawar V

*Department of Pharmacology, Nova College of Pharmaceutical Education and Research, Jaffergerud, Hayath Nagar, R.R Dist, Telangana

*Corresponding author: Bhukya Ramesh

E-mail id: ramesh.tocool@gmail.com

ABSTRACT

The present study has been undertaken to evaluate the possible role of *Mimusops elengi* Linn. flowers in experimental Alzheimer's disease (AD) in rats. Experimental AD in rats was produced by intra cerebrotal (i.c.v) administration of colchicine (Col). Various behavioural tests and biochemical analysis were performed to explore the possible role of the *Mimusops elengi* Linn. flower extract (ME)(100mg/kg & 200mg/kg doses) in AD. ME exhibited anxiolytic activity in Elevated plus maze test. In Morris' water maze test and Brightness discrimination test, ME pretreatments improved reference memory, working memory and spatial learning. ME significantly reduced the acetylcholinesterase. It reduced the Col induction increased lipid peroxidase activity, which was significantly reversed by ME (as seen from the reductions in the malondialdehyde level) and stabilized the rise in superoxide dismutase activity. ME might be effective in clinical AD by virtue of its cognition enhancement, anti-oxidant and anti-anxiety properties, which are the primary needs to be addressed in AD.

Keywords: Alzheimer's disease, Anxiolytic activity, Colchicine, Rats, *Mimusops elengi* Linn.

INTRODUCTION⁽¹⁻³⁾

The plant *mimusops elengi linn*(sapotaceae) commonly known as bakul, madhugandha, chirapushpa, pagademara is highly reputed in traditional medicine as stomachic, astringent, ulemorrhagia. Seeds contain Saponins. The seed kernels yield 16-25% of a fatty oil, used for edible and lighting purpose. The composition of the total fatty acids of the oil is as follows: Palmitic-10.97, Stearic-10.10, Behenic-0.46, Oleic-63.98, Linoleic-14.49%. Basic acid (C₃₀H₄₆O₅), the characteristic Saponin of *sapotaceae* has been isolated from the fat-free seed meal in a yield of 2.4%. A Saponin, which on hydrolysis yields rhamnose

(2mol.), arabinose(2mol.) and glucose (1mol.) has also been reported. The bark contains tannin, it is used in some parts of India for dyeing and tanning purposes. The bark and flowers are reported to contain a saponin and an alkaloid.

MATERIALS AND METHODS⁽⁴⁻¹⁰⁾

Collection and Authentication of Flowers

The flowers of *Minusops elengi* Linn. were collected from Nilgiri hills, Ooty, Tamilnadu and authentication (Voucher specimen number- PARC/2010/499) was done by Prof. P. Jayaraman, Ph.D. Plant Anatomy Research Centre, Medicinal plant Research Unit, Tambaram, Chennai-45.

Extraction of flowers of *mimusops elengi* linn

The collected flowers were cleaned, air dried at room temperature and ground to a coarse powder with an auto mix blender, passed through the sieve no:16 and stored in a deep freezer until the time for use. The powder was defatted with petroleum ether for 24 hours. Then, it was dried and cold macerated by using hydroalcoholic solvent(70% Ethanol and 30% Water) for about 5days^[45].The obtained extract(ME) was concentrated under reduced pressure and controlled temperature by Rotary evaporator at 40⁰c and stored in cool place

Experimental Animals

Thirty male Wistar rats, weighing 150-200 g were procured from King’s Institute, Guindy. The animals were maintained in the animal house under standard laboratory conditions with natural dark and light cycle (approximately 12 h light / 12 h dark cycle) and room temperature (27±1⁰C) and constant humidity (60%) in accordance with Institutional Ethical Committee rules and regulations. They were fed on a standard balanced diet and provided with water *ad libitum*. The project proposal was approved by Institutional Animal Ethical committee (IAEC 75/2009).

Table No.1 Experimental Animals

Group	No. of Animals	Treatment
I	6	Normal control(Distilled water, p.o)
II	6	Control-colchicine(15µg/rat)(I.C.V.R)
III	6	Colchicine(15µg)+ ME(100mg/kg/day, p.o)
IV	6	Colchicine (15µg)+ME(200 mg/kg/day, p.o)
V	6	Colchicine(15µg)+Donepezil(1mg/kg/day,p.o)

EXPERIMENTAL DESIGN ⁽¹¹⁻¹⁸⁾

Induction of Alzhiemer’s Disease with Colchicine

Colchicine will be administered via the intra cerebro ventricular route. The rats are anesthetised with Phenobarbital sodium and the right lateral ventricle will be cannulated and colchicine, dissolved in 5µl of artificial cerebrospinal fluid will be slowly injected into the cannulated ventricle using a 10µl Hamilton syringe. Control groups will

be subjected to the same surgical procedure and received only artificial cerebrospinal fluid.

RESULTS

Preliminary Phytochemical Investigation

The revealed results of the preliminary phytochemical screening of the hydroalcoholic extract of dried flowers of *Mimusops elengi* Linn. Results were shown below. Table no: 1. The extract gave positive results for alkaloids and saponins.

Table No: 2 Preliminary phytochemical test for ME

SL.No.	Phytochemical Tests	Results	SL. No.	Phytochemical Tests	Results
1	Test for Alkaloids	+Ve	7	Test for Flavonoids	-Ve
2	Test for Carbohydrates	-Ve	8	Test for Gums and mucilage	-Ve
3	Test for Proteins	-Ve	9	Test for Glycosides	-Ve
4	Test for Steroids	-Ve	10	Test for Saponins	+Ve
5	Test for Sterols	-Ve	11	Test for Terpenes	-Ve
6	Test for Phenols	-Ve			

*+Ve: indicates the presence of compounds *-Ve: indicates the absence of compounds

Table No: 3 Effect of ME on Elevated plus maze

Group	Time spent					
	Day 7		Day 14		Day 28	
	Open arm	Closed arm	Open arm	Closed arm	Open arm	Closed arm
I Normal	44.2±4.02**	193.12±23.18*	47.19±2.34*	192.00±23.19*	44.12±3.45*	193.23±34.56*

II Col-control	27.12±1.69	203.18±21.23	28.76±2.38	43.12±2.56	27.34±2.45	225±24.36
III ME100mg/kg	34.67±2.37*	187.32±21.45*	37.54±3.46*	198.13±1.98*	21.35±1.67*	213.78±32.13*
IV ME200mg/kg	41.23±4.38*	201.56±21.34*	44.34±2.48*	201.67±12.78**	23.56±3.89*	203.13±22.54*
V Std- Donepezil	16.38±2.06**	239.54±23.58*	15.48±4.37*	245.12±13.67**	17.45±1.63*	248.12±43.21*

Group	Number of entries					
	Day 7		Day 14		Day 28	
	Open arm	Closed arm	Open arm	Closed arm	Open arm	Closed arm
I Normal	7.13±0.65**	3.21±0.79**	6.12±0.88*	4.58±0.75**	8.06±1.39**	3.00±0.73**
II Col-control	5.12±0.60	6.98±0.78	5.76±0.73	0.63±0.56	3.31±0.76	7.12±0.06
III ME100mg/kg	8.87±0.45**	2.69±0.64**	7.98±0.76*	0.14±0.04*	0.09±0.01**	3.21±0.60**
IV ME 200mg/kg	10.37±0.97**	3.42±0.42**	9.00±0.73**	0.33±0.51*	0.17±0.70**	4.10±0.49**
V Std- Donepezil	4.12±0.51*	8.02±0.42*	3.17±0.44**	9.10±0.60**	32.74±0.60**	9.83±0.06**

Values are expressed as mean±SEM of 6 animals, *P<0.05, **P<0.01 vs Col-control (group II). Comparisons were made between Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std- Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test.

Table No: 4 Effect of ME on Radial Y-maze

Group	Acquisition					
	No. of trials			Latency period(sec)		
	Day7	Day14	Day28	Day7	Day14	Day28
I Normal	7.02±0.49*	7.78±0.01*	8.32±0.19**	100.43±2.875*	102.65±5.453*	103.50±3.890*
II Col-control	8.00±0.83	8.43±0.43	9.00±0.49	105.27±1.763	106.54±4.675	107.00±4.540
III ME (100mg/kg)	7.38±0.64*	7.45±0.51**	8.38±0.34*	137.98±14.29**	139.49±14.85**	142.87±16.68**
IV ME (200mg/kg)	6.69±0.38**	6.90±0.54**	7.19±0.98**	87.12±6.324**	88.59±6.596**	90.76±7.654**
V Std- Donepezil	7.52±0.85*	7.90±0.74*	8.02±0.85**	95.87±10.87**	96.12±12.87**	98.37±12.348*




Figure -1 After 28 days of treatment

Values are expressed as mean±SEM of 6 animals, *P<0.05, **P<0.01 vs Col-control (group II). Comparisons were made between Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std- Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test

Table No: 5 Effect of ME on Conditioned avoidance response

Group		Number of escape failures		
		Day 7	Day 14	Day 28
I	Normal	4.5±0.32**	3.1±0.23**	1.9±0.31**
II	Col-control	7.5±0.15	8.4±0.41	9.3±0.15
III	ME(100mg/kg)	6.5±0.71*	5.2±0.33*	4.1±0.84**
IV	ME(200mg/kg)	5.3±0.37**	4.1±0.23**	2.5±0.51**
V	Std-Donepezil	4.9±0.53**	3.6±0.62**	2.2±0.82**




Figure -2 After 28 days of treatment

Values are expressed as mean±SEM of 6 animals, *P<0.05, **P<0.01 vs Col-control(group II). Comparisons were made between Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std-Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test.

Table No: 6 Effect of ME on Water maze

Group	Day 7				14 th day					Day 28					
	E.L-1 Probe trial	E.L-2 New platform	E.L-3		E.L-1 Probe trial	E.L-2 New platform	E.L-3			E.L-1 probe trial	E.L-2 New platform	E.L-3			
I Normal	54.2 0±1.5*	29. 40±1.7*	14. 50±2.1*	32.7 0±1.5**	15. 20±1.2*	49.8 0±2.2*	29.2 0±1.7*	19.2 1±.5**	27.7 0±.3**	11.1 0±1.3*	51. 20±1.2*	31.6 7±.1*	18.2 1±.5*	22.4 0±.5*	13.4 2±.7*
II Col-control	58.4 0±2.1	44. 50±2.6	24. 67±1.2	15.4 2±1.7	23. 42±2.1	65.3 3±2.1	43.1 7±2.8	32.1 9±.3	13.2 5±.9	26.7 0±2.8	68. 20±1.2	48.3 7±.4	38.7 7±.4	33.4 0±.9	27.2 2±.4
III ME1	42.3 3±0.7**	18. 15±1.2*	10. 20±1.1*	20.4 0±1.6*	10. 15±1.7*	29.1 2±1.4*	9.77 ±1.2**	9.85 ±0.6*	13.1 5±.6*	8.20 ±1.2*	24. 20±0.8*	12.7 1±.6*	9.40 ±1.2*	13.7 7±.9*	9.20 ±1.8*
IV ME2	59.7 0±1.2*	27. 33±1.2*	16. 40±1.7*	31.1 7±1.8*	16. 33±1.8*	54.2 0±1.3*	22.3 3±0.9*	17.8 3±.4**	26.7 7±.7*	15.3 3±0.8*	52. 20±1.2*	26.1 7±.4*	21.3 3±.2*	25.1 7±.5*	14.2 2±.6*
V Col-Doz	56.3 3±2.1**	24. 22±1.4*	18. 40±2.6*	29.1 7±1.3**	12. 35±2.3*	55.2 2±2.1*	26.1 7±1.8*	25.4 0±.6**	34.1 7±.7*	15.8 5±.5*	58. 17±1.4*	25.7 7±.1*	28.4 0±.3*	24.2 1±.5*	17.2 8±.1*




Figure -3 After 28 days of treatment

Values are expressed as mean±SEM of 6 animals, *P<0.05 **P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std- Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dennett’s test.

Table No: 7 Effect of ME on acetyl cholinesterase (AChE) (µ moles/min/mg protein)

Group	Reaction time(sec)			Relearning Index %		
	Day 7	Day 14	Day28	Day 7	Day14	Day28
I Normal	6.8±0.8**	7.3±0.6**	8.8±0.4**	58.0±3.0**	70.0±5.0**	83.0±7.0**
II Col- Control	3.2±0.6	3.8±0.3	3.9±0.3	32.0±2.0	36.0±4.0	37.0±1.0
III ME 100mg/kg	4.3±0.2*	4.7±0.1*	4.8±0.4*	42.0±3.0**	41.0±4.0*	39.0±5.0*
IV ME200mg/kg	5.7±0.1**	7.2±0.4**	8.0±0.1**	51.0±4.0**	48.0±4.0**	46.0±1.0**
V Std- Donepezil	7.8±0.3**	9.6±0.6**	11.2±1.0**	56.0±5.0**	53.0±5.0**	51.0±1.0**




Figure -4 After 28 days of treatment

Values are expressed as mean±SEM of 6 animals *P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std- Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett’s test

Table No: 8 Effect of ME on acetyl cholinesterase (AChE) (µ moles/min/mg protein)

Group	7 th day	14 th day	28 th day
I Normal	35.13±3.65**	36.13±4.22**	40.09±3.12**
II Col-control	10.01±2.35	12.56±3.65	13.96±4.59
III ME(100mg/kg)	21.67±4.56**	26.34±3.89**	21.76±4.87**
IV ME (200mg/kg)	23.5±3.54**	29.38±5.43**	22.87±6.45**
V Std- Donepezil	29.44±4.21**	30.09±7.12**	27.13±5.34**

Values are expressed as mean±SEM of 6 animals *P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std-Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test.




Figure -5 After 28 days of treatment

Table No: 9 Effect of ME on Lipid peroxidation (LPO) in Brain tissue (units /min /mg protein)

Group	7 th day	14 th day	28 th day
I Normal	2.50±0.15**	2.1±0.301**	2.4±0.32**
II Col-control	5.3±0.36	5.7±0.52	5.56±0.22
III ME (100mg/kg)	4.2±0.09*	4.5±0.12*	4.33±0.21*
IV ME (200mg/kg)	3.0±0.15*	3.2±0.22**	3.18±0.31**
V Std-Donepezil	2.8±0.35**	2.5±0.18**	2.64±0.22**




Figure -6 After 28 days of treatment

Table No: 10 Effect of ME on Superoxide dismutase (SOD) (units/min/mg protein)

Group	7 th day	14 th day	28 th day
I Normal	7.39±0.03**	7.87±0.01**	8.08±0.53**
II Col-control	4.39±0.01	4.98±0.05	5.36±0.09
III ME (100mg/kg)	5.18±0.01*	5.91±0.02*	6.03±0.01*
IV ME (200mg/kg)	6.72±0.04*	6.88±0.04**	7.96±0.43*
V Std-Donepezil	7.87±0.05**	7.93±0.06**	8.36±0.07**

Values are expressed as mean±SEM of 6 animals, *P<0.05, **P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std-Donepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test.




Figure -7 After 28 days of treatment

Table No: 11 Effect of ME on Glutathione peroxidase (GPx) (units/min/mg protein)

Groups		7 th day	14 th day	28 th day
I	Normal	34.37±1.63**	34.31±1.06**	35.15±1.01**
II	Col-control	21.81±1.28	21.11±1.34	22.21±1.07
III	ME (100mg/kg)	25.31±1.31*	25.83±1.03*	26.13±0.63*
IV	ME (200mg/kg)	30.81±1.06*	31.91±1.51*	32.31±1.08**
V	Std-Nonepezil	32.66±1.91**	33.58±1.73**	34.31±1.09**

Values are expressed as mean±SEM of 6 animals, *P<0.05 **P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME (200

mg/kg) and Std-Nonepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dennett's test.




Figure -8 After 28 days of treatment

Table No: 11 Effect of ME on Reduced Glutathione (units/min/mg protein)

Group		7 th day	14 th day	28 th day
I	Normal	7.40±0.51**	7.03±0.63**	8.05±0.41**
II	Col-control	4.96±0.49	4.87±0.37	5.01±0.78
III	ME (100mg/kg)	6.51±0.53*	5.36±0.45*	6.03±0.03*
IV	ME (200mg/kg)	5.01±0.31*	6.82±0.26**	7.63±0.36*
V	Std-Nonepezil	7.07±0.06**	7.08±0.04**	8.77±0.73**

Values are expressed as mean±SEM of 6 animals, *P<0.05, **P<0.01 vs Col-control (group II). Comparisons were made between: Col-control group with Normal, ME (100mg/kg), ME

(200mg/kg) and Std-Nonepezil groups. Symbol represents the statistical significance done by ANOVA, followed by Dunnett's test.




Figure -9 After 28 days of treatment

SUMMARY

The preliminary phytochemical screening of ME shows the presence of various phytochemical constituents like alkaloids, saponins, carbohydrates and tannins. The neuro protective effect of ME was assessed by Elevated plus maze, Y-maze, Conditioned avoidance response, Water maze on which it showed considerable attentive effect. In elevated plus maze test, ME significantly reversed the decrease in open arm to closed arm ration induced by colchicines indicating the anxiolytic activity. In Y-Maze test, after the injection of ME, the spontaneous alteration percentage was found to be improved in 200mg/kg treated group. ME administration improves the memory deficits in the active avoidance task as it reverses the increased escape latencies with colchicines. In Water maze test, the impaired learning by colchicines and the improvement of learning by ME after injection shows the significant property of memory retention which indicates the rejuvenation potential of the extract. The time required to escape on the platform is decreased on this task indicating the hippocampal learning ability of the extract. ME improves the memory and learning of animals in Brightness discrimination test by increasing reference time and re-learning index values. The biochemical changes responsible for the cognitive impairment were assessed by the estimation of acetylcholine esterase and antioxidant enzymes. The study against the colchicines induced alzheimer's by the treatment of ME shows significant reduction in the activity of acetylcholine esterase. The antioxidant value of ME shows the regaining of the antioxidant enzymes SOD, GSH and GPx activity, it has been noted that the antioxidant properties of extract delays the generation of free radical and also showed the

reversal of the decreased antioxidant enzyme levels after the memory impairment. There was a decrease in the Lipid peroxidation levels which were alleviated by colchicine treatment.

CONCLUSION

Mimusops elengi Linn is being used in traditional medicine as a antioxidant of CNS associated disorders, still there are some scientific evolutions to be made. Hence this study is emphasized to make the evident effect of the whole plant on memory disorder representing Alzhiemer's disease. The investigation was carried out on cognitive impairment due to colchicine induced impaired behavioural performace and oxidative stress. In Elevated plus maze test and Y-maze test, ME at both 100mg/kg and 200mg/kg exhibited significant improvement than the colchicine group of animals. In conditioned avoidance response task and in Brightness discrimination test, ME at both doses indicated the improvement of memory and learning. The spatial learning in Water maze task showed the significant memory retention indicated by the decrease in escape latency at both dose levels. ME treatment had shown the significant reduction in the elevated enzyme levels of acetylcholine esterase which indicates the potential to increase cognitive function through the decreased degradation of acetyl choline. The antioxidant levels were also proved to be restored on ME treatment as there was an increase in SOD, GSH and GPx levels by decreasing the LPO level. In conclusion, the neuroprotective activity of the flowers of the plant *Mimusops elengi Linn*. On alzheimer's type of dementia may be due to the inhibiting activity against AChE, free radical scavenging activity and they can be expected to be a pivot sense in neurotoxiciry.

BIBLIOGRAPHY

- [1]. Chittaranjan Andrade, J.Suresh Chaadra. Anti-amnesic properties of Brahmi and Mandookaparni in a rat model. Indian Journal of Psychiatry. 2006; 48: 232-237.
- [2]. Sharma A, Parikh V, Singh M. Pharmacological basis of drug therapy of Alzheimer's disease. Indian Journal of Experimental Biology; 35(199): 1146-115.
- [3]. Bennett, Plum. Cecil Text book of Medicine. 20th ed: Prism India Pvt Ltd, 1996; p. 2047.
- [4]. "Alzheimer's diagnosis of AD". Alzheimer's Research Trust. <http://www.alzheimers-research.org.uk/info/diagnosis>. Retrieved 2008-02-29.
- [5]. Landes AM, Sperry SD, Strauss ME, Geldmacher DS (Dec 2001). "Apathy in Alzheimer's disease". Journal of American Geriatrics; 49 (12): 1700-7.
- [6]. Volicer L, Harper DG, Manning BC, Goldstein R, Satlin A (May 2001). "Sundowning and circadian rhythms in Alzheimer's disease". American Journal of Psychiatry; 158 (5): 704-11.
- [7]. Holmes C, Boche D, Wilkinson D, et al. (July 2008). "Long-term effects of Abeta42 immunisation in Alzheimer's disease: follow-up of a randomised, placebo-controlled phase I trial"; Lancet 372 (9634): 216-23.
- [8]. Shen ZX (2004). "Brain cholinesterases: II. The molecular and cellular basis of Alzheimer's disease". Medical Hypotheses; 63 (2): 308-21.
- [9]. M.Raghavendra, Rituparna Maiti, Shafalika Kumar, S.B. Acharya. Role of *Ocimum sanctum* in the experimental model of Alzheimer's disease in rats. Journal of Green pharmacy 2009; 6-15.
- [10]. Moan R (July 20, 2009). "MRI software accurately IDs preclinical Alzheimer's disease".
- [11]. Vaidyaratnam P.S. Varier. Indian Medicinal Plants (1)-A Compendium of 500 species. Orient Longman Pvt Ltd, 1988; p.361- 365.
- [12]. The Wealth of India (L-M), Vol: 6: Publications & Information Directorate, CSIR, 1995; p.205.
- [13]. Chopra R.N, Nayar S.L, Chopra I.C. Glossary of Indian Medicinal Plants. National Institute of Science Communications, 1956; p.167.
- [14]. The useful plants of India. Publications & Information Directorate, CSIR, 1992 ;p. 375.
- [15]. N. Raaman. Phytochemical Techniques. New India Publishing Agency, 2006; p.10, 19-24.
- [16]. Drug Discovery and Evaluation-Pharmacological assays, H.Gerhard Vogel. 2nd Ed: p.428-439.
- [17]. Pawel Boguszewski, Jolanta Zagrodska. Emotional changes related to age in rats-a behavioural analysis. Behavioural brain research 2002;133:323-332.
- [18]. Nirmal Sethi, Sanjay Dube, H.K. Singh. Effect of Chronic Administration of lithium on Memory functions in Rats. Indian Journal of Psychiatry 1983;25(2):102-106.