

TANTIA UNIVERSITY JOURNAL OF HOMOEOPATHY AND MEDICAL SCIENCE Case Report

Volume 2 Issue 2(April-June 2019)

EISSN 2581-8899

E-ISSN: 2581-8899

Homoeopathic Management of Peptic Ulcer- A case report

Dr. Rajinder Girdhar

H.O.D. Dept. of Anatomy, Sriganganagar Homoeopathic Medical College Hospital and Reserch Institute, Sriganganagar Abstract- Ulcerations in the lining of upper part of the digestive tract are known as Peptic Ulcers. The ulcers may exist in the lower part of food pipe (oesophagus), in the stomach or in the initial part of the intestine (duodenum). About 10% of all adults are affected with Peptic ulcers at some time in their life. One of the most common problems of the gastrointestinal system. The damage occurs on the lining which may be due to inflammation caused by certain bacteria, or due to erosion caused by the stomach acids. Peptic ulcer disease is a constitutional disease that finds its local expression at the level of the digestive system. Homoeopathy is holistic system of medicine that treat person as whole here a case report was a 35 yrs old female suffering from Peptic ulcer, on basis on totality of symptoms & individual peculiarity, prescribe homeopathic medicine Nux Vomica 30 & case was cured and quality of life became better.

Key word- Gastric, Duodanal, Ulcer, H. Pylori, Haematemesis

Corresponding author- Dr. Rajinder Girdhar (Prof.) H.O.D. Anatomy Dept., Sriganganagar Homoeopathic Medical College Hospital and Reserch Institute, Sriganganagar, Rajasthan

How to cite- Girdhar R., Homoeopathic Management of Peptic Ulcer - A case Report, TU J. Homo & Medi Sci. 2019; 2(2):49-52

Introduction-

peptic ulcers occur when the lining of the stomach - the esophagus or the small intestines, gets damaged. Peptic ulcers can cause intense abdominal pain, vomiting nausea, indigestion and sudden unexplained weight loss.


Causes of Peptic Ulcer:

The underlying process that results in Peptic ulcer formation is excess of acid production in the stomach and damage to the protective barrier inside the stomach. Certain things that can trigger excess acid production or can cause damage to the protective lining of the stomach are:

- Emotional stress
- Foods (spices, pungent foods, etc)
- Overuse of certain drugs, especially NSAIDs (Non-steroidal antiinflammatory drugs)
- Infections (H-Pylori)
- Hereditary
- Smoking

Types of Peptic Ulcers:

It may be Gastric ulcer or Duodenal ulcer. Peptic ulcers usually run a chronic course and complaints tend to be episodic in nature; the patient is usually free from the symptoms in between these episodes.



Common symptoms of Peptic ulcers are:

• Pain in the epigastrium (upper part of abdomen)

E-ISSN: 2581-8899

- Hunger pain- worsening of pain when the stomach is empty
- Night pain- wakes the patient from sleep
- Heartburn
- Water brash
- Loss of appetite (anorexia)
- Nausea
- Haematemesis (vomiting of blood)
- Dark or black stools

Difference between Gastric Ulcer and Duodenal Ulcer -

| Chronic Gastric Ulcer | Duodenal Ulcer |
|---|--|
| 1. Patients are usually middle age | 1. Usually young adults |
| 2. More common in males. | 2. Common In both sexes. |
| 3. Patients look thin. | 3. Patients look obese. |
| 4. Periodicity is less marked. | 4. Periodicity is well marked. |
| 5. Pain in mid epigastrium slightly. | 5. Pain in transpyloric plane To left. |
| 6. Taking food aggravates the pain. | 6. Taking food relieves the pain. |
| 7. Pain is not felt at night, just after eating | 7. Pain felt at night, after 3 hr after meal |
| 8. Vomiting is common found after food | 8. Vomiting is rare unless pyloricstenosis complicates |
| 9. Patient avoids spicy food. | 9. Any food doesn't initiate pain so He doesn't avoid. |
| 10. Patient looses weight | 10 .Patient gains weight |
| 11. Haematemesis is common | 11. Melaena is more common |

Prevention of Peptic ulcer- Certain lifestyle choices and habits can reduce your risk of developing peptic ulcers. These include:

Not drinking more than two alcoholic beverages a day

Not mixing alcohol with medication

Washing your hands frequently to avoid infections

Limiting your use of ibuprofen, aspirin, and naproxen (NSAID)

Maintaining a healthy lifestyle by quitting smoking cigarettes and other tobacco use and eating a balanced diet rich in fruits, vegetables, and whole grains will help you prevent developing a peptic ulcer.

Case Profile -

Name - Mrs. R. K.

Age - 36 year Sex - Female

Marital Status - Married

Socio-economic Status - Middle

Religion - Sikh

Present Complaints

- She complained of continuous sour eructation every day since the last 3 years.
- It would increase if she missed her meal timings or by eating spicy food, eggs, or sour food.
- She was taking an antacid daily.
- She also complained of back pain since last 2 years.
- Nausea and vomiting

History of Present Complaints

Patient come with complaint of sour eructation and pain at night since last three yrs. Patient feels burning sensation in epigastrium. She is cheerful in nature becomes angry at trifles. She was working in MNC. She lived with her in laws and children.

Past History of Illness with Treatment

- History of constipation.
- History of Appendicitis taking allopathic treatment.

Family History

- Father- MI
- Mother- DM II
- Grand Father- BPH

Patient as a Person (Personal History) General Appearance

a) Built - Normalb) Gait - Normal

c) Speech - Normal

d) Dress Sense - Good

e) Height - 5'3"

f) Weight - 57 kg.

Diet

a) Appetite - Normalb) Thirst - Normal

c) Desires - Spicy food

E-ISSN: 2581-8899

Alcohal

d) Aversion - Nothing

Discharges

Addiction

a) Stool - Constipationb) Urine - 4-5/0-1 D/N

c) Abnormal Discharges - NAD

Thermal - Chilly

Sleep -Sleepy but can't sleep

Mental Generals

- She becomes abusive during anger.
- Anger at trifles.
- Anger-violent.
- Irritability
- Fault finder

Gynaecological/Obstetrical History

Menses: LMP-20.08.18

• Time - 3-4 days, regular (28-30 days)

• Character of blood - Bright red

• Character of pain – Only 1st day

• Obs. H/O- G₂P₂A₀L₂ FTND

Physical Examination:

• Pulse - 68/min

• Blood Pressure - 120/80 mm of hg

Anemia - Absent

Eyes - Normal

• Tongue - Clean

Systemic Examination:

• GIT - Distended

• URINARY - NAD

CNS - Well Oriented

• CVS - S1, S2 Heard

• Respiratory - Bilateral Lung clear

Investigation - USG Abdomen – Duodenal ulcer

E-ISSN: 2581-8899

Provisional Diagnosis & D/D:

Gastric Ulcer

Duodenal Ulcer

Final Diagnosis - Duodenal Ulcer

Repertorial Analysis

A) Repertorial Totality with Evaluation Of Symptoms

| S. | Symptoms | Analysis |
|-----|---------------------|----------|
| No. | | |
| 1. | Anger at trifle | MG |
| 2. | Irritability | MG |
| 3. | Fault finder | MG |
| 4. | Chilly | PG |
| 5. | Desire- Spicy thing | PG |
| 6. | Pain at Epigastrium | PP |
| 7. | Nausea | PP |
| 8. | Heart Burn | PP |
| 9. | Pain after Eating, | PP |
| | Midnight | |

| 10. | Sour taste of mouth | PP |
|-----|---------------------|----|
| 11. | Pain Back | PP |

B. Analysis With Comments

Out of six highest grading medicines. Nux vomica is covering all the symptoms including thermal also. So Nux Vomica becomes the ultimate choice.

Final Prescription -

Rx

Nux Vomica 200/3 dose PL 30 /BD 4 pills for 7 days

- Oral

Mode of administration

Date - 28.08.2018

General Management

Advice to take medicine regularly. Take blond diet, avoid spicy and outside food.

Pt. Name -Mrs. R. K. 36/F Date- 28/08/2018

| 1. Clipboard 1 | × |
|--|-----------|
| 1. Mind - ANGER, general - trifles, at | (59) 1 |
| 2. Mind - IRRITABILITY, general - pai | . (21) 1 |
| Mind - FAULTFINDING, disposition | (40) 1 |
| Generals - COLD, temperature, w | (216) 1 |
| Food - SPICY, general - desires | $(75)\ 1$ |
| 6. Abdomen - PAIN, abdomen - midni, | . (14) 1 |
| Stomach - HEARTBURN, general | (233) 1 |
| 8. Stomach - NAUSEA, general - mc | (147) 1 |
| Stomach - PAIN, stomach - eating | (143) 1 |
| 10. Taste - SOUR, taste - morning | (25) 1 |
| 11. Back - PAIN, back - sitting, pain w | . (95) 1 |
| | |

| (N) | S.M. | £. & | . S. | , \$c. | 14c | . 4sz | 'n, |
|-------------|----------|---------|---------|---------|---------|---------|-----|
| 1 | 2 | 3 | 4 | 5 | 6 | 7 | |
| 11 29 | 10 20 | 9 21 | 9 18 | 9 18 | 9 17 | 9 15 | 1 |
| 4 | _ | 1 | 1 | 1 | 1 | 2 | : |
| 2 | 1 | - | 1 | - | - | - | |
| 3 | 2 | 3 | 2 | - | 1 | 2 | : |
| 3 | 2 | 3 | 3 | 3 | 3 | 2 | : |
| 2 | 3 | 1 | 3 | 3 | 1 | 1 | : |
| 1 | 1 | - | 2 | 1 | - | - | |
| 3 | 2 | 2 | 2 | 3 | 3 | 2 | : |
| 3 3 3 | 2 | 3 | - | 1 | 1 | 2 | |
| 3 | 3 | 3 | 3 | 3 | 3 | 2 | : |
| 3 | 2 | 2 | - | 1 | 2 | 1 | |
| 2 | 2 | 3 | 1 | 2 | 2 | 1 | |
| | | | | | | | |

Treatment & Follow-Up:

| Date | Follow Up | Prescription |
|------------|---|---------------------------------|
| 28-08-2018 | As per repertorization | Nux Vomica 200/3 Dose, |
| | | PL 30- 4globules/BD for 7 Days |
| 6-09-2018 | Sour eructation and sour taste of mouth | Nux Vomica 200/1 Dose, |
| | not present, pain intensity reduce | PL 30- 4globules/BD for 15 Days |
| 22-09-2018 | LMP 21-09-2018, No Nausea and | PL 30- 4globules/BD for 15 Days |
| | vomiting, pain reduce 50%, Back ache | |
| | better | |

| 10-10-2018 | Anger reducing. | PL 30- 4globules/BD for 15 Days |
|------------|---|---------------------------------|
| | No heart burn, Back ache better, no pain | |
| | after eating, Stool normal in consistency | |
| 26-10-2018 | LMP 19-10-2018 | Sulphur 30/1 Dose, |
| | No change, case progress stop | PL 30-4globules/BD for 15 Days |
| | Sulphur is complementary of Nux | |
| | Vomica | |
| 12-11-2018 | All complaints are better. | PL 30- 4globules/BD for 15 Days |
| | | |
| 29-11-2018 | LMP 20-11-2018 | PL 30- 4globules/BD for 1 Month |
| | No complaint | |

Conclusion - Homeopathic approach towards the management of Peptic ulcer disease involves the evaluation of the complaint in details while taking into account individual factors such as one's personal and family history while planning a long-term treatment. Since the treatment targets the root cause, Homeopathy offers long-term cure instead of temporary relief. It treats at a deeper level of immunity and enhances the healing capacity of the body. This helps not only cure the ulcers but also the pre disposition to developing it again. This case shows that homeopathy is effective in case of Duodenal Ulcer, if we prescribe the medicine on the basis of totality of symptoms and individualization. If this study carried out on big sample size then result show documentary evidence of effectiveness of homoeopathy.

References

- 1. API Text Book Of Medicine Of Edition 2012 Published by The association of physician of India Mumbai 400011.
- 2. Harrison's Principles Of Internal Medicine Vol.1, 18th Edition by Antony S.Fauci,Eugene Braunwald,Dennis L.Kasper, Stephen L. Hauser,Dan L. Longo,J. Larry Jameson, Joseph Loscalzo.
- 3. Boericke, William. Pocket Manual of Homoeopathic Materia Medica and Repertory comprising of the Characteristic and guiding symptoms of all Remedies (clinical and pathogenetic) including Indian Drugs. B. Jain Publishers Pvt. Ltd, New Delhi, 2007.

Conflict of Interest: None

4. Allen H.C. Allen's Keynotes, Rearranged and Classified with leading remedies of the Materia Medica & Bowel Nosodes, 9th edition, B. Jain Publisher Pvt. Ltd., New Delhi, Reprint edition 2004

E-ISSN: 2581-8899

- Hahnemann Samuel. Organon of medicine.
 Translated by William Boericke. B. Jain
 Publishers Pvt. Ltd., New Delhi, Reprint
 Edition 2002.
- 6. Kent James Tyler. Lectures on Homoeopathic Philosophy. Memorial Edition Reprint. B. Jain Publishers Pvt. Ltd., New Delhi, 2004.
- 7. Archibel Medical Software. Radar 7

Source of Support: Nil

This work is licensed under a Creative Commons Attribution 4.0 International License