



## International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR)

ISSN:2347-6567

IJAMSCR |Volume 4 | Issue 4| Oct -Dec - 2016  
www.ijamscr.com

Research article

Medical research

### Prevalence of anemia in type 2 diabetics-an overview

Prabakaran Thamilmani<sup>\*1</sup>, Kannan Rajendran<sup>2</sup>, Rajendran K<sup>3</sup>

<sup>1</sup>Resident, Department of Internal Medicine, Saveetha Medical College & Hospital Chennai, Tamilnadu, India,

<sup>2</sup>Professor, Department of Internal Medicine, Saveetha Medical College & Hospital, Chennai, Tamilnadu, India,

<sup>3</sup>Professor, Department of Internal Medicine, Saveetha Medical College & Hospital Chennai, Tamilnadu, India.

\*Corresponding Author: Prabakaran Thamilmani

Email: ptpcardio@gmail.com

#### ABSTRACT

##### Aim

Anemia is found to contribute to the development and progression of micro and macro-vascular complications of diabetes. This study aimed to determine the prevalence of anemia in patients with diabetes and the occurrence of micro and macro vascular complications in them.

##### Methods

This was a retrospective study of 227 patients with diabetes regularly followed up as outpatients for one year. Patient's demographic profile, history of stroke, ischemic heart disease (IHD), and concomitant hypertension (HT) along with the most recent values of complete blood picture and renal function tests were collected from the database of the Hospital.

##### Results

The prevalence of anemia in our study population is 55.5%, which is two to three times more common as compared to the result from previous studies. Anemia was six times more common (53.9%) compared to the previous study for the same GFR value (GFR > 60 mL/min/1.73 m<sup>2</sup>) and half of them in stage one kidney disease.

##### Conclusion

Anemia is a common accompaniment with diabetes and it is seen early even in the absence of renal impairment which necessitate early screening of anemia and further studies to know the possible etiology.

**Keywords:** Chronic kidney disease, Hypertension, Stroke, Heart disease.

## INTRODUCTION

Diabetes is a major public health problem worldwide. Globally it is estimated that 382 million people suffer from diabetes with a prevalence of 8.3%. India with a prevalence of 8.7% is one of the top 10 countries with higher prevalence of diabetes [1]. Several studies suggest that anemia is twice as common in diabetics compared with non-diabetics [2]. Despite these facts, anemia is unrecognized in 25% of the diabetic patients [3]. Anemia also develops earlier and is more severe in patients with diabetes than in patients with renal impairment from other causes [4]. Additionally, the risk of anemia is higher in people with diabetic nephropathy compared with people with nephropathy from other causes and is associated with a more rapid decline in the GFR [5, 6]. However, it develops whilst

Serum creatinine is within the normal range in 5% of the cases [5]. Recent studies have linked anemia with relatively low serum erythropoietin in persons with either type 1 or type 2 diabetes, even without advanced kidney disease or overt uremia [7]. The etiology of anemia in diabetes is multifactorial and includes inflammation, nutritional deficiencies, concomitant autoimmune diseases, drugs, and hormonal changes in addition to kidney disease [3]. Anemia is found to contribute to the development and progression of micro- and macro-vascular complications of diabetes, which has a negative impact on the quality of life and an additional burden on the health of the patients [4,8]. It is therefore important to diagnose and correct anemia among diabetic patients early. This study aimed to determine the prevalence of anemia in diabetics and the occurrence of micro and macro vascular complications in them.

## MATERIALS AND METHODS

This was a retrospective study of diabetic patients regularly followed up as outpatients at hospital. Data was collected with the approval of Health center's ethics committee.

### Subjects

All diabetic patients attending the Out Patient Department between 1st January 2015 and 31st

December 2015 were included in the study evaluation.

### Determination of variables

Patient's demographic profile, history of stroke, ischemic heart disease (IHD), and concomitant hypertension (HT) along with complete blood picture and renal function tests were collected from the database of the hospital. The most recent values were used for the analysis. Estimated glomerular filtration rate (GFR) using Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) equation formula was used to define renal failure.

### Definition of diabetes

Diabetes was diagnosed, when the fasting glucose value was  $>125$  mg/dL, or random blood glucose  $>200$  mg/dL or patients on treatment for diabetes. The diagnosis of diabetes was based on the "Definition and description of diabetes mellitus" from American Diabetes Association 2010 [9].

### Anemia

Was considered as per the World Health Organization's gender specific criteria, ( $<13$  g/dL in men and  $<12$  g/dL in women) [10]. Anemia was defined as normocytic with a mean corpuscular volume (MCV) of 80 to 100 FL, microcytic with the MCV  $<80$  FL, and macrocytic with the MCV  $>100$  FL [11]

### Renal failure

Patients were classified based on estimated glomerular filtration rate (EGFR). EGFR values  $<90$  mL/min/1.73 m<sup>2</sup> were considered abnormal (CKD 2–5). Staging of CKD was based on the Kidney Disease Outcomes Quality Initiative (K/DOQI) guidelines:  $\geq 90$  (CKD1), 60–89 (CKD2), 30–59 (CKD3), 15–29 (CKD4) and  $<15$  (CKD5) ml / min 1.73 m<sup>2</sup>. Creatinine clearance (CCr) was determined using the (CKD-EPI) equation formula to define renal failure [12-14]

### Statistical analysis

Descriptive analysis were used to characterize the study sample by demographics, including sex, age, prevalence of hypertension, stroke, heart disease in relation to anemia and kidney function. The result was presented in the form of tables, using frequencies percentage, mean and standard

deviation to describe the study sample in relation to relevant variables. P-value less than 0.05 was taken as statistical significant. Chi-square statistics was used for testing the association between the categorical variables. All the data were manually checked for its clarity and completeness, then coded, entered and transported to IBM SPSS version 22.0 software package for analysis.

## RESULTS

### Study groups

Two hundred and twenty seven diabetic patients were evaluated, 112 (49.3%) were men and 115 (50.7%) were women. Mean age of the patients was  $57.8 \pm 17.1$ . Clinical characteristics of the patients are shown in Table 1

**Table 1:** Characteristics of Study Participants. Mean  $\pm$  standard deviation (SD)


PARAMETER	FREQUENCY	PERCENTAGE (%)
<b>GENDER</b>	MALE-112	49.3
	FEMALE- 115	50.7
<b>AGE</b>	57.8+/-17.1	
<b>HYPERTENSION</b>	138	60.8
<b>HEART DISEASE</b>	50	22
<b>STROKE</b>	34	15
<b>RENAL REPLACEMENT THERAPY</b>	6	2.6

### Prevalence

Of the 227 patients with diabetes, 126 (55.5%) patients had anemia. Fifty-six (44.4%) were males and 70 (55.55%) were females. Among the anemic patients, 67 (53.17%) had normal mean corpuscular volume (MCV), while 58 (46.03%) had  $MCV < 80$  FL. while only one patient (0.7%) had  $MVC > 100$  FL.

### The effect of diabetic kidney disease

Of the total 227 patients with diabetes, 91 (40.1%) patients had  $GFR > 90$  mL/min/1.73 m<sup>2</sup> While 136 (59.9%) patients had  $GFR < 90$  mL/min/1.73 m<sup>2</sup> (Figure 1). Amongst patients with impaired kidney function, 88 (64.7%) had anemia while 48 (35.3%) had no anemia (chisquare 11.76,  $P < 0.05$ ).




**Figure 1:** Distributions of total sample over the stages of chronic kidney disease. Of the total 227 patients with diabetes, 40.1% had GFR>90 mL/min/1.73 m<sup>2</sup>, While the remaining 59.9% are distributed over the other stages of chronic kidney disease.

**Effect on vascular complications**

Diabetic patient with anemia were found to have heart disease (62%) compared with patients without anemia (38%) (Chi-square 14.44, P-value<0.05). Twenty-two diabetic patients with anemia (64.7%) had stroke compared with 12

(35.3%) patients without anemia (chi-square 8.47, P-value<0.05). Out of 138 diabetic patients with hypertension, 82 (59.4%) patients were anemic as compared to 56 (40.6%) patients who were non-anemic. (2,138, P-value<0.05) (Figure 2).




**Figure 2:** Distribution of vascular complications with respect to anemia. Diabetic patient with anemia found to have vascular complications more as compared with patients without anemia (P value <0.05).

**DISCUSSION**

Anemia is common in diabetes, potentially contributing to the pathogenesis of diabetes complications [15]. The prevalence of anemia in our study population is 55.5%, which is two to three times more common as compared to the result from previous studies (19-25%) [5,16]. Approximately one in two patients (126/227) had anemia as defined by gender-specific WHO guidelines. Prevalence of anemia in patients with diabetes correlated with renal impairment. The percentage of anemic patient with GFR<60mL/min/1.73 m2 (46%) which is comparable to the result from other study (36%) [5]. However, it is about two time more when compared to nonanemic patient in the same stages (17.8%) and this is may be explained that anemia increase the progression

of renal impairment as compared to those without anemia who are allocated in the early stages of kidney disease (Figure 3). Although anemia in diabetic patient correlated with degree of renal impairment, in our study anemia was six times more common (53.9%), compared to the previous study (9%) for the same GFR value (GFR>60 mL/min/1.73 m2) and half of them in stage one kidney disease unlike in previous study [5,16,17]. Despite the predominance of patients with anemia in the early stages of chronic kidney disease, their numbers are progressively decreasing with progression of kidney disease and this may be explained by increased morbidity and mortality as anemia closely associated with vascular complications.



**Figure 3:** Distributions of anemic and non-anemic patient over the stages of chronic kidney disease. The percentage of anemic patient with GFR<60 mL/min/1.73 m2 (46%) which is about two time more when compared to non-anemic patient in the same stages (17.8%).

Though the current recommendations for screening for anemia in the UK, target patients with stage 3 CKD (eGFR<60 ml / min per 1.73 m2), In our study it would fail to detect 53.9% of patients with diabetes and anemia with who have an EGFR>60 ml/min per 1.73 m2 [5]. So, ideally our patients with diabetes should be screened for anemia from the onset of diabetes.

Our study has several limitations. First, although the prevalence of anemia is high as compared to the previous studies, the main etiology for anemia was not investigated which is very crucial, though hemoglobinopathies are quiet rare and the variation of measured prevalence depends on the population studied and definition used [18]. Secondly, the interval between the diagnosis of

diabetes and the onset of anemia was not known in this retrospective study. Therefore, the ideal timing for anemia screening remains unclear or vague.

## CONCLUSION

Anemia is a common accompaniment with diabetes and it is seen early even in the absence of renal impairment. So, it may have further role in

the development and progression of both micro and macrovascular complications.

## Acknowledgement

I am grateful to SAVEETHA MEDICAL COLLEGE AND HOSPITAL for their support by using patients clinical and laboratory data. I would also like to extend my appreciation to study participants for providing necessary information for this study.

## REFERENCES

- [1]. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, et al. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract* 103, 2014, 137–149.
- [2]. Wright JA, Oddy MJ, Richards T (2014) Presence and characterization of anaemia in diabetic foot ulceration. *Anemia* 2014, 104-214
- [3]. Abate A, Birhan W, Alemu A Association of anemia and renal function test among diabetes mellitus patients attending Fenote Selam Hospital, West Gojam, Northwest Ethiopia: a cross sectional study. *BMC Hematol* 13(6), 2013.
- [4]. Thomas MC, Cooper ME, Rossing K, Parving HH Anaemia in diabetes: is there a rationale to TREAT. *Diabetologia* 49, 2006, 1151–1157.
- [5]. Jones SC, Smith D, Nag S, Bilous MT, Winship S, et al. Prevalence and nature of anaemia in a prospective, population-based sample of people with diabetes: teesside anaemia in diabetes (TAD) study. *Diabet Med* 27, 2010, 655–659.
- [6]. Shokoufeh B, Mohammad V, Mohammad G The prevalence of anemia in Iranian type two diabetic patients and the role of nephropathy. *Saudi J Kidney Dis Transpl* 22, 2011, 286-290.
- [7]. El-Achkar TM, Ohmit SE, McCullough PA, Crook ED, Brown WW, et al. Higher prevalence of anemia with diabetes mellitus in moderate kidney insufficiency: The Kidney Early Evaluation Program. *Kidney Int* 67, 2005, 1483-1488.
- [8]. New JP, Aung T, Baker PG, Yong sheng G, Pylypczuk R, et al. The high prevalence of unrecognized anaemia in patients with diabetes and chronic kidney disease: a population-based study. *Diabet Med* 25, 2008, 564–569.
- [9]. American Diabetes Association Diagnosis and classification of diabetes. *Diabetes care* 33, 2010, S62-S69.
- [10]. Beulter E, Waalen J The definition of anemia: what is the lower limit of normal of the blood hemoglobin concentration? *Blood* 107, 2006, 1747-50.
- [11]. Lam AP, Gundabolu K, Sridharan A, Jain R, Msaouel P, et al. Multiplicative interaction between mean corpuscular volume and red cell distribution width in predicting mortality of elderly patients with and without anemia. *Am J Hematol.* 88, 2013, E24.
- [12]. Isakov E, Fromm P, Henig C, Barak M Anemia and estimated glomerular filtration rates. *Ann Clin Lab Sci.* 44, 2014, 419-424.
- [13]. Liu X, Qiu X, Shi C, Huang H, Huang J, et al. Modified Glomerular Filtration Rate-Estimating Equations Developed in Asiatic Population for Chinese Patients with Type 2 Diabetes. *Int J Endocrinol* 9, 2014.
- [14]. Levey AS, Eckardt KU, Tsukamoto Y, Levin A, Coresh J, et al. Definition and classification of chronic kidney disease: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 67, 2089–2100.
- [15]. Thomas MC, MacIsaac RJ, Tsalamandris C, Molyneaux L, Goubina I, et al. The burden of anaemia in type 2 diabetes and the role of nephropathy: a cross sectional audit. *Nephrol Dial Transplant* 19, 2004, 1792–1797.
- [16]. Merlin CT, Richard JM, Con T, Lynda M, Inna G, et al. Anemia in patients with type 1 diabetes. *J Clin Endocrinol Metab* 89, 2004, 4359–4363.

- [17]. Thomas MC Anemia in diabetes: marker or mediator of microvascular disease? Nat Clin Pract Nephrol 3, 2007, 20-30.
- [18]. Alsaeed AH Prevalence of hemoglobinopathy disorders in adult patients sent for diagnosis of anemia in Saudi Arabia. Genet Test Mol Biomarkers 16, 2012, 25-29.

**How to cite this article:** Prabakaran Thamilmani, Kannan Rajendran, Rajendran K. Prevalence of anemia in type 2 diabetics-an overview. Int J of Allied Med Sci and Clin Res 2016; 4(4): 617-623.  
**Source of Support:** Nil. **Conflict of Interest:** None declared.