# Original Article



Online 2231 – 3656

# Abstract

# PHARMACOLOGICAL REVIEW OF LANTANA CAMARA .L REVIEW ARTICLE

<sup>\*</sup>Kumarasamyraja D, Jeganathan N S, Manavalan R Annamalai University, Annamalai Nagar, Chidhambaram, Tamil nadu, India – 608 002.

Herbal medicine is the oldest form of healthcare known to mankind. Herbs had been used by all cultures throughout history. India has a rich tradition of plant based knowledge in health care. Among the large number of herbal drugs existing in India, very few have been studied systematically so far. Lantana camara is a rugged evergreen shrub from the tropics. The plant extract are used in folk medicine for the treatment of chicken pox, measles, asthma, ulcers, swellings, eczema and high blood pressure. Decoction of plant used for tetanus, rheumatism, malaria and Decoction of fresh leaves used for gargle for toothaches. Pounded leaves used for cuts, ulcers and swelling. The present study was attempted to investigate the wide pharmacological activities of *lantana camara .L.* 

Key words: Herbal medicine, Lantana camara, Pharmacoloaical activities.

#### Introduction

The genus Lantana camara (Verbenaceae) as described by Linnaeus in 1753 contained seven species, six from South America and one from Ethiopia. Lantana camara from the Latin lento means to bend. Probably derives from the ancient Latin name of the genus Viburnum. Which it resembles a little in foliage and inflorescence<sup>1</sup> Lantana camara.L commonly known as weed or red sage, unniceeti (Tamil), pulikampa (Telugu) and caturang (Hindi) is a significant weed commonly found throughout in India<sup>2</sup>. It is ever green strong smelling shrub, with stout recurred prickles, leaves opposite, ovate or ovate -oblong, acute or sub acute, crenate serrate, scab rid on both side<sup>3</sup>. It is grew up to a height of 1-2m, occurs luxuriously in elevation up to 2000m in tropical<sup>4</sup>, sub tropical and temperate regions. It was introduced in India as an ornamental plant but entirely naturalized and found throughout India<sup>3</sup>, it was listed one of the important medicinal plants of the world<sup>6</sup>.

#### Author for Correspondence:

Kumarasamyraja D, Annamalai University, Annamalai Nagar, Chidhambaram, Tamil nadu – 608 002, India. Email: ksrajapharma83@gmail.com The fruits are useful in fistula, pustules, tumors and rheumatism.<sup>6-9</sup> The essential oil of Lantana camara showed a wide spectrum of antibacterial, antimicrobial and antifungal activity.<sup>10-12</sup> Lantana camara.L contains triterpenes; related compounds were lantandene C, reduced lantandene A, lantandene D, lantanolic and lantic acid<sup>13</sup>. Since very long time lantana camara.L reported to be used in traditional medicine system for itches, cuts, ulcers, swelling, bilious<sup>8,9</sup> fever, tetanus, and carminative<sup>3,14</sup> cold, head ache, uterine hemorrhage, chicken box, eye injuries, whooping cough, asthma<sup>5</sup>, bronchitis and arterial hypertension.<sup>15,16</sup> A tea prepared from the leaves and flowers was taken against fever, influenza and stomach - ache. In central and south America, the leaves were in to poultice to treat sores, chicken pox and measles and high blood pressure were treated with preparation from the plant<sup>16</sup>.

# Characteristics, Pharmacological Effects and Constituents

**Root:** sweet and bitter tasting, refrigerant, antifibrile. **Leaves:** minty tasting, cooling natured, antiphlogisitic, anti dermatoses.

Flowers: sweet tasting, mildly cooling, hemostastic.

1

Plant considered antiseptic, antispasmodic, vulnerary, diaphoretic and carminative.

# Antibacterial Activity

The extract of flower, leaf, stem and root of lantana showed antibacterial activity against camara.L Escherichia coli, pseudomonas aeruginosa, staphylococcus aureus. and staphylococcus saprohiticus. The antibacterial activity was determined by disc diffusion method<sup>17</sup>, tube dilution technique<sup>18</sup>, agar well diffusion method, micro distillation method<sup>19</sup>. Essential oil of lantana had considerable effect camara റെ antibacterial activity. It was observed that stains of E.coli and staphylococcus aureus were more susceptible to essential<sup>18</sup>. Lantana Camara flower extract posses strong antibacterial activity All few types' yellow, lavender, red and white lantana camara.L flowers displayed almost similar antibacterial activities. Petroleum ether root extract shown less antibacterial activity on pseudomonas aeruginosa and staphylococcus saprophiticus. The chloroform extract produced a moderate inhibition zone against staphylococcus aureus (5m). Chloroform stem extract showed inhibitory effect against staphylococcus saprophiticus<sup>19</sup>.

In other study, the essential oil of Lantana camara exhibited prominent antibacterial activity against all the bacterial strains tested. Gram positive Bacillus cereus, Bacillus subtilis and Staphylococcus aureus were the most sensitive strains to L. camara essential oil. Nevertheless, Gram negative Klebsiella pneumonia and Pseudomonas aeruginosa were not susceptible to the essential oil at lower concentration. A matter-of-fact, Gram-positive bacterium was more sensitive to the essential oils than gram-negative bacteria<sup>20</sup>.

# **Biochemical Composition and Antibacterial Activity**

Study of the leaves and flowers of four *lantana camara* plants with yellow, red, lavender and white flowers showed three of the four to have similar carbohydrates and lipid compositions. The carbohydrate levels were higher in the flowers than the leaves, and the lipids higher in the leaves expect for the lavender and white flowered kinds. The carbo in lavender L camara was very low. Leaf protein electrophoresis also showed similarities and differences. Antibacterial activities varied according to the type of tissue used<sup>18</sup>.

# Cytotoxicity

In vitro cytotoxicity activity of *lantana* camara *Linn* Study showed leaf extract of lantana camara is cytotoxic in nature and may possess antitumor activity that may be due to the presence of toxic lantanoids and alkaloids. Some taxa of lantana camara are toxic to ruminants and poisoning has been reported from Australia, india, new Zealand, south Africa and Americas<sup>21, 22, 36</sup>.

# Anti cancer Activity

Anti cancer effect of lantana camara root and leaf extracts against jurkat leukemia cell was investigated by MTT assay. Lantana camara leaf extract and root extract had roughly equaled anti proliferative activity on human leukemia jurkat cells. Morphological examinations indicated apotosis induction of the mechanisam of activity on jurkat cells. The lantana camara root and leaf extracts might be subjects for further fractionation and identification to find new anti cancer agents<sup>23</sup>.

# Anti fungal Activity

Lantana camara leaves evaluated for anti microbial and anti fungal activity and *L.camara* oil was also evaluated for larvicidal activity against different mosquito larvae<sup>37</sup>. The essential oil of lantana camara, tested against seven bacteria and eight fungi, showed wide spectrum of antifungal activities<sup>24</sup>.

# Anti helmentic Activity

Helminth infection is among the most common infection in men. Successive leaf extracts of *lantana* camara showed significant anthelmintic activity on selected warms. Anthelmintic activity was assessed using earth warms by the reported methods. Ethanolic extract found to be more active compared to remaining extracts<sup>4</sup>. Methanol extracts from the leaves, stems and roots of lantana camara were investigated for their anthelmintic activity against pheritima posthuma. The methnolic extract of stems of *lantana* camara were found to be most active<sup>25</sup>.

# Anti hyperglycemic Activity

A recent literature survey showed that lantana camara.L has clear the diabetes. Methanolic extracts of lantana camara. L administered orally in Alloxan induced diabetic rates, the results showed significant reduction in the blood glucose concentration in dose dependent manner and also promising anti hyperglycemic activity against Alloxan induced diabetic rates<sup>3</sup>.

#### Anti motility Activity

Evaluation of anti motility activity was done in intestine of mice treated with *lantana camara.L* leaf powder, neostigmine used as a pro motility agent. Intestinal motility was assessed by charcoal meal test. The anti diarrheal effect of *lantana leaf* extract was studied against castor oil induced diarrhea model in mice. When the plant extract were administered intra peritonealy. There was significant reduction in fecal output compared with castor oil treated mice<sup>26</sup>.

#### Anti micro-bacterial Activity

Chloroform and methanol extracts of *lantana camara.L* claimed for antimicrobial activity. The leaf extract of *lantana camara.L* was screened against three strains of mycobacterium tuberculosis using agar – well diffusion method. The MIC and MBC were determined using agar dilution method. Rifampicin kept as a standard dug for this study. Methanolic extract of *lantana camara.L* showed highest antimicrobial activity but it was much less activity then Rifampicin. The chloroform extract of *lantana* camara.L showed activity against all three strains of mycobacterium tuberculosis but it was less active than Methanolic extract<sup>27</sup>.

#### Anti oxidant Activity

The anti oxidant activity of Methanolic extract of *lantana camara.L* has been reported. The study showed anti oxidant potential of leaves of *lantana camara* extract measured in terms of reducing and scavenging activity. The Methanolic extract prepared from leaves I and III position exhibited significantly higher anti oxidant activity than leaves present from IV to V position<sup>28</sup>.

#### Anti ulcerogenic Activity

Methanolic extract of lantana camara.L has been reported anti ulcerogenic activity. Oral administrations of methanol extract of lantana camara.L exhibited dose dependent inhibition percentage respectively compared to ulcer control. Pre treatment with methanol extract of lantana camara.L leaves produced significant anti ulcer effect which can compared by aspirin induced ulcer<sup>3</sup>. The methanolic extract of lantana camara administered orally in pyrolic ligated rates, ethanol induced gastric ulcer and cysteamine induced duodenal ulcer. The lantana camara shown healing of gastric ulcer and also prevent the development of duodenal ulcer in rates<sup>29</sup>.

#### Wound healing Activity

Wounds are inescapable events of life, which arise due to physical injury, chemical injury or microbial infections<sup>30</sup> lantana camara used as an herbal medicine for the treatment of antiseptic for wounds and superficially for leprosy and scabies<sup>29</sup>.

The Ethanolic extract of *lantana* camara leaf was evaluated for their wound healing Potential in rates. Animals were experimentally wound in posterior neck area and treated with thin layer of blank placebo and placebo containing 5 and 10% lantana camara extract. Result of this topical application was assessed. Placebo containing leaf extracts significantly healed earlier than those treated with blank placebo histological analysis confirmed the results<sup>32</sup>.

In other study, investigation of burn wound healing activity of leaf extract of lantana camara on rates. Animals were anesthetized with ketamine. The burn wounds were created by pouring hot molten wax in to a metal cylinder placed on the back of the rates. The test group of the animals treated with ethanol extracts of lantana camara applied topically and control group left untreated. Wound healing assessed by rate of wound contraction and period of epithelialization. The extract showed antimicrobial activity aaainst staphylococcus aureus, klebsiella pneumoniae and E.coli. Extracted treated wounds were healed, which is not distinct from control and promoting the wound healing activity on burn wounds<sup>31</sup>.

#### Other Pharmacological activities

The Ethanolic extract of *lantana* camara ointment prepared for topical treatment on chronic crusty or acute lesions of Dermatophilosis<sup>33</sup>. Mosquito larvicidal activity and Phytochemical screening of methanol and ethanol extract of leaves and flowers of lantana camara larvicidal activity on III & IV instars larvae of mosquito species aedes aegypti and culexquinquefasciatus had been investigated an dose dependent manner for 24 h<sup>34</sup>. Essential from leaves of lantana camara possesses adulticidal activity against different mosquito species that was utilized for development of oil based insecticide as supplementary to synthetic insecticides <sup>35</sup>.

## Conclusion

Lantana camara is rugged evergreen shrub from the tropics. Despite its popularity as an ornamental plant, commonly found throughout in India. Lantana camara mainly used as an herbal medicine and in some areas as firewood and mulch. In some countries it is planted as a hedge to contain or keep out livestock. Leaf extracts of lantana exhibit antimicrobial, fungicidal, insecticidal and nematicidal properties. Which possess antimicrobial, immunosuppressive and anti tumor activity. Lantana oil is sometimes used for the treatment of skin itches, as an antiseptic for wounds and externally for leprosy and scabies. It has been used as natural insecticides before the discovery of synthetic organic insecticides<sup>34</sup>. The presence of alkaloids in L. camara has been indicated<sup>36</sup>. In future, there is tremendous scope in research for this plant.

#### References

- 1. Munir AA. J Adelaide Bot Gard 1996; 17:1.
- Ganesh T,Saikat sen, Thilagam G, Loganathan T, Raja Chakaraborty, Pharmacognostic and anti hyperglycemic evaluation of lantana camara (L) var. aculeate leaves in alloxan – induced hyperglycemic rats.Int.j.Res.Pharm. Sci. vol – 1, Issue -3, 2010, 247-252.
- Thamotharan G, Sekar G, Ganesh T, Saikat sen, Raja chakraborty, Senthil kumar N, Anti-Ulcerogenic effect of lantana camara Linn. Leaves on in vivo test models in Rats: Asian journal of pharmaceutical and clinical research, 3(4), 2010, 57-60.
- Jitendra patel, G.S. Kumar, Deviprasad S.P.., Deepika S., MD Shamim Qureshi, Phytochemical and Anthelmintic Evaluation of lantana camara (L.) var. aculeate leaves against pheretima posthuma, Journal of Global Trends in Pharmaceutical Sciences; 2(1), 2011, 11-20.

- Ross, I.A., Medicinal plants of the world. Chemical constituent, traditional and modern medical uses: New Jersey, Humana Press, 1999.
- 6. Kirtikar KR, Basu BD. Indian medicinal plants; vol. III, 1935, 1914.
- Chopra RN, Nayar SL, Chopra IC. Glossary of Indian medicinal plants. New Deli: CSIR, 1956:149.
- 8. CSIR. Wealth of India, vol. VI. Council of Scientific and Industrial Directorate, New Delhi, 1962:31.
- CSIR. The useful plants of India. Publication and Information Directorate, CSIR, New Delhi, 1992: 316.
- Siddiqui, B.S., Raza, S.M.Begum, S., Firdous, .S..Pentacyclic triterpenoids from Lantana camara.Phytochemistry, 38, 1995, 681–685.
- Barre, J.T., Bowden, B.F., Coll, J.C., Jesus, J.D., De La Fuente, V., Janairo, G.C., Ragasa, C.Y., A bioactive triterpene from Lantana Camara: Phytochemistry, 45, 1997, 321–324.
- Kumar, V.P., Chauhan, N.S., Padh, H., Rajani, M., Search for antibacterial and antifungal agents from selected Indian medicinal plants: Journal of Ethnopharmacology 107, 2006, 182–188.
- Abdel-Hady, Abdei-Halim AS, Al-Ghadban AM Chemical composition and insecticidal activity of the volatile oils of leaves and flowers of camara L. cultivated In Egypt: J Egypt Soc Parasitol, 35(2), 2005, 687-698.
- 14. Agarwal VS. Drug plants of India; vol. II. Ludhiana: Kalyani Publishers, 1997:426.
- Rastogi RP, Mehrotra BN. Compendium of Indian medicinal plants; vol. 1. Central Drug Research Institute, New Delhi: Lucknow and Publication and Information Directorate, CSIR, 1995:238.
- E.L.Ghisalberti,Lantana camara.L(Verbinaceae); A Review: Fitoterapia, 71, 2000, 467- 486.
- Mary kensa V. Studies on phytochemical screening and antibacterial activities of Lantana Camara Linn: Plant science Feed 2011-1(5): 74 – 79.
- Smaranika Pttnaik and Banita Pattnaik., A study of lantana camara. Linn aromatic oil as an antibacterial agent: Intr. Jpharmasci; 01(01), 2010, 32.
- Deepak Ganjewala, Silviya Sam, Kishwar Hayat Khan, Biochemical compositions and antibacterial activities of Lantana Camara plants with yellow,

Lavender, Red and white flowers: Eur. Asia J BioSci, 3, 2009, 69-77.

- Anil kumar saikia and Ranjan Kumar Sahoo, Chemical composition and Antibacterial Activity of Essential oil of Lantana camara L:.Middle – East journal of scientific research, 8 (3), 2011, 602.
- 21. Pass, M.A., Current ideas on the pathophysiology and treatment of Lantana poisonig of ruminants: Australian Veterinary Journal, 63, 1986,169–171.
- Seawright, A.A., Hrdlicka, J., The oral toxicity for sheep of triterpene acids isolated from *Lantana camara*: Australian Veterinary Journal, 53, 1977, 230–235.
- 23. Badakhshan Mahdi Pour, Lachimanan Yoga Latha 2 and Sreenivasan Sasidharan. Cytotoxicity and Oral Acute Toxicity Studies of Lantana camara Leaf Extract: Molecules, 16, 2011, 3663-3674.
- 24. M.J.Deena, J.E.Thoppil, Anti microbial activity of the essential oil of lantana camara, 71, 2000, 453-455.
- 25. A.S.Girme, R.D.Bhalke, P.B. Ghogare, V.D. Tambe, R.S.Jadhav and S.A. Nirmal, Comparative in vitro anthelmintic Activity of metha piperita and lantana camara from western india: Dhaka Univ, J.pharm. Sci. 5(1-2),2006, 5-7.
- 26. Lenika Sagar, Rajesh Sehgal and Sudarshan Ojha. Evaluation of antimotility effect of Lantana camara L. var. acuelata constituents on neostigmine induced gastrointestinal transit in mice:BMC Complementary and Alternative Medicine, 5, 2005, 18.
- 27. Claude Kirimuhuzya, Paul Waako, Moses Joloba, Olwa Odyek, The anti-mycobacterial activity of Lantana camara a plant traditionally used to treat symptoms of tuberculosis in South-western Uganda: African Health Sciences, 9(1), 2009.
- Dipita Bhakta and Deepak Ganjewala, Effect of Leaf Positions on Total Phenolics, Flavonoids and Proantho-cyanidins Content and Antioxidant Activities in Lantana Camara (L): J. Sci. Res, 1(2), 2009, 363-369.
- R. Sathish, Bhushan vyawahare, K. Natarajan, anti ulcerongenic activity of lantana camara leaves on gastric and duodenal ulcers in experimental rates, Journal of Ethanopharmacology, 134, 2011, 195-197.
- Sandeep B. Patil, N.S. Naikwade, M.S.Kondawar, C.S.Magdum and V.B. Awale. Traditional uses of

plants for wound healing in the sangli district, Mahastra: international journal of pharm Tech Research, 1(3), 2009, 876-878.

- 31. Nayak B.S, Raju S.S, Ramsubhag. A, Investigation of wound healing activity of Lantana camara L. in Sprague dawley rats using a burn wound model: International Journal of Applied Research in Natural Products, 1(1),2008, 15-19.
- Mahmood Ameen Abdulla, Pouya Hassandarvish, Hapipah Mohd Ali, Research journal of medicinal sciences, 3(2), 2009, 75-79.
- 33. N. Ali-Emmanuel, M. Moudachirou, J.A. Akakpo, J. Quetin-Leclercq, Treatment of bovine dermatophilosis with Senna alata, Lantana camara and Mitracarpus scaber leaf extracts: Journal of Ethnopharmacology, 86, 2003, 167–171.
- 34. M. Sathish Kumar and S. Maneemegalai, Evaluation of Larvicidal Effect of Lantana Camara Linn against Mosquito Species Aedes aegypti and Culex quinquefasciatus: Advances in Biological Research, 2 (3-4), 2008, 39-43.
- V.K.Dua, A.C. Pandey & A.P.Dash, Adulticidal activity of essential oil of Lantan canara Leaves against mosquitoes. Indian Med Res., 131, 2010, 434 – 439.
- Sharma OP, Makar HPS, Dawra RK: A Review of the noxious plant Lantana camara. Toxicon, 26, 1988, 975-987.
- 37. Vidya SS Dharmagadda, Mamta Tandonb and Padma Vasudevan: Biocidal activity of the essential oils of Lantana camara, Ocimum sanctum and Tagetes: journal of scientific & Industrial Research, 64, 2005, 53-56.