



Original Research Article

Quantitative determination of Trazodone hydrochloride in bulk and pharmaceutical dosage form by Zero order and AUC by UV-spectrophotometry

P S Jain^{1,*}, Santosh B Bodkhe¹¹Dept. of Pharmaceutical Chemistry, RC Patel Institute of Pharmaceutical Education & Research, Shirpur, Maharashtra, India

ARTICLE INFO

Article history:

Received 08-08-2020

Accepted 14-09-2020

Available online 28-10-2020

Keywords:

Trazodone hydrochloride(TZH)

UV spectrometric

Zero order

AUC

ABSTRACT

Two simple UV-Spectrophotometric methods have been developed for estimation of trazodone HCl (TZH) in bulk and pharmaceutical formulation form by using Zero order UV spectrometric absorbance method (method I) and Zero order UV spectrometric method using area under curve technique (method II). Water was used as solvent. In these Method λ max was found to be 246.0 nm of TZH and AUC between two wavelengths 235.20 nm – 256.60 nm. In both method Linearity in the concentration range of 05-35 μ g/ml ($r^2 > 0.99$) for TZH. Proposed methods were applied for oral dosage and amounts of TZH estimated by method I were found to be 99, by Method II were found to be 98.80 respectively. Both these methods were validated statistically and by recovery experiments.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

Trazodone HCl, 2-[3-[4-(3-chlorophenyl)-1-piperazinyl]propyl] - 1,2,4 - triazolo [4,3 - a] pyridin - 3 - (2H)-one mono hydrochloride, is a anti-depressant. Trazodone binds at 5-HT₂ receptor. It acts as a serotonin agonist at high doses and a serotonin antagonist at low doses. Like Fluoxetine, trazodone's antidepressant activity likely results from blockage of serotonin reuptake by inhibiting serotonin reuptake pump at the presynaptic neuronal membrane. The sedative effect of trazodone is likely the result of alpha-adrenergic blocking action and modest histamine blockade at H₁ receptor. It weakly blocks presynaptic alpha₂-adrenergic receptors and strongly inhibits postsynaptic alpha₁ receptors. Trazodone does not affect the reuptake of nor-epinephrine or dopamine within the CNS. In literature survey many analytical methods includes RP-HPLC¹⁻⁵ UV-spectroscopic¹⁻⁵ and HPTLC¹ methods have been reported for the estimation of TZH in bulk, pharmaceutical formulation n and in biological samples. In present study, simple, economical, accurate reproducible analytical method with better detection range for estimation of TZH

in its pure form and its pharmaceutical dosage were developed. This paper describes a UV-Spectrophotometric and Zero Order Derivative methods for estimation of TZH in bulk and pharmaceutical dosage form using Area Curve (AUC) technique. To our knowledge no method has been reported using water solvent on zero order UV-spectrometric method for estimation of TZH in Bulk and Pharmaceutical dosage form. Further methods validated as per ICH Q2R1 guideline.

2. Materials and Methods

2.1. Instrumentation

1. Spectrophotometer UV-2450 Shimadzu, Japan
2. Software UV Probe 2.21
3. Sample cell 1 cm matched quartz cell
4. Lamp Deuterium Lamp Wavelength range 200 - 400 nm
5. Detector Silicon Photodiode, Photomultiplier R-928
6. Scan speed Medium
7. Spectral slit width 1.0 nm
8. Weighing Balance Shimadzu AUX – 120

* Corresponding author.

E-mail address: [prishash79@yahoo.com](mailto:pritash79@yahoo.com) (P. S. Jain).

2.2. Selection of common solvent

R.O. water was selected as common solvent for developing spectral characteristics of TZH. The selection was made after evaluating the solubility of TZH in different solvents ACN, MeOH, 0.1N HCL

The stock standard solution was prepared by dissolving 10 mg of TZH in 100 mL of water to obtain a concentration of 100 $\mu\text{g/mL}$. The working standards were prepared by dilution of the stock standard solution.

2.3. Determination of λ_{max} and linearity study

Taken portion 1.0 mL of TZH were transferred to 10 mL volumetric flask, diluted to mark with water to obtain concentration of 10 $\mu\text{g/mL}$ for TZH. The resultant solution was scanned in UV range (200-400 nm) in 1.0 cm cell against solvent blank. The λ_{max} of TZH was found to be 246 nm. In Method I absorbance at 246 nm was considered for analysis while for Method II two wave lengths 235.20 nm to 256.60 nm were selected for determination of Area Under Curve [AUC] (Figure 2).

2.4. Linearity curve

For linearity study, seven solutions of TZH of different concentrations (5-35 $\mu\text{g/ml}$) were prepared using stock standard solution, analysed by proposed methods and the obtained data were utilized to plot calibration curves results was reported in Table 2 for both methods.

2.5. Analysis of pharmaceutical formulation: =

An accurately measured weight of 20 tablet equivalent to 10mg of TZH was transfer into 100 mL volumetric flask and volume was made up to the mark with water, filtered through 0.45 μm Whatman filter paper. A suitable volume of solution was further diluted with water to obtain concentration 15 $\mu\text{g/mL}$ of TZH. AUC were recorded in between the selected wavelengths and the concentrations were determined using respective linear regression equations. The analysis procedure was repeated for six times with same concentration.

2.6. Method validation

The method was validated terms of accuracy, precision and ruggedness, sensitivity, repeatability.

2.7. Accuracy

The accuracy of tablet of TZH by both methods was evaluated through recovery experiments. To the pre-analyzed sample solutions of concentration 15 $\mu\text{g/mL}$; a known amounts of stock standard solutions were added at different levels, that is, 80%, 100%, and 120%. The solutions were re-analyzed by the proposed methods. The




Fig. 1: Calibration curve




Fig. 2: Zero order spectrum of TZH showing lambda max and wave length selection between 235.20 and 256.60nm of AUC

experiments were performed for three times at each level for each method.

2.8. Precision (Intra-day and Inter-day precision)

Precision was determined as intra-day and inter-day variations.

Intra-day precision was determined by analysing the 10, 15 and 20 $\mu\text{g/mL}$ of TZH solution for three times in the same day morning, afternoon, and evening. Inter-day

Table 1: Analysis of Tablet (TZH)

	Label Claim(mg)	Conc ($\mu\text{g/ml}$)	% Amount found [n=6]	SD	%RSD
Method I	100	15	99.41%	0.004	0.49
Method II	100	15	99.72%	0.006	1.08

Table 2: Validation parameter

TZH		
Parameter	Method I	Method II
Working wavelength	246 nm	235.20 -256.60 nm
Linearity range ($\mu\text{g/ml}$)	05-35	05-35
Precision (%RSD)		
Inter-day[n=3]	0.09-0.6746	0.19-0.39
Intra-day[n=3]	0.20 – 1.03	0.17-1.4
Repeatability [n=6]	0.74	0.45
Ruggedness (%RSD)		
Analyst I [n=6]	0.59	0.47
Analyst II [n=6]	0.82	0.42
% Recovery		
Tablet	99.0	98.80
% RSD	0.38-0.77	0.18 – 0.25

precision was determined by analysing 10,15 and 20 $\mu\text{g/mL}$ of TZH drug solution daily for three consecutive days over a period of week results are reported in Table 2 for bulk as well as pharmaceutical formulation by both methods.

2.9. Repeatability

Repeatability was determined by analysing 15 $\mu\text{g/mL}$ of TRZ for six times and results reported in Table 2 for bulk and formulation by both methods.

2.10. Ruggedness

Ruggedness of the proposed method was determined by analysis of aliquots from homogenous slot by two analyst using same operational and environmental conditions and the results are reported in Table 2 for bulk and pharmaceutical formulation. by both methods.

2.11. Sensitivity

The sensitivity of proposed methods was determining in terms of Limit of Detection (LOD) and Limit of Quantification (LOQ) which were calculated using formulae “ $\text{LOD} = 3.3 \times \text{N/B}$ ” and “ $\text{LOQ} = 10 \times \text{N/B}$ where N is average standard deviation of the absorbance or peak areas of the TZH and “B” is the slope of the corresponding calibration curve. LOD and LOQ was found to be 0.38 and 1.18 respectively for method I and LOD and LOQ was found to be 0.27 and 0.84 respectively for method II.

3. Result and Discussion

In water TZH obeyed linearity in the concentration range of 05- 35 $\mu\text{g/mL}$, a correlation coefficient ($r^2 > 0.999$)

and equation was found in $Y = 0.032x - 0.0049$ for method I and a correlation coefficient ($r^2 > 0.995$) and equation was found in $Y = 0.1014x - 0.089$ for method II Marketed brand of tablet form were analysed. The amounts of (TZH) determined by ‘Method I’ was found to be 100% and 99.5%, for bulk and pharmaceutical dosage form respectively; and The amounts of TZH determined by ‘Method II’ was found to be 99.76 and 98.43%, for tablet and parenteral dosage form respectively in both these methods precision was studied as repeatability ($\% \text{RSD} < 2$) and inter and intra-day variations ($\% \text{RSD} < 2$) for drugs. The accuracy of method was determined by calculating mean percentage recovery. It was determined at 80,100 and 120% level. The ruggedness of the methods was studied by two different analysts using the same operational and environmental conditions. The % recovery, Table 2.

4. Conclusion

The proposed method is rapid, accurate, precise and sensitive for the quantification of trazodone hydrochloride and its pharmaceutical dosage forms by the spectrophotometric method. The method based on the use of simple working procedure comparable to that achieved by sophisticated and expensive technique such as HPLC, and hence this method can be routinely used in quality control for analysis of trazodone hydrochloride in tablets.

5. Source of Funding

None.

6. Conflict of Interest

None.

7. Acknowledgement

The authors are thankful to R.C. Patel Institute of Pharmaceutical Education and Research, Shirpur (M.S.), India for providing the required facilities to carry out this research work.

References

1. In: International conference of Harmonization (ICH) 1997 Q2R1 : Validation of analytical Procedure methodology. vol. 62;. p. 27463.
2. Beckett AH, Stenlake JB. Practical Pharmaceutical Chemistry. New Delhi: CBS Publication and distributors; 1997.
3. British Pharmacopoeia. Department of Health and Stationary office under the cotroller of majesty officer for Health minister UK; 2004.
4. ICH Steering Committee: International Conference on Harmonization (ICH) of Technical Requirements for Registration of Pharmaceutical for Human Use. Validation of Analytical Procedure-Methodology, Geneva, 2006.
5. The United States Pharmacopoeia, 24 revision, Asian Edition, United States Pharmacopoeial Convention. Inc, Twinbrook Parkway, Rockville, MD; 2000.

Author biography

P S Jain Professor

Santosh B Bodkhe PG Student

Cite this article: Jain PS, Bodkhe SB. Quantitative determination of Trazodoe hydrochloride in bulk and pharmaceutical dosage form by Zero order and AUC by UV-spectrophotometry. *Int J Pharm Chem Anal* 2020;7(3):145-148.