



Original Research Article

Medico-legal autopsy study of alleged dowry deaths - a two year cross-sectional study

Priyanka Patil A^{1,*}, Deepak Sangle J¹, Bhalchandra Chikhalkar²

¹Dept. of Forensic Medicine and Toxicological, Government of Medical College, Nagpur, Maharashtra, India

²Grant Medical College, Mumbai, Maharashtra, India



ARTICLE INFO

Article history:

Received 15-06-2020

Accepted 01-07-2020

Available online 06-08-2020

Keywords:

Alleged dowry deaths

Suicides

Menstruation

Asphyxia

ABSTRACT

Background: In India, dowry is a rising social evil contributing to female mortality.

Aims: To evaluate the socio-demographic profile of alleged dowry victims.

Materials and Methods: Cross-sectional study of 84 alleged dowry deaths stratifying the socio-demographic pattern of alleged dowry deaths.

Results: 57.1% were from 21to30 years of age group. Maximum females were illiterate (39.3%), housewives (77.4%) and from lower socio-economic strata of society. Victims were brought dead with 50% cases involving 71-90% of total body surface area. 29.8% were hospitalized. In 88.57 % of suicidal cases, menstruation was noted. 39.29% deaths were within 1st year of marriage. In 84.84% cases, dowry was provoking factor. In 50 % cases, Asphyxia was the cause of death.

Conclusion : Domestic abuse and dowry were significant provocative factors for dowry deaths.

© 2020 Published by Innovative Publication. This is an open access article under the CC BY-NC license (<https://creativecommons.org/licenses/by-nc/4.0/>)

1. Introduction

"Women are the architects of the society", as quoted by Harriet Beecher Stowe, an American Author. In India, women fall victims against violent crimes at high rates.¹ The purpose of dowry practice was probably to help new couples start their life in comfort which has recently grown into a social evil of bride- burning and suicides leading to social corruption.² According to National Crime Record bureau of India, total dowry deaths in the year 2007 and 2011 were 8093, 8618 respectively.³ This study enlightens the issue of dowry deaths in a metropolitan city of Maharashtra.

2. Materials and Methods

Cross-sectional study with a study period of two years conducted at the Department of Forensic Medicine and Toxicology, in a tertiary care centre in a metropolitan city. This study was approved by the Institutional Ethics Committee. Reference population included cases reported

and referred to this study centre, while study population included Unnatural female deaths within 7 years of marriage with the allegation of dowry death. 241 unnatural female deaths were examined in the study period, of which in 84 cases allegation of dowry death was observed. Various relevant socio-demographic and medico-legal aspects were stratified and analysed with the resource being Postmortem examination reports, documents of investigating agencies like police inquest Panchnama, Magistrate inquest Panchnama wherever applicable, Accidental Death Report and statements of relatives taken by police during investigation, Spot Panchnama and suicide notes if applicable. All findings were explained and portrayed in tabular form and inference was extracted. After entry, data was analysed using frequency distribution and statistical test with the help of computer generated software (S.P.S.S.16.0).

3. Results

Out of 241 Unnatural female deaths observed in the present study centre, 84 alleged dowry deaths are Analysed.

* Corresponding author.

E-mail address: pppriyankapatil1@gmail.com (P. Patil A).

Table 1 depicts age group distribution among alleged dowry deaths dowry victims (57.1%) belonged to the age group of 21 to 30 years of age and minimum (7.1%) belonged to the age group of 41 to 50 years of age.

Table 1: Age group distribution among alleged dowry deaths

| Age Group (Years) | Frequency | Percent age |
|-------------------|-----------|-------------|
| 11 to 20 | 11 | 13.1 |
| 21 to 30 | 48 | 57.1 |
| 31 to 40 | 19 | 22.6 |
| 41 to 50 | 6 | 7.1 |
| Total | 84 | 100.0 |

Table 2 describes occupational status of alleged dowry deaths. Maximum number of female victims of alleged dowry are housewives (77.4%) and minimum number of professionals (2.4%) with 1 victim who was working as house help.

Table 2: Occupational status among alleged dowry deaths

| Occupation | Frequency | Percentage |
|--------------|-----------|------------|
| Professional | 2 | 2.4 |
| Clerical | 3 | 3.6 |
| House help | 1 | 1.2 |
| Housewife | 65 | 77.4 |
| Laborer | 10 | 11.9 |
| Tailor | 3 | 3.6 |
| Total | 84 | 100.0 |

Table 3 shows distribution of educational status among alleged dowry victims. Maximum female victims are illiterate (39.3 %). Only (3.6%) female victims are Professionals and (2.4%) received intermediate school education.

Table 3: Educational status among alleged dowry victims

| Educational status | Frequency | Percentage |
|---------------------|-----------|------------|
| Illiterate | 35 | 39.3 |
| Primary | 7 | 8.3 |
| Middle | 26 | 31.0 |
| High school | 6 | 9.5 |
| Intermediate | 2 | 2.4 |
| Graduate | 5 | 6.0 |
| Professional degree | 3 | 3.6 |
| Total | 84 | 100.0 |

Table 4 describes socioeconomic status of alleged dowry victims according to Kuppaswamy's classification. Maximum female victims of alleged dowry death belonged to upper lower socioeconomic class (56 %) and minimum belonged to upper socio-economic class (21.4 %).

Table 5 depicts distribution of cases on the basis of hospitalisation. Out of 84 alleged dowry victims, 29.8% female victims are hospitalised and 59(70.2%) female victims are brought dead.

Table 4: Socio-economic status among alleged dowry deaths

| Socioeconomic status | Frequency | Percentage |
|-------------------------|-----------|------------|
| Upper (Class I) | 2 | 1.2 |
| Middle (Class III) | 19 | 21.4 |
| Upper Lower (Class IV) | 45 | 56.0 |
| Lower (Class V) | 18 | 21.4 |
| Total | 84 | 100.0 |

Table 5: Hospitalisation Among Alleged Dowry Deaths

| Hospitalisation | Frequency | Percentage |
|-----------------|-----------|------------|
| Yes | 25 | 29.8 |
| No | 59 | 70.2 |
| Total | 84 | 100.0 |

Table 6 denotes distribution of alleged dowry victims according to the place of incidence. Out of 84 female victims of alleged dowry , in 59 female victims (70.2%) , the incidence occurred at their own /in-laws residence , while in 25 female victims 29.8% cases , the incidence occurred at places other than their own residence.

Table 6: Place of Incidence Among Alleged Dowry Victims

| Place of incidence | Frequency | Percentage |
|--------------------|-----------|------------|
| Own /in-laws house | 59 | 70.2 |
| Other | 25 | 29.8 |
| Total | 84 | 100.0 |

Table ?? describes the distribution of percentage of burns injury among alleged dowry deaths and survival period distribution. In maximum burns cases , victims are brought dead 36.36% and percentage of burns in maximum cases(50%) is 71-90% of Total body surface area.

(Figures mentioned in parenthesis are percentages)

Table 8 shows association of menstruating uterus and suicidal dowry deaths. Out of a total of 35 menstruating female victims , 31 (88.57 %) suffered from suicidal deaths and 4 (11.42 %) are non- suicidal deaths. Out of 49 non-menstruating female victims , 33 (67.34 %) suffered non-suicidal deaths ,16 (32.66 %) suffered from suicidal deaths.

(Figures mentioned in parenthesis are percentages)

Table 9 Shows distribution of manner of death according to provocative factors in alleged dowry deaths.

Out of 47 suicidal deaths reported, in maximum cases (36 or 76.6%) , dowry was the provocative factor, in 6 victims (12.77%), domestic abuse was a provocative factor, in 2 cases (4.26%), infertility was an important provocative factor. 1 (2.13 %) female victim each has RVD and Psychiatric condition like depression is an important provocative factor for suicide.

In maximum suicidal cases within 7 years of marriage, dowry and domestic abuse are important provocative factors and RVD and Psychiatric diseases are least provocative factors for deaths.

Table 7: Distribution of percentage of burns according to survival period

| Percentage of burns | | 31 to 50 | 51 to 70 | 71 to 90 | >90 | Total |
|---------------------|---------------|----------|----------|----------|---------|-------|
| Survival period | Brought Dead | 0 | 3(37.5) | 4(50) | 1(12.5) | 8 |
| | 0 to 6 Hrs. | 0 | 1(100) | 0 | 0 | 1 |
| | 7 to 12 Hrs. | 0 | 2(33.33) | 4(66.66) | 0 | 6 |
| | 13 to 18 Hrs. | 0 | 1(33.33) | 2(66.66) | 0 | 3 |
| | 19 to 24 Hrs. | 0 | 1(100) | 0 | 0 | 1 |
| | 1 to 7 Days | 1(33.33) | 1(33.33) | 1(33.33) | 0 | 3 |
| Total | | 1 | 9 | 11 | 1 | 22 |

Table 8: Association of menstruating uterus and female deaths among alleged dowry deaths

| Uterus | Non-suicidal | Suicidal deaths |
|------------------|--------------|-----------------|
| Menstruating | 4 (11.42) | 31 (88.57) |
| Non-menstruating | 33 (67.34) | 16 (32.66) |
| Uterus | Non-suicidal | Suicidal deaths |

Out of 37 homicidal deaths within 7 years of marriage, in 2 (5.41 %) female deaths, the provocative factor for death is love affair, 9(24.32 %) female victims suffered homicidal deaths due to domestic abuse, 2 (5.41 %) female victims suffered homicidal deaths due to economical crisis like when involved in property disputes or high paying insurance policies. In 20 (54.05 %) female victims dowry is an important provocative factor for homicidal intent. 1 (2.7 %) female victim suffered homicidal death due to unemployment as a provocative factor, wherein the reason for impulsive anger of the accused is the lack of economical input from the female victim to support her family. In 3 (8.11 %) female victims, the cause of homicidal intent is not known.

In Maximum homicidal deaths within 7 years of marriage, dowry and domestic abuse are major provocative factors, whereas unemployment, love affair and economical crisis are least provocative factors for homicidal intent.

(Figures mentioned in parenthesis are percentages)

Table 10 shows distribution of marriage years according to provocative factors in Alleged dowry deaths. Maximum cases of alleged dowry deaths occurred in 1st year of marriage (39.29%). Dowry was a provocative factor in 84.84% of deaths within first year.

4. Discussion

Out of 84 female victims of alleged dowry death, maximum alleged female dowry victims were from the age group of 21 to 30 years of age and minimum were from the age group of 41 to 50 years of age.

Similar findings were noted in a study of Rajesh Kumar Verma et al,⁴ where in, maximum number of deaths (56.56%) occurred between 18-25 years of age and least (5.43%) in the age-group 31-35 years.

Following studies gave similar results- Shrivastava⁵ AK et al, Kulshreshtha⁶ P et al, Sharma⁷ BR et al, Harish D et al,⁸ Ghaffar UB⁹ et al, Kumar V,¹⁰ et al, Saha KK¹¹ et al

and Agnihotri A et al.¹²

The high incidence of unnatural deaths in young emotionally charged females is most probably due to unending demands of dowry (cash/ kinds) by their husbands and/or in-laws, for which they sometimes kill or torture the bride in such a way that she commits suicide, which is consistent with other authors findings.

And brides belonging to the age group of 41 to 50 years are those who committed a late marriage, which is noted mostly among educated independent working females and hence, the incidence of allegedly dowry deaths is comparatively less. This being a comparatively emotionally mature age group, alleged dowry deaths are comparatively less.

Maximum number of female victims of alleged dowry were housewives (77.4%) which may be attributed to financial dependency of housewives and minimum number of professionals (2.4%) with 1 victim who was working as house help.

Similar findings were noted in Prajapati P et al¹³ and Kumar T N S¹³ et al studies.

Maximum female victims were illiterate (39.3 %) .Only (3.6%) female victims are Professionals and (2.4%) received intermediate school education, which results in maximum dependant female population leading to exploitation and abuse of these females. In Saha et al¹¹ and Kumar V et al,¹⁴ wherein maximum female dowry victims are illiterate and minimum female victims are professionals similar to present study.

Maximum female victims of alleged dowry death belonged to upper lower socioeconomic class (56 %) and minimum belonged to upper socio-economic class (21.4 %). Dowry is considered as an evil of lower socio-economic group due to illiteracy and ignorance. Sharma BR et al⁷ and Kumar V et al¹⁴ states similar results.

Out of 84 alleged dowry victims, 29.8% female victims are hospitalised and 59 (70.2%) female victims are brought dead. Dowry victims in most cases are deprived from

Table 9: Distribution of manner of death according to provocative factors in alleged dowry deaths

| Provocative Factors | | Manner of death | | Total |
|---------------------|---------------------|-----------------|-----------|-------|
| | | Homicide | Suicide | |
| | Not known | 3(8.11) | 0 | 3 |
| | Love failure/affair | 2(5.41) | 1(2.13) | 3 |
| | Domestic abuse | 9(24.32) | 6(12.77) | 15 |
| | Economic crisis | 2(5.41) | 0 | 2 |
| | Dowry | 20(54.05) | 36(76.60) | 56 |
| | Aids | 0 | 1(2.13) | 1 |
| | Psychiatry | 0 | 1(2.13) | 1 |
| | Infertility | 0 | 2(4.26) | 2 |
| | Unemployment | 1(2.70) | 0 | 1 |
| | Total | 37 | 47 | 84 |

Table 10: Distribution of duration of years according to provocative factors in alleged dowry deaths

| Provocative factors | <1 | 1 to 2 | 2 to 3 | 3 to 4 | 4 to 5 | 5 to 6 | 6 to 7 | Total |
|----------------------|------------|-----------|------------|----------|----------|----------|----------|-------|
| Not known | 1(3.03) | 1(5) | 1(7.14) | 0 | 0 | 0 | 0 | 3 |
| Love Failure/affairs | 0 | 1(5) | 0 | 0 | 2(33.33) | 0 | 0 | 3 |
| Domestic abuse | 4(12.12) | 2(10) | 5(35.71) | 0 | 1(16.67) | 2(40) | 1(50) | 15 |
| Economic Crisis | 0 | 1(5) | 0 | 0 | 1(16.67) | 0 | 0 | 2 |
| Dowry | 28(84.84) | 15(75) | 6(42.86) | 2(50) | 1(16.67) | 3(60) | 1(50) | 56 |
| Aids | 0 | 0 | 0 | 1(25) | 0 | 0 | 0 | 1 |
| Psychiatry | 0 | 0 | 1(7.14) | 0 | 0 | 0 | 0 | 1 |
| Infertility | 0 | 0 | 0 | 1(25) | 1(16.67) | 0 | 0 | 2 |
| Unemployment | 0 | 0 | 1(7.14) | 0 | 0 | 0 | 0 | 1 |
| Total | 33 (39.29) | 20 (23.8) | 14 (16.67) | 4 (4.76) | 6 (7.14) | 5 (5.95) | 2 (2.38) | 84 |

(Figures mentioned in parenthesis are percentages)

hospital services which may be because of ignorance of their relatives to hospitalise the victims.

Out of 84 female victims of alleged dowry, in 59 female victims (70.2%), the incidence occurred at their own residence, while in 25 female victims 29.8% cases, the incidence occurred at places other than their own residence. Similar findings are noted by Dasari et al.¹⁵ Sixty one percent females (105) suffered burns in their in-laws house while the rest at other places.

Similar findings were noted by Jaswinder Singh¹⁶ et al, wherein 67.34% incidences occurred in in-laws house, while remaining occurred in other places.

In maximum burns cases, victims are brought dead 36.36% and percentage of burns in maximum cases(50%) is 71-90% of Total body surface area. With increasing percentage of body surface area involved in burns injury, the fatality rate increases.

Similar to Sharma B R¹⁵ and Modi JP¹⁶ also states that death in burns injury may occur within 24 -48 hours, but the first week is most fatal.

Out of a total of 35 menstruating female victims, 31 (88.57 %) suffered from suicidal deaths and 4 (11.42 %) are non-suicidal deaths.

Out of 49 non-menstruating female victims, 33 (67.34 %) suffered non-suicidal deaths, 16 (32.66 %) suffered from suicidal deaths.

Strong association between menstruating uterus and suicidal deaths is observed.

The first (or menstrual) and fourth (or premenstrual) week of the menstrual cycle may be associated with many suicide attempts in women, stated by Baca –Garcia¹⁷ et al.

Out of 47 suicidal deaths within 7 years of marriage, in 1 (2.13 %) female victim, the provocative factor for death is love failure/ affair of the husband, 6 (12.77 %) female victims committed suicide due to domestic abuse by the in-laws and 36(76.60 %) female victims committed suicide with dowry as a provocative factor. 2 (4.26 %) female victims committed suicide with infertility as a provocative factor and 1 (2.13 %) female victim each has RVD and Psychiatric condition like depression is an important provocative factor for suicide. In maximum suicidal cases within 7 years of marriage, dowry and domestic abuse were important provocative factors and RVD and Psychiatric diseases are least provocative factors for deaths.

Out of 37 homicidal deaths within 7 years of marriage, in 2 (5.41 %) female deaths, the provocative factor for death is love affair, 9(24.32 %) female victims suffered homicidal deaths due to domestic abuse, 2(5.41 %) female victims suffered homicidal deaths due to economical crisis like when involved in property disputes or high paying insurance policies. In 20 (54.05 %) female victims dowry is an important provocative factor for homicidal intent.1

(2.7 %) female victim suffered homicidal death due to unemployment as a provocative factor, wherein the reason for impulsive anger of the accused is the lack of economical input from the female victim to support her family. In 3 (8.11 %) female victims, the cause of homicidal intent was not known.

In Maximum homicidal deaths within 7 years of marriage, dowry and domestic abuse are major provocative factors, whereas unemployment, love affair and economical crisis are least provocative factors for homicidal intent.

Similar to Biswadeep Paul et al,¹⁸ family disputes pertaining to Dowry deaths including failure in love accounted for highest number of burns injury deaths. In 2010, according to NCRB, a total of 8391 and in 2016, 8618 dowry deaths are reported in India, while in Assam according to Crime Statistics, 121 female deaths are observed and according to National Crime Bureau, a dowry death occurs every 77 minutes.

Similar to the findings of Guntheti et al,¹⁹ where in dowry and harassment from in-laws are important provocative factors. Similar findings were noted in Dere Rajesh²⁰ et al and Meera Tet al.²¹

According to T.C.Siddaramanna²² et al, family issues are important cause of poisoning and Chakrabarthi²³ et al, more than one-third of deaths due to domestic conflict, marital disharmony, quarrel.

Another study with similar findings is Shrivastav⁵ et al, with dowry (25%) and domestic abuse (26.39%) being an important provocative factor for suicidal deaths and infertility (4.17%) and poverty (5.56%) being the provocative factors for least suicidal deaths among newly married females.

Maximum cases of alleged dowry deaths occurred in 1st year of marriage (39.29 %) with dowry as an important provoking factor in 84.84% of deaths in 1st year of marriage.

Dowry and domestic abuse were important provocative factors for death after marriage within 7 years in alleged dowry victims. Dowry was an important factor for death within first three years of marriage. While infertility is an important provocative factor in the 3rd and 4th decade of married life, when the family pressure for bearing a child increases and hence, is a trigger factor for suspicious death.

This relationship of provocative factors and duration of marriage years was studied uniquely in this study.

5. Conclusions

Incidence of dowry deaths among unnatural deaths is 34.85 % which is higher considering the fact that the study is conducted within city.

Maximum cases of alleged female dowry deaths were from younger and reproductive group which is emotionally labile.

Maximum housewives, illiterate belonging to upper lower socioeconomic class suffered death.

Maximum deaths occurred within first year of marriage and did not receive hospitalization services.

Dowry was an important provocative factor in most of the deaths along with domestic abuse and infertility.

Most of the deaths occurred at their own or in-laws residence.

Most of the alleged cases of dowry were suicidal and most of the suicides were during the menstruation phase of cycle.

Asphyxia was the cause of death in most cases and a significant number of Burns injury cases were also noted.

6. Source of Funding

None.

7. Conflict of Interest

None.

References

- Mangoli RN, Tarase GM. Crime against woman in India. A Statistical review. *Int J Criminol Sociol Theory*. 2009;2(2):292–302.
- Park K. Park's Textbook of Preventive and Social Medicine, Jabalpur. BanarsidasBhanot Publishers; 2011. p. 525–645.
3. Crime in India 2011 Compendium. New Delhi: Government of India, Minister of Home Affairs, National Crime Record Bureau Report; 2012. p. 190.
- Datta D. "The new laws of marriage". India Today; 2013.
- Shrivastava AK, Arora P. Suspicious Deaths in Newly Married Females-A Medicolegal Analysis. *J Indian Acad Forensic Med*. 2007;29(4):63–7.
- Kulshreshtha P, Sharma RK, Dogra TD. Study of Sociological and Demographical Variables of Unnatural Deaths Among Young Women in South Delhi within seven years of marriage. *J Punjab Acad Forensic Med Toxicol*. 2002;2:7–17.
- Sharma BR, Singh VP, Sharma R, Sumedha. Unnatural Deaths In Northern India-A Profile. *J Indian Acad Forensic Med*. 2004;26(4):140–6.
- Sharma BR, Harish D, Sharma A, Singh H. Accidental burns in Indian kitchen; Are they really accidents? *JIAFM*. 2006;28(1):14–7.
- Ghaffar UB, Hussain M, Rizvi SJ. Thermal Burn: An Epidemiological Prospective Study. *J Indian Acad Forensic Med*. 2008;30(1):10–4.
- Koulapur VV, Yoganarsimha K, Gouda H, Mugadlimath A, Kumar V. Analysis of Fatal Burns Cases-A 5 Year study At Sri B M Patil Medical college. *Medico-Legal Update*. 2011;11(2):107–9.
- Saha KK, Mohanty S. Alleged Dowry Death: A study of homicidal burns. *Med Sci Law*. 2006;46(2):105.
- Agnihotri A, Aggarwal A. The Epidemiological Study of Dowry Death Cases with Special references to burn cases in Allahabad zone. *Internet J F M T*. 2001;2(1):201–5.
- Kumar T, Mahathi K, Thirthala S. Role of Constitution in Women Empowerment in Corporate Sector. *Arabian J Bus Manag Rev*. 2016;6:267.
- Kumar V, Tripathi CB, Kanth S. Burn Wives - A Sociologic Study. *Int J Med Toxicol Legal Med*. 1999;2(1):18.
- Sharma BR, Harish D, Sharma V, Vij K. Kitchen accidents vis-a-vis dowry deaths. *Burns*. 2002;28:250–3.
- Modi JP. Textbook of Medical Jurisprudence and Toxicology. Lexis Nexis; 2016. p. 159–62.
- Baca-Garcia E, Diaz-Sastre C, Ceverino A, Perez-Rodriguez MM, Navarro-Jimenez R, Lopez-Castroman J, et al. Suicide attempts

- among women during low estradiol/low progesterone states. *J Psychiatr Res.* 2010;44(4):209–14.
18. Paul B, Mahanta P, Tamuli RP. A Statistical Analysis of Death Victims brought for Medico-legal Autopsy. *Medico-Legal Update.* 2013;13:98–101.
 19. Guntheti BK, Singh UP. The pattern of Poisoning in Khammam. *J Indian Acad Forensic Med.* 2011;33(4):296–300.
 20. Dere RC, M CRK. Study of Unnatural Deaths in Females A Medicolegal Study at Rural Medical College. *J Indian Acad Forensic Med.* 2011;33(3):211–3.
 21. Meerath, Singh M. Pattern of neck findings in suicidal hanging a study in Manipur. *Journal of Indian Academy of Forensic Medicine.* 2011;33(4):352–354.
 22. Siddaramanna TC, Kumar RD, Rudramurthy S. Deaths due to organophosphorus poisoning-A retrospective study. *J Indian Acad Forensic Med.* 2016;38(3):278–9.
 23. Chakraborty P, Tudu NK, Saren AB, Dwari AK, Haldar D, Samanta S, et al. Epidemiology of unnatural death from suspected poisoning :

an autopsy based study. *IOSR-J Dent Med Sci.* 2015;p. 1–8.

Author biography

Priyanka Patil A Assistant Professor

Deepak Sangle J Assistant Professor

Bhalchandra Chikhalkar Professor and Head

| |
|---|
| <p>Cite this article: Patil A P, Sangle J D, Chikhalkar B. Medico-legal autopsy study of alleged dowry deaths - a two year cross-sectional study. <i>IP Int J Forensic Med Toxicol Sci</i> 2020;5(2):64-69.</p> |
|---|