

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Orthopaedics Surgery

Journal homepage: <https://www.ijos.co.in/>

Original Research Article

Comparison of early clinico-radiological outcome of posterolateral fusion and transforaminal lumbar interbody fusion techniques

Uday Phute¹, Gajanan Deshmukh^{1,*}, Sushant Sopan Tuse¹¹Dept. of Orthopaedic, Seth Nandlal Dhoot Hospital, Aurangabad, Maharashtra, India

ARTICLE INFO

Article history:

Received 05-01-2022

Accepted 17-05-2022

Available online 15-06-2022

Keywords:

Posterolateral fusion

Transforaminal lumbar interbody

fusion

ABSTRACT

Study Design: Retrospective cohort study.**Objective:** The objective is to analyse clinical & radiological outcome with respect to functional disability, pain, fusion rate of patients treated with TLIF compared to instrumented PLF techniques for lumbar spinal stabilization.**Materials and Methods:** Monocentric retrospective study with an average follow up of 36 months in patients who underwent surgery from January 2016 to December 2017. Out of 140 participants, 78 males and 62 females with mean age 52.22 (± 11.97) years; 78 underwent PLF and 62 underwent TLIF. Assessment was done using VAS score and ODI score before surgery and post surgery at 3, 6, and 18 weeks and thereafter at 18, 24 and 36 months. Radiologically sagittal and coronal angles were measured both preoperatively and postoperatively and evaluation of correction in sagittal and coronal angle was calculated.**Results:** We observed highly significant reduction in the post-operative VAS score, ODI Score compared to pre-operative scores in both TLIF & PLF group but radiologically TLIF gives better correction in sagittal balance and rotational alignment compared to PLF.**Conclusion:** We conclude that in the short term duration of our study, both the procedures done with proper technique in duly indicated patients shows satisfactory clinical outcome. However, radiologically TLIF patients had better outcome. We expect better outcome in long term with TLIF compared to PLF. In presence of insignificant blood loss, surgical duration and better 360° fusion TLIF is preferred over PLF.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

With increase in life expectancy the degenerative spinal disorder have also increased like spinal stenosis, degenerative disc disease, and degenerative deformities.^{1,2} Most of the patients are initially treated with conservative approach. Those who do not have relief with conservative approach or in presence of neurological deterioration, surgical option is considered in the form of spinal fusion along with decompression.

Instability due to degenerative spine disease is treated surgically by achieving fusion either posterolateral or by interbody technique. Both methods are equally effective but some researchers believe TLIF provides better mechanical stability and maintains it.

2. Materials and Methods

2.1. Study design

Retrospective cohort study.

Patients who have been operated for spinal instability, Spinal stenosis, degenerative disc disease, spondylolisthesis

* Corresponding author.

E-mail address: gajanan9690@gmail.com (G. Deshmukh).

with neurological symptoms were included in the study. Patients who had severe osteoporosis, infection, or malignancy were excluded from the study.

Written consent of all the patients were taken and study is approved by Ethics Committee of Seth Nandlal Dhoot Hospital, Aurangabad.

Duration of the study was between January 2016 and December 2017. Patients were followed up retrospectively till Dec 2019 with an average follow up of 36 months. Total sample size was 140 patients out of which 78 (n=78, 55.71%) were male and 62 (n=62, 44.29%) were female.

2.2. *Plf/lif*

Out of 140 patient 78 underwent PLF and 62 underwent TLIF (Tables 1 and 2). The mean age of the patients in the TLIF group was 52.82 (± 12.23), and the mean age of the patients in the PLF group was 51.47 (± 11.68). Out of 78 PLF patient 47 are male and 31 are female and out of 62 TLIF patients 31 are male and 31 are female. Clinical and radiological outcomes were serially evaluated upto 36 months retrospectively in each cohort.

2.3. *Surgical procedure*

After standard soft tissue dissection pedicle screws were inserted, decompression was done in the form of laminectomy, excision of hypertrophied Ligamentum Flavum and medial facetectomy if required for all the patients.

In PLF group distraction was done and rods were placed and then decortication of transverse process and pars was done. Locally harvested autograft was placed posterolaterally for fusion.

In TLIF group distraction was done and rods were placed. Discectomy done and end plates curated, locally harvested autograft placed anteriorly and then bone graft impacted cage was inserted.

2.4. *Assessment and outcome*

Patient's outcome was assessed on the basis of data which was taken from Medical record department of hospital,

1. Clinically patient's pain and well-being was assessed using VAS score and ODI score before surgery and post-surgery at 3 weeks, 6 weeks and 18 weeks and thereafter at 18, 24 and 36 months.
2. Radiologically lumbar sagittal and coronal angles were assessed both preoperatively and postoperatively on AP and lateral X ray views. Evaluation of correction in sagittal and coronal angle were done by Cobbs angle.
3. Radiologically the fusion was assessed by Lenke's method.

The four grades according to Lenke's method was judged by anteroposterior radiograph which are as follows:-

Grade A:- bilateral definitely solid stout fusion masses present;

Grade B:- probably solid with a unilateral stout fusion mass and contralateral thin fusion mass;

Grade C:- probably not solid with a thin unilateral fusion mass; and

Grade D:- definitely not solid with thin fusion masses bilaterally with obvious pseudoarthrosis or bone graft dissolution bilaterally.

Brand Tegan and Steffee method of grading interbody fusion which was modified to describe the Fraser definition of locked pseudoarthrosis (BSF scale) was used.

In order to eliminate any possible errors regarding the x ray interpretation, all x-rays were evaluated by the operating team and by another orthopaedic surgeons & result were analysed statistically.

3. Results

The following table shows comparison of pre-operative and post-operative VAS Score, ODI Score, Coronal Angle and Lumbar Lordosis Angle in PLF group.

VAS Score: The mean pre-operative VAS score was 6.048 (± 1.487) and mean post-operative VAS score was 2.839 (± 1.074). The t-test analysis indicates a highly significant reduction in the post-operative VAS score ($p < 0.001$).

ODI Score: The mean pre-operative ODI score was 54.774 (± 9.855) and mean post-operative ODI score was 12.177 (± 2.917). The t-test analysis indicates a highly significant reduction in the post-operative ODI score ($p < 0.001$).

Coronal Angle: The mean pre-operative coronal angle was 16.097 (± 1.676) and mean post-operative coronal angle was 4.677 (± 0.937). The t-test analysis indicates a highly significant reduction in the post-operative mean coronal angle ($p < 0.001$).

Lumbar Lordosis Angle: The mean pre-operative Lumbar Lordosis Angle was 20.790 (± 1.794) and mean post-operative Lumbar Lordosis Angle was 33.016 (± 1.937). The t-test analysis indicates a highly significant increase in the post-operative mean Lumbar Lordosis Angle ($p < 0.001$).

The following table shows comparison of pre-op and post-op VAS Score, ODI Score, Coronal Angle and Lumbar Lordosis Angle in TLIF group.

VAS Score: The mean pre-operative VAS score was 5.833 (± 1.189) and mean post-operative VAS score was 2.872 (± 0.998). The t-test analysis indicates a highly significant reduction in the post-operative VAS score ($p < 0.001$).

ODI Score: The mean pre-operative ODI score was 56.782 (± 8.294) and mean post-operative ODI score was

Table 1:

Study Variable	Posterolateral fusion		Transforaminal lumbar interbody fusion		Total	
	n	%	n	%	n	%
Occupation of patient						
House wives	21	26.92	17	27.42	38	27.14
Farmers	21	26.92	14	22.58	35	25.00
Drivers	15	19.23	12	19.35	27	19.29
Manual labourer	14	17.95	10	16.13	24	17.14
Computer professional	2	2.56	6	9.68	8	5.71
Company workers	3	3.85	2	3.23	5	3.57
Mechanic	2	2.56	1	1.61	3	2.14
Patient condition						
Spondylolysthesis	20	25.64	22	35.48	42	30.00
Spinal stenosis	22	28.21	17	27.42	39	27.86
Degenerative disc disease	18	23.08	16	25.81	34	24.29
Spinal instability	18	23.08	7	11.29	25	17.86
Age of patient (in Years)						
< 40	11	14.10	9	14.52	20	14.29
40-59	38	48.72	34	54.84	72	51.43
>= 60	29	37.18	19	30.65	48	34.29
Mean Age (\pm SD)	51.47 (\pm 11.68)		52.82 (\pm 12.23)		52.22 (\pm 11.97)	
Gender of patient						
Male	47	60.26	31	50.00	78	55.71
Female	31	39.74	31	50.00	62	44.29
Total	78	55.71	62	44.29	140	100.00

Table 2:

Pre-Post comparison of study variables in PLF group							
Study Variable		Mean	N	SD	SEM	t-stat	p-value
VAS	Pre-operative	6.048	62	1.487	0.189	25.425	<0.001**
	Post-operative	2.839	62	1.074	0.136		
ODI	Pre-operative	54.774	62	9.855	1.252	38.900	<0.001**
	Post-operative	12.177	62	2.917	0.370		
Coronal Angle	Pre-operative	16.097	62	1.676	0.213	53.689	<0.001**
	Post-operative	4.677	62	0.937	0.119		
Lumbar Lordosis Angle	Pre-operative	20.790	62	1.794	0.228	-39.914	<0.001**
	Post-operative	33.016	62	1.937	0.246		

Operation Procedure = Posterolateral Fusion (PLF);

** : Sig. at 1 % level of significance

13.218 (\pm 3.238). The t-test analysis indicates a highly significant reduction in the post-operative ODI score ($p < 0.001$).

Coronal Angle: The mean pre-operative coronal angle was 13.205 (\pm 1.976) and mean post-operative coronal angle was 4.500 (\pm 1.114). The t-test analysis indicates a highly significant reduction in the post-operative mean coronal angle ($p < 0.001$).

Lumbar Lordosis Angle: The mean pre-operative Lumbar Lordosis Angle was 20.038 (\pm 2.371) and mean post-operative Lumbar Lordosis Angle was 30.038 (\pm 2.230). The t-test analysis indicates a highly significant

increase in the post-operative mean Lumbar Lordosis Angle ($p < 0.001$).

The Table 3 indicates significance of the mean difference between the pre-operative and post-operative mean VAS score, ODI score, coronal angle and lumbar Lordosis angle.

VAS Score: The mean difference between pre-operative and post-operative VAS score in PLF group was 3.210 (\pm 0.994) and in TLIF group it was 2.962 (\pm 0.959). The result of t-test indicates no significant difference in the mean difference in two groups ($p = 0.137$).

ODI Score: The mean difference between pre-operative and post-operative ODI score in PLF group was 42.598

Table 3: Pre-post comparison of study variables in TLIF group

Study Variable		Mean	N	SD	SEM	t-stat	p-value
VAS	Pre-operative	5.833	78	1.189	0.135	27.261	<.001**
	Post-operative	2.872	78	0.998	0.113		
ODI	Pre-operative	56.782	78	8.294	0.939	52.413	<.001**
	Post-operative	13.218	78	3.238	0.367		
Coronal Angle	Pre-operative	13.205	78	1.976	0.224	37.352	<.001**
	Post-operative	4.500	78	1.114	0.126		
Lumbar Lordosis	Pre-operative	20.038	78	2.371	0.268	-40.959	<.001**
	Post-operative	30.038	78	2.230	0.252		

Angle

Operation Procedure = Transforaminal Lumbar Interbody Fusion (TLIF);

** : Sig. at 1 % level of significance

(±8.622) and in TLIF group it was 43.564 ((±7.341).The result of t-test indicates no significant difference in the mean difference in two groups (p=0.475).

Coronal angle: The mean difference between pre-operative and post-operative Coronal angle in PLF group was 11.419 (±1.675) and in TLIF group it was 8.705((±2.058).The result of t-test indicates a highly significant difference in the mean difference in two groups (p<.001).

Lumbar Lordosis Angle: The mean difference between pre-operative and post-operative Lumbar Lordosis Angle in PLF group was -12.226 (±2.412) and in TLIF group it was -10.000 ((±2.156).The result of t-test indicates a highly significant difference in the mean difference in two groups (p<0.001).

The Table 5 indicates the significance of association between LENKES grading and operation procedure. The analysis of chi-square shows no significant difference in the LENKES grading according to the operation procedure (Chi-square = 0.817, p=0.845).

The Table 6 indicates the significance of association between BSF scale and operation procedure. The analysis of chi-square shows no significant difference in the BSF scale according to the operation procedure (Chi-square = 0.952, p=0.621).

4. Discussion

We have retrospectively analysed a case series of 140 TLIF and PLF patients. Several non-randomized studies and one small randomized controlled trial comparing TLIF and PLF, suggests that PLF is associated with fewer complications, less blood loss, shorter operative time and hospital duration, leading to lower health care costs and an improvement in quality of life but the reoperation rate was statistically higher for PLF.³ In the last decade, with various studies superior results of interbody spinal fusion has increased steeply,⁴ and with the increase in life expectancy of population this is expected to rise further. Therefore we attempted to verify these presumed advantages of TLIF over PLF.

In contrast to the literature we did not find difference in estimated intraoperative blood loss or duration of hospitalization between both procedures. A shorter surgical time was observed in the PLF group compare to TLIF group which was not significant.

Complications like Superficial infection was seen in 3 cases of PLF group and 2 cases of TLIF group. They were treated with appropriate antibiotics and resolved completely. There was one patient of hardware failure in PLF group however patient was clinically asymptomatic. With both the techniques, there is significant reduction in back as well as leg pain post operatively.

Comparing our series to other reported series we observed some similarities and some differences.




Fig. 1: Preoperative and postoperative pain assessment by Oswestry disability index




Fig. 2: Radiographic assesment of the lumbar alignment

Table 4: Significance of mean difference between pre-op and post-op values of study variables

Study variable	Operation Procedure		Mean	MD	SD	SEDM	p-value
VAS Score	PLF	Pre-operative	6.048	3.210	0.994	0.126	0.137, NS
		Post-operative	2.839				
	TLIF	Pre-operative	5.833				
		Post-operative	2.872				
ODI Score	PLF	Pre-operative	54.774	42.597	8.622	1.095	
		Post-operative	12.177				
	TLIF	Pre-operative	56.782				
		Post-operative	13.218				
Coronal Angle	PLF	Pre-operative	16.097	11.419	1.675	0.213	<0.001**
		Post-operative	4.677				
	TLIF	Pre-operative	13.205				
		Post-operative	4.500				
Lumbar Lordosis Angle	PLF	Pre-operative	20.790	-12.226	2.412	0.306	
		Post-operative	33.016				
	TLIF	Pre-operative	20.038				
		Post-operative	30.038				

Table 5: Association between LENKES grading and operation procedure

LENKES Grading	Operation Procedure				Total	
	PLF		TLIF		n	%
	n	%	n	%	n	%
Grade A	57	91.94	68	87.18	125.0	89.29
Grade B	1	1.61	2	2.56	3.0	2.14
Grade C	1	1.61	2	2.56	3.0	2.14
Grade D	3	4.84	6	7.69	9.0	6.43
Total	62	100	78	100	140	100

Chi-square = 0.817, df=3, p=0.845, not significant

Table 6: Association between BSF scale and operation procedure

BSF Scale	Operation Procedure				Total	
	PLF		TLIF		n	%
	n	%	n	%	n	%
BSF 1	2	3.23	3	3.85	5	3.57
BSF 2	3	4.84	7	8.97	10	7.14
BSF 3	57	91.94	68	87.18	125	89.29
Total	62	100.00	78	100.00	140	100.00

Chi-square = 0.952, df=2, p=0.621, not significant

Audat et al.⁵ compared PLF, PLIF, and TLIF for degenerative disc disease in 81 patients between 2003 to 2006 and found that no significant difference existed between the three groups in terms of clinical, radiological outcome and complications. However, the best radiological outcome was found in patients treated with TLIF, with Radiographic fusion rates 91.9% at the end of 3 years follow-up. Comparing to our study we also found highly clinical outcome & Radiographic fusion in both TLIF group and PLF group with slightly better radiographic fusion in TLIF group at the end of 36 months.

Kristian Høy et al.⁶ studied 100 patients in Prospective randomized clinical study from November 2003 to November 2008 to analyse outcome, with respect to

functional disability, pain, fusion rate and complications of patients treated with TLIF was compared to instrumented PLF alone in low back pain patients. They concluded that TLIF did not significantly improve functional outcome in patients compared to PLF. Both groups improved significantly in all categories compared to preoperatively. We found similar results in our present study.

Videbaek TS et al.⁷ did Prospective randomized clinical study with a 5 to 9-year follow-up period to analyse the long-term outcome with respect to functional disability, pain and general health of patients treated by means of circumferential lumbar fusion in comparison with those treated by means of instrumented posterolateral lumbar fusion alone from April 1996 to November 1999 a total

of 148 patients were included in study. They found 5 to 9 years after surgery; the circumferentially fused patients had a significantly improved outcome compared with those treated by means of posterolateral fusion. These results emphasize the superiority of circumferential fusion in the complex pathology of the lumbar spine. In our present study we compared TLIF which is a type of circumferential fusion with PLF. Follow up of this study is 5 to 9 years and follow up of our study is 3 years which is shorter in duration and results of our study shows highly significant post-operative improvement in clinical outcome and radiological outcome in both PLF group and TLIF group. However we are assuming superiority in long term results of TLIF over PLF; as TLIF enables stabilization of the deranged motion segment, neural decompression, reconstruction of the disc height, restoration of the sagittal plane translation and rotational alignment better than PLF.

We observed significant reduction in the post-operative VAS score, ODI Score compare to pre-operative scores in both TLIF & PLF but the mean difference between pre-operative and post-operative Coronal angle in PLF group was 11.419 (± 1.675) and in TLIF group it was 8.705 (± 2.058) & The mean difference between pre-operative and post-operative Lumbar Lordosis Angle in PLF group was -12.226 (± 2.412) and in TLIF group it was -10.000 (± 2.156); which shows radiologically TLIF gives better correction in sagittal balance and rotational alignment compare to PLF. Because of this we assume that there will be better clinical and radiological outcome in TLIF group in long term.

This study has several limitations. First this is monocentric retrospective study; secondly our follow up period is short & also the sample size of 140 patients is less to form a definitive conclusion. A randomized controlled trial of sufficient sample size with longer follow up is needed for further evaluation.

5. Conclusions

We conclude that in the short term duration of our study, both the procedures done with proper technique in duly indicated patients shows satisfactory clinical outcome. However, radiologically TLIF patients had better outcome. We expect better outcome in long term with TLIF compared to PLF. In presence of insignificant blood loss, surgical duration and better 360° fusion TLIF is preferred over PLF.

6. Source of Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

7. Conflict of Interest

We declare that there is no conflict of interest.

References

1. Issack PS, Cunningham ME, Pumberger M, Hughes AP, Cammisia FP. Degenerative lumbar spinal stenosis: evaluation and management. *J Am Acad Orthop Surg*. 2012;20(8):527–35.
2. Atlas SJ, Keller RB, Wu YA, Deyo RA, Singer DE. Long-term outcomes of surgical and nonsurgical management of lumbar spinal stenosis: 8 to 10 year results from the maine lumbar spine study. *Phila Pa 1976*. 2005;30:936–43.
3. Bydon M, Macki M, Abt NB. The cost-effectiveness of interbody fusions versus posterolateral fusions in 137 patients with lumbar spondylolisthesis. *Spine J*. 2015;15(3):492–498.
4. Rajae SS, Bae HW, Kanim LEA, Delamarter RB. 1976. Available from: <http://dx.doi.org/10.1097/BRS.0b013e31820cccfb>.
5. Audat Z, Moutasem O, Yousef K, Mohammad B. Comparison of clinical and radiological results of posterolateral fusion, posterior lumbar interbody fusion and transforaminal lumbar interbody fusion techniques in the treatment of degenerative lumbar spine. *Singapore Med J*. 2012;53(3):22434292–22434292.
6. Høy K, Bünger C, Niederman B, Helmig P, Hansen ES, Li H. Transforaminal lumbar interbody fusion (TLIF) versus posterolateral instrumented fusion (PLF) in degenerative lumbar disorders: A randomized clinical trial with 2-year followup. *Eur Spine J*. 2013;22(9):2022–9.
7. Videbaek TS, Christensen FB, Soegaard R, Hansen ES, Høy K, Helmig P, et al. Circumferential fusion improves outcome in comparison with instrumented posterolateral fusion: long-term results of a randomized clinical trial. *Spine (Phila Pa 1976)*. 2006;31(25):2875–80.

Author biography

Uday Phute, HOD

Gajanan Deshmukh, Consultant

Sushant Sopan Tuse, Senior Resident

Cite this article: Phute U, Deshmukh G, Tuse SS. Comparison of early clinico-radiological outcome of posterolateral fusion and transforaminal lumbar interbody fusion techniques. *Indian J Orthop Surg* 2022;8(2):148-153.