

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Pathology and Oncology

Journal homepage: www.ijpo.co.in

Original Research Article

Evaluation of serum adenosine deaminase activity with reference to dyslipidemia in ischemic heart disease

Pampa Reddy B Kollur¹, Raghunandana R², Girish M Desai³, Netravati Sajjan⁴, Sunilkumar Biradar^{5,*}

¹Al-Azhar Medical College and Super Specialty Hospital, Kumaramangalam, Kerala, India

²Sri Siddhartha Institute of Medical Sciences & Research, Tumakuru, Karnataka, India

³Gulbarga Institute of Medical Sciences, Kalaburagi, Karnataka, India

⁴Navodaya Medical College, Raichur, Karnataka, India

⁵Dept. of Microbiology, M R Medical College, Gulbarga, Karnataka, India



ARTICLE INFO

Article history:

Received 08-09-2021

Accepted 07-02-2022

Available online 28-05-2022

Keywords:

Adenosine deaminase

Ischemic heart disease

Atherosclerosis

Lipids

ABSTRACT

Background: Ischemic heart disease is a major health and a modern epidemic which is increasing in India. Dyslipidemia is a major risk factor for atherosclerosis. Serum adenosine deaminase activity is associated with coronary artery disease.

Aims and Objectives: The present work was done to understand the relation of serum adenosine deaminase activity with dyslipidemia in ischaemic heart disease.

Materials and Methods: This cross-sectional study comprised 200 subjects; healthy and myocardial infarction. Serum ADA and lipids were estimated. Obtained data was statistically analysed.

Results: Increase in serum ADA activity correlating with dyslipidemia was found in Myocardial infarction subjects.

Discussion: Increased ADA has role in LDL oxidation, dyslipidemia, foam cell transformation etc. leading to atherosclerosis and ischemic heart disease.

Conclusion: Adenosine deaminase has a crucial role in the evolution of ischemic heart disease through dyslipidemia and coronary artery disease.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Ischemic heart disease is a major health problem. It is the single largest cause of mortality worldwide accounting for 30% of global deaths.^{1,2} An estimated 7.4 million people died from cardiovascular disease in 2015 worldwide and 2.59 million people died from cardiovascular disease in India during 2016.³ Coronary artery disease is considered a modern epidemic and is increasing in India.⁴ Ischemic heart disease is a condition where there is inadequate supply of blood and

oxygen to a portion of myocardium. The most common cause of myocardial ischemia is atherosclerotic disease of epicardial coronary artery disease sufficient to cause a regional reduction in myocardial blood flow and inadequate perfusion of myocardium supplied by involved coronary artery.⁵

Atherosclerosis results from endothelial dysfunction lipoprotein infiltration through arterial endothelium into intima, inflammation and altered reverse cholesterol transport.⁵ The risk factors for cardiovascular disease are cigarette smoking, dyslipidemia, obesity, hypertension, diabetes mellitus, alcohol etc.⁶ Lipoproteins and dyslipidemia rank among the most prominent and modifiable risk factors for

* Corresponding author.

E-mail address: drsnilkb11@gmail.com (S. Biradar).

atherosclerosis.⁷ Dyslipidemia is associated with adenosine deaminase in diseases.⁸ Serum adenosine deaminase activity is associated with coronary artery disease.⁹

The present work was done to understand the relation of serum adenosine deaminase activity with dyslipidemia in ischaemic heart disease.

2. Objectives

1. To compare serum Adenosine deaminase activity, lipid profile & indices in myocardial infarction and healthy subjects.
2. To correlate the serum Adenosine deaminase activity with serum lipid profile in myocardial infarction and healthy subjects.

3. Materials and Methods

This observational cross-sectional study was carried out for a period of 1 year from October 2013 to September 2014 after obtaining institutional ethical committee clearance.

The study subjects were selected by random sampling. They comprised 200 subjects divided into 2 groups – controls and cases. The healthy volunteers were included in the control group and myocardial infarction patients were included in the cases group. The study subjects were selected by the following inclusion and exclusion criteria. Myocardial infarction patient of both sex, between age 30-60 years were included in cases group. Patients with cardiac risk factors, hepatic diseases, rheumatoid arthritis, tuberculosis, recent surgery or injury, endocrine disturbances, nutritional disorders, pregnant women, those on drugs affecting lipid metabolism were excluded.

The equipment used for the study were syringes, BD vacutainers, Biohit autopipettes, Remi centrifuges, ERBA CHEM 7 semi autoanalyser and ROCHE C111 autoanalyser. The reagent kits were obtained from Tulip and Roche Diagnostics company. The material used for the study was a well-structured questionnaire and blood sample. The volunteers were explained about the study and written informed consent was taken from them. A detailed history was taken and through physical clinical examination was done by a cardiologist. Diagnosis was based on changes in ECG and cardiac biomarkers. Overnight fasting 2 ml blood was collected aseptically from the median cubital vein using a 22 gauge needle syringe into a plain vacutainer. Blood was allowed to clot and centrifuged at 4000 RPM for 15 minutes to obtain serum. The obtained serum were subjected to the following biochemical analysis (Figure 1).

4. Results

The data from the above analysis were compiled into two table's i.e. (Tables 1 and 2).

Compares the mean value of variables in healthy volunteers and myocardial infarction patients. The

difference for the age and sex was not statistically significant. Highly significant statistical differences were seen for serum Adenosine deaminase activity, Cholesterol, triglycerides, High density and low density lipoproteins between the study groups.

Figure 1 compares the mean value of variables in healthy volunteers and Myocardial infarction patients. The difference for the age and sex was not statistically significant. Highly significant statistical differences were seen for serum Adenosine deaminase activity, cholesterol, triglycerides, high density and low density lipoproteins.




Fig. 1: Comparison of ADA in healthy and MI subjects




Fig. 2: Comparison of serum lipids in both healthy and MI subjects

Shows the association of serum adenosine deaminase activity with cholesterol, triglycerides, HDL and LDL. The enzyme activity showed statistical significant correlation with serum lipids in myocardial infarction subjects.


Table 1: Comparison of serum adenosine deaminase activity and serum lipids

Parameters	Controls	Cases	P value	Interpretation
ADA (IU/L)	7.4	11.7	0.0014	Highly Significant
TC (mg/dl)	158	180.6	0.019	Highly Significant
TAG (mg / dl)	121.8	176.4	0.0003	Highly Significant
HDL(mg/dl)	46.5	37.82	0.0001	Highly Significant
LDL (mg/dl)	89.3	132.2	0.0009	Highly Significant

ADA = Adenosine deaminase, TC = Total cholesterol, TAG = Triglycerides, HDL = High density lipoproteins, LDL = Low density lipoproteins

Table 2: Correlation of ADA with lipid profile

Parameters	Controls (r value)	Cases (r value)
TC	0.2	0.58
TAG	-0.45	0.72
HDL	0.23	0.55
LDL	0.31	0.58

**Fig. 3:** Correlation of ADA with total cholesterol

5. Discussion

This study was done to understand the relationship between adenosine deaminase activity and dyslipidemia in Ischemic heart disease (Myocardial infarction). In the present study we have compared the levels of adenosine deaminase and serum lipids in healthy subjects and myocardial infarction patients. Also we correlated the adenosine deaminase activity with serum lipids. We found an increase in ADA activity, serum lipids and correlation of ADA with serum lipids in myocardial infarction. The findings of our study were similar to other studies.^{10–20}

Increase in serum ADA levels of Myocardial infarction patients was also seen in works done by A Jyothy¹¹ [Mean ADA levels : 44.07±12.46 in patients and 20.71±5.63 in controls] Asiya¹² [Case 48.28±12.63 U/L, Control 16.39±3.63 U/L], Torrelas¹³ [Case 120±19.59 nmol/ml] and decreased in work done by Chao⁹ [Cases 10.08 ±3.57 U/L, Control 11.71±4.20 U/L]. Increase in serum cholesterol triglycerides, LDL and decrease in HDL in Myocardial subjects has been reported by Ferdous¹⁷ and Ganesh Draupad Ghuge.¹⁸ ADA had shown positive correlation with total Cholesterol (+ 0.2), Triglycerides (+

0.4) and negative correlation with HDL (-0.3) in work done by P Vijaylaxmi.¹⁹ Similar findings were reported by Shashikala Magadi Dasegowda²⁰ for ADA vs TG (0.858), ADA vs LDL (0.953), and ADA vs HDL (-0.604).

Altered cell mediated immunity increases adenosine deaminase, decreases Adenosine and adenosine receptor activation leading to decreased lipolysis, increased lipogenesis and LDL oxidation causing dyslipidemia. Also adenosine deaminase plays a role in foam cell transformation and vascular proliferation. The above actions of adenosine deaminase leads to atherosclerosis and ischemic heart disease.^{21–25}

The strength of this work is standardized protocols, examination by experienced cardiologist and biochemist. The limitations of the work are cross-sectional study design, small sample size, subjects are not sex matched and single measurements.

6. Conclusion

Adenosine deaminase has a crucial role in the evolution of ischemic heart disease through dyslipidemia and coronary artery disease. It should be measured in suspected patients of ischaemic heart disease. Therapeutic strategies against it can be developed for prevention of ischemic heart disease.²⁵

7. Source of Funding

None.

8. Conflict of Interest

None.

Acknowledgments

We would like to thank the Dean, Medical Superintendent, staff, Cardiologist, statistician, Post graduates, technicians, patients and volunteers of our hospital for their cooperation

in helping to carry out the study.

References

- Kumar V, Abbas AK, Aster J. Robbin and Cottran Pathologic Basis of Disease. 10th ed. India: Elsevier; 2020.
- Mohan H. Textbook of Pathology. 8th ed. Delhi, India: Jaypee Brother Medical Publishers; 2019.
- Park K. Park's textbook of Preventive and Social Medicine. 25th ed. India: M/S Banarasi Das Bhanot Publication; 2019. p. 395.
- Suryakanta AH. Community Medicine. 3rd ed. New Delhi: Jaypee Brother Medical Publishers; 2014. p. 529.
- Disorders of Cardiovascular system. In: Longo DL, Kasper DL, Jameson JL, Fauci AS, Hauser SL, Loscalzo J, editors. Harrison's Principles of Internal Medicine. vol. Vol 1. McGraw-Hill Professional; 2011. p. 243.
- The Pathogenesis, prevention and treatment of Atherosclerosis. In: Longo DL, Kasper DL, Jameson JL, Fauci AS, Hauser SL, Loscalzo J, editors. Harrison's Principles of Internal Medicine. vol. Vol 1. McGraw-Hill Professional; 2011. p. 241.
- Epidemiology of Cardiovascular Disease. In: Goldman L, Ausiello DA, Arend W, Armitage JO, Clemmons D, Drazen J, et al., editors. Cecil Medicine. United States: Saunders; 2007.
- Vijayalakshmi P, Ranjan S. Altered levels of Serum ADA in Type 2 Diabetes Mellitus. *Natl J Lab Med*. 2019;8(2):9–11.
- Xuan C, Tian QW, Zhang SY, Li H, Tian TT, Zhao P, et al. Serum Adenosine Deaminase activity and Coronary Artery Disease : A retrospective case control study based on 9929 participants. *Ther Adv Chronic Dis*. 2019;10:1–10. doi:10.1177/2040622319891539.
- Glusti G, Galanti B. Methods of Enzymatic Analysis. In: Adenosine Deaminase. New York: Academic Press; 1974. p. 1092–9.
- Jyothy A, Rani HS, Rao VD, Reddy PP. Serum Adenosine Deaminase Activity in Myocardial Infarction. *Int J Hum Genet*. 2003;3(1):65–7.
- Naaz A. Estimation of Serum Homocysteine, Adenosine Deaminase And Hs-CRP In Myocardial Infarction Patients. *J Biotechnol Biochem*. 2019;5(3):60–4.
- Shatha A, AL-Shammaree SAW, Saadon S. Adenosine Deaminase activity in Iraqi patients with Myocardial Infarction. *Asian J Pharm, Nurs Med Sci*. 2017;2(1):9–14.
- Torrellas Y, Careman M, Aguidar P, Ramas B. Increased Activity in patients with acute Myocardial Infarction. *Latin Am J Hypertens*. 2010;15(2):38–42.
- Surekha RH, Dayasagar RV, Shiva PM, Jyothy A. Serum Adenosine Deaminase activity and C Reactive protein in Unstable Angina. *Indian J Hum Genet*. 2003;9(1):17–20.
- Tang R, Ma CS, Dong JZ, Liu XP, Liu X. Does adenosine deaminase play a key role in coronary artery disease. *Med Arch*. 2006;67(5):374–7.
- Ferdous B, Sultana N, Ahmed S, Khan E, Sultana S & Parvin T. Serum Lipid profile in Ischemic heart disease a cross sectional analytical study Bangladesh. *J Med Biochem*. 2014;7(1):14–6.
- Ghugre GD, Zine R, Mogrekar M. Lipid profile of patients with Ischemic heart disease from rural area of Marathwada region, India. *Int J Biomed Adv Res*. 2012;3(10):767–9.
- Vijaylakshmi P, Supriya. Altered level of adenosine deaminase in type 2 diabetes mellitus. *Natl J Lab Med*. 2019;8(3):9–11.
- Dasegowda S, Jeppu AK, Sushith, Kumar K. Relation of Serum Adenosine Deaminase to Some Risk Factors in Type 2 Diabetes Mellitus. *J Health Sci Res*. 2016;1(1):13–7.
- Bottini FG, Sarfranow K, Bancietal M, Binczak-Kuleta A, Saccucci P, Ciechanowicz A. The association of Adenosine Deaminase with Coronary artery disease : Effect of Gender and Diabetes. *Health*. 2013;15(2):166–9.
- Bottini FG, Banci M, Saccucci P, Lucarini N, Ianniello F, Paradisi G, et al. Coronary artery disease : a study on joint role of Birth weight, Adenosine deaminase and Gender. *Cardiol Res Pract*. 2009;2009:860328. doi:10.4061/2009/860328.
- Koupenova M, Johnston-Cox H, Raia K. Regulation of Atherosclerosis and Associated Risk Factors by Adenosine and Adenosine Receptors. *Curr Atheroscler Rep*. 2012;14(5):460–8.
- Kaul A, Sethi R, Misra MK. Erythrocyte Adenosine Deaminase in Post Myocardial Infarction angina patients. *Indian J Clin Biochem*. 2009;24(1):49–51.
- Kutryb-Zajac B, Mateuszuk L, Zukowska P, Jaszal A, Zabielska MA, Toczek M, et al. Increased activity of vascular adenosine deaminase in atherosclerosis and therapeutic potential of its inhibition. *Cardiovasc Res*. 2016;112(2):590–605.

Author biography

Pampa Reddy B Kollur, Professor and Head

Raghunandana R, Associate Professor

Girish M Desai, Professor and Head

Netravati Sajjan, Assistant Professor

Sunilkumar Biradar, Associate Professor

Cite this article: Kollur PRB, Raghunandana R, Desai GM, Sajjan N, Biradar S. Evaluation of serum adenosine deaminase activity with reference to dyslipidemia in ischemic heart disease. *Indian J Pathol Oncol* 2022;9(2):128–131.