

Content available at: <https://www.ipinnovative.com/open-access-journals>

International Journal of Pharmaceutical Chemistry and Analysis

Journal homepage: <https://www.ijpca.org/>

Original Research Article

The trend of serum sodium and potassium levels among post-surgical trauma among patients admitted in rural tertiary care unit

Avni Bhadja¹, Akshaya N. Shetti^{1,*}, Bhavika Singla², Vaijayanti K. Badhe¹¹Dept. of Anaesthesiology and Critical Care, Rural Medical College, PIMS, Loni, Maharashtra, India²Dept. of Anaesthesiology, Shri Guru Ram Rai Institute of Medical and Health Sciences, Shri Mahant Indires Hospital, Dehradun, Uttarakhand, India

ARTICLE INFO

Article history:

Received 29-10-2021

Accepted 15-11-2021

Available online xx xx xxxx

Keywords:

Critical care

Electrolytes

postoperative

Potassium

Sodium

Trauma

ABSTRACT

Background: Operative trauma affects the physiology of fluid and electrolytes within the body. Magnitude and duration of this response is directly proportional to the severity of the trauma. Inorganic electrolytes like sodium, potassium, chloride and calcium are very important constituents of the body. There are preoperative, intraoperative and postoperative changes in both fluid volume and electrolyte compositions.

Material and Methods: The cases were selected among the patients operated in the Tertiary Care Hospital and then getting admitted to ICU from November 2020 to April 2021. Males and females, more than 18 years of age, got operated in same hospital were included in the study. The latest preoperative serum sodium and potassium levels had been noted. The postoperative serum sodium and potassium levels along with postoperative day had been noted till the patient was in ICU.

Results: There was gradual rise in the serum sodium level till post-operative day -6 and then it started decreasing and attained its normal preoperative value on post-operative day-11. The preoperative sodium level was restored in all cases by the 11th postoperative day.

There was a gradual fall in serum potassium level from post-operative day-1 to post-operative day-6 and then it started increasing and attained pre-operative value by post-operative day-11. So, the response of serum sodium and serum potassium is contrary to each other.

Conclusion: Based on the study results, increased sodium and decreased potassium levels were observed in postoperative cases. Normalization of these two electrolyte are seen postoperative day seven onwards.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Fluid balance regulation, tissue excitability and acid base equilibrium, cellular function and its survival and other manifestations are associated with presence of inorganic electrolytes in our body. These inorganic electrolytes like sodium, potassium, chloride and calcium are very important constituents of the body.¹ There are preoperative, intraoperative and postoperative changes in both fluid volume and electrolyte compositions.² Any operative

trauma leads to series of changes which is referred to as a metabolic response to injury. Magnitude and duration of this response is directly proportional to the severity of the trauma.^{3,4} Operative trauma affect the physiology of fluid and electrolytes within the body. Moore's metabolic care of the surgical patients is a milestone in the field of modern surgery.⁵ Due to surgical intervention the normal physiology gets distorted. This decides the successful post-operative result. Surgical care is based on an understanding of the normal adaptation and to treat it when abnormal.^{6,7} Fluid and electrolyte management has thus been an integral part of care of each and every surgical

* Corresponding author.

E-mail address: aksnsdr@gmail.com (A. N. Shetti).

patient.⁸

Plasma sodium concentrations disorders are very common in the patients who are hospitalised for a long duration of time. Patients who undergo major surgeries are more prone. In peri-operative period there will be major stress response and fluid loss will occur. The previous studies have showed hospital outcomes associated with dysnatraemias.^{9,10} There are small changes like measurement artefact or diurnal variations and large changes usually associated with iatrogenic factors or endocrine response. These can induce osmotic disequilibrium and cerebral dysfunction.




Fig. 1: The change in serum sodium levels in relation with preoperative levels




Fig. 2: The change in serum potassium levels in relation with preoperative levels

2. Materials and Methods

The study was conducted in intensive care unit (ICU) of tertiary care hospital, Loni, Maharashtra, India. The cases were selected among the patients operated in the same hospital and then getting admitted to ICU from 1st of November 2020 to 30th April 2021. The written, verbal and informed consent of patients or his/her relative was taken. The patients including males and females of age more than 18 years undergoing elective or emergency surgery getting admitted in rural critical care unit were included in the

study. The latest preoperative serum sodium and potassium levels were recorded. The postoperative serum sodium and potassium levels along with postoperative day were noted till the patient was in ICU. The diagnosis, type and duration of the surgery had been noted from the records.

3. Results

This study involves the post-operative patients admitted in rural critical care unit from November 2020 to April 2021. Total 111 cases were studied out of which 101 were major surgeries including 72 males and 29 females and 10 were minor surgeries including 2 males and 8 females. These groups were further divided on the basis of age group less than 60 years and more than 60 years as shown in Table 1. The trend of postoperative average serum sodium and potassium is depicted in Table 2.

4. Discussion

In the present study, the average preoperative level of serum sodium was 134.26 meq/L and serum potassium was 4.05 meq/L as shown in Table 2. Preoperative serum electrolyte levels of both serum sodium and potassium levels were within normal limits. In the present study we observed, there was gradual rise in the serum sodium level till post-operative day — 6 and then it started decreasing and attained its normal preoperative value on post-operative day-11. The preoperative sodium level was restored in all cases by the 11th postoperative day as shown in the Figure 1.

There was a gradual fall in serum potassium level from post-operative day-1 to post-operative day-6 and then it started increasing and attained pre-operative value by post-operative day-11 as shown in Figure 2. So, the response of serum potassium and serum potassium is contrary to each other as studied previously.^{11,12}

The hormonal response to surgery leads to release of catecholamines which stimulate hypothalamo-pituitary-adrenal axis and increase in cortisol, aldosterone and ADH. These cause oliguria and retention of sodium and chloride and increase urinary excretion of potassium. This whole mechanism can be explained as production of extensive inflammatory exudates around traumatized tissue. These exudates are rich in water, sodium, chloride and protein all of which necessary for healing process. Therefore, water, sodium and chloride retention occur. Breakdown of protoplasm of injured tissue (especially skeletal muscle) lead to Potassium release, thus causing slight rise of serum potassium, as it is being produced in such a great quantity that it may not be excreted by kidney with sufficient speed.^{13–15} This explains Potassium paradox i.e. increased plasma level despite negative metabolic balance. In our study we did observe there is an increase in serum potassium levels postoperatively for initial two days and gradually reached to normal values by postoperative day 6 to 7.

Table 1: Type of surgery and demographic details

Type of Surgery	Gender	Age	No. of Patients
Major surgery 101	Males (72)	< 60 years	52
		>= 60 years	20
	Females (29)	< 60 years	21
		>= 60 years	8
Minor surgery 10	Males (2)	< 60 years	0
		>= 60 years	2
	Females (8)	< 60 years	7
		>= 60 years	1
Total patients			111

Table 2: Observation of changes in serum sodium and potassium level in post-operative period

Pre-operative day	Total Patients admitted:111	134.26	4.5
Postoperative day	No. of remaining patients out of 111 on day	Average Serum Sodium level	Average Serum Potassium level
Day 1	85	134.72	4.52
Day 2	33	136.27	4.18
Day 3	25	139.08	4.02
Day 4	22	138.79	3.80
Day 5	18	139.25	3.853
Day 6	11	142.09	4.07
Day 7	10	135.74	3.98
Day 8	7	139.6	3.66
Day 9	6	136.45	4.16
Day 10	5	134.22	3.62
Day 11	5	137.74	4.56
Day 12	2	130.5	4.05
Day 13	2	126.5	3.85

5. Limitation of the study

The study was done on limited number of samples (111). We recommend conducting such study in large number of patients. We have not taken into consideration of intravenous fluid administered in perioperative period and not studied other electrolytes.

6. Conclusions

Based on the study results, the sodium levels were seen in increasing trend in the post-surgical patients as compared to the preoperative levels. The potassium levels were of decreasing trend in postoperative levels. Normalization of these serum sodium and potassium levels are seen postoperative day seven onwards. One should carefully monitor the electrolytes postoperatively and treat accordingly. The type of fluid selection in postoperative period is an important aspect to avoid life threatening complications related to electrolyte imbalances.

7. Source of Funding

None.

8. Conflict of Interest

None.

References

- Terry J. The major electrolytes: sodium, potassium, and chloride. *J Intraven Nurs.* 1994;17(5):240–7.
- Piper GL, Kaplan LJ. Fluid and electrolyte management for the surgical patient. *Surg Clin North Am.* 2012;92(2):189–205. doi:10.1016/j.suc.2012.01.004.
- Minto G, Biccard B. Assessment of the high-risk perioperative patient. *BJA Educ.* 2014;14(1):12–7. doi:10.1093/bjaceaccp/mkt020.
- Burton D, Nicholson G, Hall G. Endocrine and metabolic response to surgery. *Contin Educ Anaesth Crit Care Pain.* 2004;4(5):144–51.
- Moore FD. Metabolic care of surgical patient; 1959. p. 394.
- Surwit EA, Tam TY. Postoperative Care Glob. *Libr Women's med.* 2008;doi:10.3843/GLOWM.10036.
- Piper GL, Kaplan LJ. Fluid and electrolyte management for the surgical patient. *Surg Clin North Am.* 2012;92(2):189–205.
- soni N. British consensus guidelines on intravenous fluid therapy for adult surgical patients-GIFTASUP. *Anaesthesia.* 2008;64(3):235–8. doi:10.1111/j.1365-2044.2009.05886_1.x.
- Funk GC, Lindner G, Druml W. Incidence and prognosis of dysnatremias present on ICU admission. *Intensive Care Med.* 2010;36(2):304–15. doi:10.1007/s00134-009-1692-0.
- Wald R, Jaber BL, Price LL, Upadhyay A, Madias NE. Impact of hospital-associated hyponatremia on selected outcomes. *Arch Intern Med.* 2010;170(3):294–302. doi:10.1001/archinternmed.2009.513.
- Steele A, Gowrishankar M, Abrahamson S, Mazer CD, Feldman RD, Halperin ML. Postoperative hyponatremia despite near isotonic saline infusion: a phenomenon of desalination. *Ann Intern Med.* 1997;126(1):20–5. doi:10.7326/0003-4819-126-1-199701010-00003.

12. Luckey AE, Parsa CJ. Fluid and electrolytes in the aged. *Arch Surg.* 2003;138(10):1055–60. doi:10.1001/archsurg.138.10.1055.
13. Finnerty CC, Mabvuure NT, Ali A, Kozar RA, Herndon DN. The surgically induced stress response. *J Parenter Enteral Nutr.* 2013;37(50):21–9. doi:10.1177/0148607113496117.
14. Şimşek T, Şimşek HU, Cantürk NZ. Response to trauma and metabolic changes: posttraumatic metabolism. *Turkish J Surg.* 2014;30(3):153–62. doi:10.5152/UCD.2014.2653.
15. Hiner HH, Suki WN. Fluid and electrolyte disorders in surgical patient. *Ther Renal Dis Related Disord.* 1991;2:263–76. doi:10.1007/978-1-4613-0689-4_16.

Akshaya N. Shetti, Professor

Bhavika Singla, Assistant Professor

Vaijayanti K. Badhe, Professor and HOD

Cite this article: Bhadja A, Shetti AN, Singla B, Badhe VK. The trend of serum sodium and potassium levels among post-surgical trauma among patients admitted in rural tertiary care unit. *Int J Pharm Chem Anal* 2021;8(4):167-170.

Author biography

Avni Bhadja, Resident