

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Indian Journal of Conservative and Endodontics

Journal homepage: <https://www.ijce.in/>

## Original Research Article

# Comparative evaluation of antimicrobial efficacy of natural extracts and sodium hypochlorite against persistent root canal pathogens: An in-vitro study

S Anitha Rao<sup>1</sup>, Avula Sindhuja<sup>1,\*</sup><sup>1</sup>Dept. of Conservative Dentistry And Endodontics, Mamata Dental College, Khammam, Telangana, India

## ARTICLE INFO

## Article history:

Received 28-02-2022

Accepted 02-03-2022

Available online 31-03-2022

## Keywords:

E faecalis

Carica papaya

Moringa oleifera

## ABSTRACT

**Background:** The purpose of the present study was to evaluate the antimicrobial efficacy of natural extracts and Sodium hypochlorite against *E. faecalis*.

**Aim:** The present study aimed to evaluate the antimicrobial efficacy of natural extracts (*Moringa oleifera* and *Carica papaya*) and Sodium hypochlorite against *E. faecalis*.

**Materials and Methods:** *Carica papaya* and *Moringa oleifera* leaf extracts were obtained. 30 extracted incisor teeth were taken and access cavities were prepared, canals were prepared using the crown-down technique. *E. faecalis* suspension was placed into root canals and was incubated. Teeth were divided into three groups based on irrigating solutions: 1. 3% NaOCl, 2. *Moringa* leaf extract, 3. *Papaya* leaf extract. Dental shavings were collected on BHI agar and were incubated. Colonies of *E. faecalis* were counted.

**Results:** Differences were found in colony count for different irrigating solutions. p-value < 0.05 is considered statistically significant.

**Conclusions:** *Moringa oleifera* has good antimicrobial properties compared to Sodium hypochlorite and *Carica papaya*.

**Relevance for patients:** The results should be considered for further research in minimizing the use of chemical substances and increasing the use of effective herbal extracts in root canal treatment.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprint@ipinnovative.com](mailto:reprint@ipinnovative.com)

## 1. Introduction

The main objective of root canal treatment is to eliminate the microorganisms from the complex root canal system. The persistence of the microorganisms leads to failure of the root canal treatment.<sup>1</sup> *Enterococcus faecalis* is a gram-positive, facultative anaerobic coccus that can survive in various environments with different pH, and is a pathogen that is seen in persistent peri-radicular lesions even after root canal treatment.<sup>2</sup>

Sodium hypochlorite is the most commonly used endodontic irrigant, has excellent tissue dissolution and antimicrobial properties. It has undesirable effects like

tissue toxicity, pain, swelling due to accidental injection beyond the apex and it leads to the development of secondary infection and paresthesia.<sup>3</sup>

*Carica papaya* commonly known as the Papaya tree has good medicinal properties in its fruits and leaves. The young Papaya leaves contain papain which is a proteolytic enzyme and is used to soften the meat. Leaves of the Papaya tree contain alkaloids, saponins, tannins, glycosides which have both antibacterial and antiviral properties. Flavonoids present in the leaves are known as 'biological response modifiers' as it protects against oxidative and free-radical damage.<sup>4,5</sup>

*Moringa oleifera* commonly known as Moringa has curative properties. The leaves contain alkaloids, flavonoids, saponins, triterpenoids that contribute to the antibacterial

\* Corresponding author.

E-mail address: [sindhujaavula95@gmail.com](mailto:sindhujaavula95@gmail.com) (A. Sindhuja).

properties.<sup>6,7</sup>

The objective of the present study was to evaluate the antimicrobial efficacy of the *Moringa oleifera* and *Carica papaya* leaf extract against *E. faecalis*, a persistent root canal pathogen. The outcomes of the research may help in giving scope for further research on the herbal products uses in endodontics thus, eliminating the undesirable effects caused by chemical irrigants.

## 2. Materials and Methods

*Moringa oleifera* leaf extract, *Carica papaya* leaf extract, 3% Sodium hypochlorite, *E. faecalis*, BHI broth, and agar were used in this study.

### 2.1. Preparation of leaf extracts

*Carica papaya* and *Moringa oleifera* leaves were collected and were thoroughly washed in distilled water and were dried in daylight. The dried leaves were ground into fine powder. 40 grams of Papaya leaf powder was taken in a cloth pouch and 250 ml of Ethanol was taken (Figure 1). Both the ingredients were placed in the Soxhlet apparatus and the temperature was maintained at 60°C for four days. The resultant liquid was filtered using Whatman no.1 filter paper and was concentrated in an evaporator at 40°C to obtain a concentration of 30 mg/ml. Similarly, *Moringa* leaf extract was prepared with the same concentration as mentioned above.



**Fig. 1:** Soxhelt apparatus

### 2.2. Preparation of the specimens

Thirty extracted human incisor teeth with a single canal were collected and stored in the normal saline solution until use (Figure 2). Access cavities were prepared on the lingual surfaces of the teeth with diamond points using a high-speed airtor handpiece (Figure 3). Canal patency was confirmed using the #15k file and working length was determined (Figure 4). The root canals were prepared using the crown-down technique with Protaper hand files. The canals were irrigated with 3% NaOCl in between instrumentation. Then the root apices of the teeth were sealed with nail varnish and then sterilized in an autoclave (Figure 5).



**Fig. 2:** Extracted incisor teeth




**Fig. 3:** Access cavity preparation in incisors

**Table 1:** Antimicrobial efficacy of the irrigants in terms of colony-forming units


Extracts	Mean ±SD	SE	95% CI for Mean	
			Lower Bound	Upper Bound
Sodium Hypochlorite	3.30 ± 1.70	0.54	2.08	4.52
Moringa oleifera	1.10 ± 0.74	0.23	0.57	1.63
Carica papaya	5.60 ± 2.07	0.65	4.12	7.08

**Table 2:** Pairwise comparisons of the three irrigants for antimicrobial efficacy in terms of CFU


Extracts	Mean ±SD	Mean ranks	U-value	Z-value	p-value
Sodium Hypochlorite	3.30 ± 1.70	14.65			
Moringa oleifera	1.10 ± 0.74	6.35	8.50	-3.1371	0.0017*
Sodium Hypochlorite	3.30 ± 1.70	7.15			
Carica papaya	5.60 ± 2.07	13.85	16.50	-2.5324	0.0113*
Moringa oleifera	1.10 ± 0.74	5.50			
Carica papaya	5.60 ± 2.07	15.50	0.00	-3.7796	0.0002*



**Fig. 4:** Confirmation of canal patency



**Fig. 6:** Turbid suspension of E. Faecalis



**Fig. 5:** Rootapices sealed with varnish

**2.3. Preparation of Enterococcus faecalis suspension and inoculation into the specimens**

A turbid suspension of E. faecalis was obtained by growing E. faecalis cells in the BHI (Brain heart infusion) broth for 6 hours at 36.5°C (Figure 6). This suspension was inoculated into the root canals and the teeth were incubated for 24 hours at 36.5°C (Figure 7).

**2.4. Grouping based on the irrigants and formation of E faecalis colonies**

Teeth were divided into three groups (n=10).

Group 1: 3% NaOCl.


Group 2: 30 mg/ml Moringa leaf extract.



**Fig. 7:** Inoculation into teeth



**Fig. 8:** Collection of shavings by drilling



**Fig. 9:** Comparison of three extracts with antimicrobial efficacy in terms of CFU.

Group 3: 30 mg/ml Papaya leaf extract.

In all three groups, canals were irrigated with 2 ml of the respective irrigant solutions using a sterile needle for one minute. Then a cavity was made at the middle third of the proximal surface of the root into the root canal using a sterile round bur. The shavings were collected on the BHI agar plate and then incubated for 48 hours at 36.5°C (Figure 8).

*E. faecalis* were expressed as colony-forming units (CFU). The obtained data were statistically analyzed using the Mann-Whitney U test.  $p$ -value < 0.05 is considered statistically significant.

### 3. Results

Table 1 shows the antimicrobial efficacy in terms of the colony-forming units for all three irrigants individually. It shows that the colony-forming units with *Moringa oleifera* leaf extract as irrigant is low compared to the other two irrigating solutions indicating its good antimicrobial potential against *E. faecalis*. The second lesser colony count is seen with 3% NaOCl as an irrigant, with the highest number of colonies for papaya leaf extract.

Table 2 Shows the pair-wise comparisons of the irrigants (NaOCl and *Moringa*, NaOCl and Papaya, *Moringa* and Papaya). It shows that the difference in antimicrobial efficacy of *Moringa* and Papaya is more statistically significant compared to other pairs.

### 4. Discussion

Sodium hypochlorite has been used as the most effective root canal irrigant due to its high antimicrobial efficacy and the ability to eliminate persistent endodontic microorganisms. However, it has few disadvantages like the inability to remove the smear layer, unpleasant taste, allergic potential, and toxicity.<sup>5</sup> According to Grigoratos D et al., there was a decrease in elastic and flexural strengths of dentin after immersion in NaOCl for 2 hours or when used for a longer duration.<sup>8</sup>

*Moringa oleifera* leaves contain phytochemicals that have anti-cancer, anti-inflammatory, and antibacterial properties. According to Lucio Arevalo Hijar et al, *Moringa oleifera* demonstrated antimicrobial effect against *E. faecalis* without any toxicity when used at low concentrations.<sup>9</sup>

*Carica papaya* contains papain, alkaloids that have toxic properties against microbes that can effectively kill bacteria and viruses. According to Snigdha Shubham et al, the antimicrobial efficacy of crude Papaya extract is comparable to Sodium hypochlorite and CHX.<sup>5</sup>

The antimicrobial properties of the plant extracts are due to the presence of components known as Bioactive compounds like alkaloids, flavonoids, saponins. Alkaloids are the main group of bioactive components present in the leaves of the *Moringa* and *Papaya* tree.<sup>10,11</sup> According

to Pedro et al, the leaf extracts of plants contain the components that have the highest antibacterial and antiviral properties.<sup>5</sup>

It has been demonstrated that the solvents used for the extraction also influence the antibacterial activity of herbs. Various solvents like distilled water, ethanol, phenol can be used for the extraction of bioactive components. Ethanolic extracts showed more antimicrobial potential compared to other extracts. According to Nwah NS et al., when aqueous and ethanolic extracts of papaya leaves were compared, the ethanolic extracts showed a significant antimicrobial effect whereas aqueous extract showed no antibacterial effect.<sup>12</sup>

In this study, the order of antimicrobial efficacy is Moringa oleifera > Sodium hypochlorite > Carica papaya. The clinicians must understand the relationship of a microorganism in the development of endodontic disease to plan for better treatment. The use of herbal alternatives as root canal irrigants is proved to be advantageous considering the several undesirable characteristics of chemical irrigants. Further studies are required to evaluate other desirable properties of herbal irrigants.

## 5. Conflict of Interest

The authors declare that there is no conflict of interest.

## 6. Source of Funding

None.

## References

1. Neera, Kundabala M, Shenoy, Shalini, Kamath, Sarita, et al. Evaluation of antimicrobial efficacy of 0.5% IKI, 3% NaOCl, and 0.2% CHX, when used alone and in combination as intracanal irrigants against *Enterococcus faecalis*. *Endodontology*. 2009;21(2):6–16.
2. Stuart CH, Schwartz SA, Beeson TJ, Owatz CB. *Enterococcus faecalis*: Its role in root canal treatment failure and current concepts in retreatment. *J Endod*. 2006;32(2):93–8.
3. Mohammadi Z. Sodium hypochlorite in endodontics: an update review. *Int Dent J*. 2008;58(6):329–41.
4. Wilda Y, Satryani F, Maemonah S, Badriyah, Christyaningsih J. The inhibitory effect of *Carica papaya* cv. Thailand leaf extract to the growth of *Enterococcus faecalis* in-vitro. *Sch J Dent Sci*. 2017;4(6):262–6.
5. Shubham S, Samant PS, Khanal R, Gautham V, Biring OJS, Arora C, et al. Ravish Mishra, Laxmi Kandel. Papaya extract as a new endodontic irrigant. *J Dent Med Sci*. 2018;17(9):23–8.
6. Sopandani P, Iskandar BO, Ariwibowo T, Djamil MS. Antibacterial effects of moringa oleifera leaf extract against *Enterococcus faecalis* in-vitro. *Sci Dent J*. 2020;4:16–20.
7. Elgamily H, Moussa A, Elboraey A, El-Sayed H, Al-Moghazy M, Abdalla A, et al. Microbiological Assessment of Moringa Oleifera Extracts and Its Incorporation in Novel Dental Remedies against Some Oral Pathogens. Open Access Maced. *J Med Sci*. 2016;4(4):585–90.
8. Grigoratos D, Knowles J, Ng YL, Gulabivala K. Effect of exposing dentine to sodium hypochlorite and calcium hydroxide on its flexural strength and elastic modulus. *Int Endod J*. 2001;34(2):113–9. doi:10.1046/j.1365-2591.2001.00356.x.
9. Arévalo-Hfjar L, Aguilar-Luis M, Gracia NC, Valle-Mendoza JD. Antibacterial and Cytotoxic Effects of Moringa oleifera (Moringa) and *Azadirachta indica* (Neem) Methanolic Extracts against Strains of *Enterococcus faecalis*. *Int J Dent*. 2018;p. 1071676. doi:10.1155/2018/1071676.
10. Yusha'u M, Onuorah FC, Murtala. In-vitro sensitivity pattern of some urinary tract isolates to *Carica papaya* extracts. *Bayero J Pure Appl*. 2009;2(2):75–8.
11. Jimenez MV, Almatrafi MM, Fernandez ML. Bioactive components in Moringa oleifera leaves protect against Chronic diseases. *Antioxidants (Basel)*. 2017;16(4):91. doi:10.3390/antiox6040091.
12. Awah NS. Antibacterial Activities of the Aqueous and Ethanolic Extracts of the Male and Female *Carica papaya* Leaves on Some Pathogenic Bacteria. *Bioeng Biosci*. 2017;5(2):25–9. doi:10.13189/bb.2017.050201.

## Author biography

**S Anitha Rao**, Professor and HOD

**Avula Sindhuja**, 3rd Year MDS Student

**Cite this article:** Rao SA, Sindhuja A. Comparative evaluation of antimicrobial efficacy of natural extracts and sodium hypochlorite against persistent root canal pathogens: An in-vitro study. *IP Indian J Conserv Endod* 2022;7(1):11-15.