

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Comprehensive and Advanced Pharmacology

Journal homepage: <https://www.ijcap.in/>

Original Research Article

Self medication among under-graduate students in IGIMS, Patna - An observational & prospective study

Shambhu Kumar Yadav, Saajid Hameed^{1,*}¹Dept. of Pharmacology, Indira Gandhi Institute of Medical Sciences, Patna, Bihar, India

ARTICLE INFO

Article history:

Received 25-11-2021

Accepted 04-12-2021

Available online 21-12-2021

Keywords:

SelfMedication

Students

Questionnaire

Drugs

Prescription

ABSTRACT

Background: Inapt medical use is a major problem to ensure effective and safe treatment. Most common health diseases are treated by the people themselves without medical supervision, commonly referred to as self-medication (SM). However, it is also known that responsible SM must be accompanied by relevant health information. It has become a serious problem that raises concerns about misdiagnosis and drug reactions as well.

Aim: To determine the pattern of SM procedures, to identify common diseases and common drugs used, SM causes and to examine the relationship between the level of medical education and SM practices among undergraduate medical students.

Materials and Methods: A structured and validated questionnaire was used for the study to collect information regarding age, gender, awareness of SM practice, type of drugs self-medicated and source of information. Students were also interviewed to check their knowledge and attitude towards self-medication. Results obtained from this study were presented in tabular form and data were interpreted by using Microsoft Excel 365 software.

Results: Out of 387, 365 students (94.32%) students accepted self-medication practice. Most of the self-medicators used the medicines for fever (24.11%) followed by common cold (17.26%) and headache (16.71%). Paracetamol was used by 51.78% of students. Most of the students took self-medications for quick relief and for minor ailments.

Conclusion: our study shows that SM is most prevalent in medical students of North India. Schedule H drugs have also been used without a prescription. This highlights the need for improved drug control.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Inapt medical use is a major problem to ensure effective and safe treatment. Most common health diseases are treated by the people themselves without medical supervision, commonly referred to as self-medication (SM). SM is a human behavior where; a person uses drugs to treat symptoms or debilitating diseases that are found to have the potential to do good and harm as it involves the use of drugs. It is widely practiced worldwide for people in

urban and rural areas including developing countries such as India because many drugs are given over the counter without prescription and offer an alternative that costs less people.¹ It may be due to a number of factors such as socioeconomic status, lifestyle, drug-free access, and greater availability of available medical products in developing countries.² It is an important issue regarding the health of the individual. The practice of Self-Medication should be based on false medical knowledge to avoid unnecessary drug use, which can also lead to loss of resources, increase resistance to viruses and can lead to serious health risks such as long-term suffering, drug reactions and drug dependence.

* Corresponding author.

E-mail address: saajid36@gmail.com (S. Hameed).

Self-medication is now considered part of self-care.³ Conversely, when properly applied, it can be used effectively to treat minor ailments, save time, and save lives. The World Health Organization (WHO) has also suggested that proper Self-Medication can help prevent and treat diseases that do not require consultation and may provide a cheaper alternative to treating common ailments. However, it is also known that responsible SM must be accompanied by relevant health information.⁴

Unregistered drugs among young people, especially students, are being abused as a result of media exposure and advertising. It has become a serious problem that raises concerns about misdiagnosis and drug reactions as well. As future medical professionals, SM has a special impact on medical students. The prevalence of SM was found to vary among medical students from different countries in previous studies⁵⁻⁹ conducted to obtain an SM procedure among medical students. The most common reasons for SM reported in previous study were previous technologies,⁸⁻¹¹ lack of consultation with doctors,^{12,13} rapid relief,¹⁴ and time saving.¹⁵ As most of the studies were conducted in countries other than India, the pattern of SM practices in our country remained unknown.

With this in mind, the current study was designed to determine the pattern of SM procedures, to identify common diseases and common drugs used, SM causes and to examine the relationship between the level of medical education and SM practices among undergraduate medical students.

2. Materials and Methods

This study was conducted at IGIMS Patna, after approval by institutional ethics committee of IGIMS, Patna (Bihar).

2.1. Sample size - 387

2.2. Study design

Observational & Prospective Study

2.3. Study duration

3 Months from January 2020 to March 2020

2.4. Source of data

Medical College, IGIMS Patna, (Bihar)

2.5. Materials

Answers given by Under-Graduate students to the Questionnaire on Self-Medication

2.6. Methodology

A structured and validated questionnaire was used for the study to collect information regarding age, gender,

awareness of SM practice, type of drugs self-medicated and source of information.

Prior to administrating the questionnaire, the students were addressed regarding the purpose and process of data collection. They were informed that data collected would be anonymous and their participation would be voluntary. Questionnaires were distributed among the participants after taking informed consent. Students were also interviewed to check their knowledge and attitude towards self-medication.

Results obtained from this study were presented in tabular form and data were interpreted by using Microsoft Excel 365 software.

FORMAT	
Questionnaire on Self-Medication	
Name:	Age:
Sex:	Session:
Mobile No.:	E-Mail:
1. Indications for self-medication:	
2. Drugs used for self-medication:	
3. Source of drug information:	
4. Reasons for self-medication. Choose from given options:	
a. Quick relief	c. Economical
b. Minor ailments	d. Self-Diagnosis & Previous expertise
5. Source for drug procurement Choose from given options:	
a. Pharmacy	c. Unused medicines at home
b. Batchmates/Seniors	d. Free Physician Sample
6. Do you know the term "OTC (Over the Counter) Drugs"	Yes/No
7. Do you know the recommended dose & frequency of the drug	Yes/No
8. Do you know common adverse effect of the drug	Yes/No

3. Results

Table 1: Comparison of different indication for self-medication

Indications	Number of Students	Percentage of Students
Fever	88	24.11
Common Cold	63	17.26
Headache	61	16.71
GI upset	40	10.96
Diarrhea	37	10.14
Constipation	26	7.12
Nausea & Vomiting	23	6.30
Wound	16	4.38
Sleep disorder	11	3.01
Total	365	100.00

4. Discussion

SM is the use of drugs by individuals alone without competent medical guidance. In developing countries such as India, many episodes are handled by SM due to the easy availability of prescription drugs. It is a major obstacle to ensuring the safe and effective use of drugs. It may not be possible without full knowledge even though it is common practice these days especially for undergraduate students.^{5-11,16,17}

Out of 387, 365 students (94.32%) students accepted self-medication practice. However, various studies have

Table 2: Comparison of drugs used for self-medication

Drugs	Number of Students	Percentage of Students N=365
Paracetamol	189	51.78
Levocetirizine	51	13.97
Cetirizine	17	4.66
Montelukast	42	11.51
Omeprazole	23	6.30
Pantoprazole	13	3.56
Ranitidine	09	2.47
Dicyclomine	21	5.75
Norfloxacin	39	10.68
Tinidazole	39	10.68
Metronidazole	23	6.30
Liquid paraffin	15	4.11
Sodium pico-sulfate	15	4.11
Milk of magnesia	15	4.11
Polyethylene glycol	16	4.38
Probiotics	18	4.93
Ondansetron	12	3.29
Domperidone	09	2.47
Metoclopramide	07	1.92
Wound	16	4.38
Azithromycin	47	12.88
Amoxicillin	31	8.49
Amoxicillin + Clavulanate	23	6.30
Cefixime	13	3.56
Ciprofloxacin	09	2.47
Levofloxacin	03	0.82
Clonazepam	09	2.47
Zopiclone	02	0.55

Table 3: Comparison of different sources of drug information for self-medication

Sources	Number of Students	Percentage of Students
Book	105	28.77
Internet	98	26.85
Lecture	55	15.07
Seniors	41	11.23
Colleague	35	9.59
Family	31	8.49
Total	365	100.00

Table 4: Comparison of Different Reason for Self-Medication

Reasons	Number of Students	Percentage of Students
Quick relief	164	44.93
Economical	77	21.10
Minor Ailment	112	30.68
Self Diagnosis & Previous Expertise	12	3.29
Total	365	100.00

Table 5: Comparison of Different Sources of Drug Procurement for Self-Medication

Sources	Number of Students	Percentage of Students
Pharmacy	216	59.18
Unused medicines at home	67	18.36
Batchmates/Seniors	49	13.42
Free Physician Sample	33	9.04
Total	365	100.00

Table 6: Knowledge regarding self-medication among undergraduate students

Knowledge domain	Number of Students having Adequate Knowledge (%)	Number of Students having Inadequate Knowledge (%)
Over the counter drugs	247 (67.67%)	118 (32.33%)
Recommended dose & frequency of the drug	296 (81.10%)	69 (18.90%)
Adverse effect of the drug	208 (56.99%)	157 (43.01%)

reported different distribution figures ranging from 43.2 to 91%.^{8–12,16–19} It is very difficult to compare the increase in diversity of education with the current study due to different demographic characteristics, different approaches, and different socio-economic status.

Among the self-medicators, the majority followed allopathic system of medicine followed by ayurvedic and homeopathic system of medicine which is in conformity with earlier studies,^{12–14} which might be due to easy access of allopathic medicines. The reason attributed to the use of other system of medicines have been stated as practice from childhood, earlier prescriptions available for same ailments, easy accessibility, belief in those systems.

Most of the self-medicators used the medicines for fever (24.11%) followed by common cold (17.26%) and headache (16.71%). Gastrointestinal problem were indications for 34.52% of total indications. The same was reported in other studies,^{7–12,14–17,20–22} although it differs from a study conducted in South India that revealed cough and cold as the most common cause.¹³

Paracetamol was used by 51.78% of students. Various researchers^{7,9–13,15–17,19–22} have identified antipyretics and analgesics as the most commonly used SM drugs, consistent with the current study and, research conducted a medical college in West Bengal reported that antibiotics are the most widely used drugs.¹⁸ 135.52% students used antimicrobial agent without prescription. This highlights the poor regulation of sale of schedule H drug.

Most of the students have procured information about self medicated drugs from books (28.77%), internet (26.85%) and lecture (15.07%) while rest of the students

have procured it from seniors, colleague and family. This is similar to previous studies.^{7,9,13} In contrast, some studies have identified a source of information such as decision-making,¹⁰ adults,^{16,20} family and friends,^{11,15} and previous instructions.^{12,14,17,21}

Most of the students took self-medications for quick relief and for minor ailments. The findings are similar to those found in the literature,^{16,20,21} however some employees have reported previous experiences,^{8–11} lack of time to consult a doctor,^{12,13} immediate relief,¹⁴ and time savings¹⁵ as common reason.

Pharmacy was the most common source of drug purchases (59.18%). The free physician sample was used by 9.04% of students. This was similar to the results of previous studies.^{9–11,16,20} The reason could be easy availability and previous treatment for illness by medication obtained from a pharmacy. Although students have easy access to doctors, but a complex consultation process, getting prescription drugs and going to the pharmacy increases the practice of SM.

Our findings regarding non-pharmacological awareness are consistent with the findings of studies conducted in Maharashtra and Jammu.^{7,12} Findings from various studies^{7,11} regarding knowledge on dosing are similar to the current study. The majority of students regardless of the academic year reported that they were aware of the side effects of their medications, similar to other studies.^{7,11–13,15,19,20,22}

5. Conclusion

Therefore, our study shows that SM is most prevalent in medical students of North India. Schedule H drugs have also been used without a prescription. This underscores the need for improved drug control. Although it is difficult to eliminate SM, various measures can be taken to discourage such practices. If no action is taken, the risk of drug interaction and side effects may increase. With this in mind, an awareness program should be developed to educate students about the various aspects of SM. This will raise awareness of students about SM drug abuse and will also lead to the development of a variety of health education strategies, which are needed to educate students and the community at large.

6. Acknowledgement

We are thankful to the healthcare workers (faculty members) of IGIMS, Patna for their support.

7. Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

8. Source of Funding

None.

References

- Hussain S, Malik F, Hameed A, Riaz H. Exploring health seeking behavior, medicine use and self-medication in rural and urban Pakistan. *South Med Rev.* 2010;3(2):32–4.
- World Health Organization. The Role of pharmacist in Health Care System; 1998. [Last accessed on 2013 Jan 29]. Available from: <http://www.apps.who.int/medicinedocs/en/d/Jwhozip32e>.
- Hughes CM, McElnay JC, Fleming GF. Benefits and risks of self-medication. *Drug Saf.* 2001;24(14):1027–37. doi:10.2165/00002018-200124140-00002.
- World Health Organization (WHO) Guidelines for the regulatory assessment of Medicinal Products for use in self-medication. [Last accessed on 2013 Feb 5]. Available from: <http://www.apps.who.int/medicinedocs/en/d/Js2218e/>.
- Zafar SN, Syed R, Waqar S, Zubairi AJ, Waqar T, Shaikh M, et al. Self-medication amongst University Students of Karachi: Prevalence, Knowledge and Attitudes. *J Pak Med Assn.* 2008;58(4):214–7.
- Klemenc-Ketis Z, Hladnik Z, Kersnik J. Self-medication among healthcare and non-healthcare students at University of Ljubljana. *Med Princ Pract.* 2010;19(5):395–401. doi:10.1159/000316380.
- Sontakke SD, Bajait CS, Pimpalkhute SA, Jaiswal KM, Jaiswal SR. Comparative study of evaluation of self-medication practices in first and third year medical students. *Int J Biol Med Res.* 2011;2(2):561–4.
- James H, Handu SS, Khaja KAA, Ootom S, and RPS. Evaluation of the knowledge, attitude, and practice of self-medication among first-year medical students. *Med Princ Pract.* 2006;15(4):270–5. doi:10.1159/000092989.
- Abay SM, Amelo W. Assessment of self-medication practices among medical, pharmacy, and health science students in Gondhar University. *J Young Pharm.* 2010;2(3):306–10. doi:10.4103/0975-1483.66798.
- Gutema GB, Gadisa DA, Kidanemariam ZA, Berhe DF, Berhe AH, Hadera MG, et al. Self-medication practices among health sciences students: The case of Mekelle University. *J Appl Pharm Sci.* 2011;1(10):183–9.
- Patel PM, Prajapati AK, Ganguly B, Gajjar BM. Study on impact of Pharmacology teaching on knowledge, attitude and practice on self-medication among medical students. *Int J Med Sci Public Health.* 2013;2(2):181–6.
- Kumari R, Kiran, Kumar D, Bahl R, Gupta R. Study of knowledge and practices of self-medication among medical students at Jammu. *J Med Sci.* 2012;15(2):141–4. doi:10.33883/jms.v15i2.252.
- Gaikwad NR, Patil AB, Khan TA. Comparative evaluation of knowledge, attitude and practice of self-medication among first and second year medical students. *J Datta Meghe Inst Med Sci Univ.* 2010;5:157–62.
- Pandya RN, Jhaveri KS, Vyas FI, Patel VJ. Prevalence, pattern and perceptions of self-medication in medical students. *Int J Basic Clin Pharmacol.* 2013;2(3):275–80.
- Badiger S, Kundapur R, Jain A, Kumar A, Patanashetty S, Thakolkaran N, et al. Self-medication patterns among medical students in South India. *Australas Med J.* 2012;5(4):217–20. doi:10.4066/AMJ.2012.1007.
- Girish HO, Divya HM, Prabhakaran S, Venugopalan PP, Koppad R. A cross-sectional study on self medication pattern among medical students at Kannur, North Kerala. *J Evol Med Dent Sci.* 2013;2(45):8693–700.
- Kumar N, Kanchan T, Unnikrishnan, Rekha T, Mithra P, Kulkarni V, et al. Perceptions and practices of self-medication among medical students in coastal South India. *PLoS One.* 2013;8(8):e72247. doi:10.1371/journal.pone.0072247.
- Banerjee I, Bhadury T. Self-medication practice among undergraduate medical students in a tertiary care medical college, West Bengal. *J Postgrad Med.* 2012;58(2):127–31. doi:10.4103/0022-3859.97175.
- Ezz NFE, Ez-Elarab H. Knowledge, attitude and practice of medical students towards self medication at Ain Shams University, Egypt. *J*

- Prev Med Hyg.* 2011;52(4):196–200.
20. Mumtaz Y, Jahangeer SM, Mujtaba T, Zafar S, Adnan S. Self-medication among university students of Karachi. *J Liaquat Univ Med Health Sci.* 2011;10(3):102–5.
21. Bashir MS, Bansod KA, Khade A, Konnoju M, Rani U, Vadala KK, et al. Self-medication A comparative study between 2nd and 3rd year medical students. *Int J Basic App Med Sci.* 2013;3:1–7.
22. Shukla AK, Anand M, Chugh Y, Sharma A, Yadav VS, Charausia RC, et al. Self-medication pattern among medical students in MLN medical college, Allahabad. *Indian J Pharmacol.* 2008;40(S2):S61–2.

Author biography

Shambhu Kumar Yadav, Ex-Senior Resident

Saajid Hameed, Senior Resident

Cite this article: Yadav SK, Hameed S. Self medication among under-graduate students in IGIMS, Patna - An observational & prospective study. *IP Int J Comprehensive Adv Pharmacol* 2021;6(4):185-189.