

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP International Journal of Medical Paediatrics and Oncology

Journal homepage: <https://www.ijmpo.com/>

Original Research Article

Study of prevalence of anaemia and haematological parameters in children below 12 years

Shivendra Varshney¹, Nilesh T Shah^{1,*}, Umang Patel²¹Dept. of Paediatrics, Shiv Multispeciality Hospital, Ahmedabad, Gujarat, India²Dept. of Forensic Medicine, Vedanta Institute of Medical Science, Palghar, Maharashtra, India

ARTICLE INFO

Article history:

Received 03-05-2022

Accepted 23-05-2022

Available online 30-06-2022

Keywords:

Haemoglobin

Iron deficiency

Thalassemia

Serum Fe

Serum B 12

ABSTRACT

Background: Anaemia is a major global health problem; it results in significant morbidity and mortality among children**Materials and Method:** 85 children below 12 years having anaemia were studied. Complete blood count was done to determine PCV, MCV, MCH, MCHC, RDW, Hb %, along with Peripheral smear (PS) study, Reticulocyte count and Serum iron (Fe), Serum B₁₂ and Folic acid levels were done**Results:** Types of anaemia were 50(58.8%) Iron deficiency anaemia, 12(14.1%) Thalassemia, 8 (9.41%) Megaloblastic anaemia, 7(8.23%) Anaemia of acute haemorrhage, 5 (5.88%) Sickle cell anaemia, 2(2.35%) Aplastic anaemia, 1(1.17%) Leukemia. The clinical manifestations were 85 (100%) pallor, 71(83.5%) weakness and fatigability, 32(37.6%) fever, 23(27.05%) 17(20%) shortness of breath, 15(17.6%) hepatomegaly, 12(14.1%) cough, 11(12.9%) history of pica, 11 (12.9%) splenomegaly, 9(10.5%) petechiae, 8(9.4%) Nausea/vomiting, 8(9.4%) koilonychia, 4(4.7%) hyper pigmentation, 5(5.8%) tremors. Out of the total number of patients screened 18(21.1%) had mild, 39(45.8%) moderate and 28(32.9%) had severe anaemia.**Conclusion:** The present study of anaemia had Iron deficiency as the major cause followed by Thalassemia, Megaloblastic anaemia, acute Haemorrhage, Sickle cell anaemia and Aplastic anaemia. This study will be helpful to paediatricians to treat such patients efficiently to avoid morbidity and mortality.This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.For reprints contact: reprint@ipinnovative.com

1. Introduction

Anaemia is a term used for all pathological conditions in which the blood haemoglobin concentration decreases to an abnormally low level, due to a deficiency in one or more micro nutrients. The main nutrients involved in the synthesis of haemoglobin are iron, folic acid, and vitamin B12. Anaemia is a serious health problem in India and under developed countries.¹ About two million people (30% of world population) are anaemic and it's mainly due to iron deficiency; in developing countries like India this figure is exacerbated by malaria worm infestations and

mal-nutrition.² As per the NFHS (National Family Health survey) report prevalence of anaemia among children less than 12 years of age is 60-70%.³

Anaemia usually develops as breast milk is replaced by food that are deficient in micro nutrients including iron, vitamin B₁₂ and folic acid, during early childhood, bad feeding habits, especially during weaning period, exacerbate the problem. Impaired cognitive function, growth and psychomotor development are the manifestation of anaemia in children.⁴ Hence an attempt was made to evaluate the clinical manifestations, grades of anaemia in different age groups so that the study will be a guide line for paediatricians.

* Corresponding author.

E-mail address: dr.nilesh27@yahoo.com (N. T. Shah).

2. Materials and Methods

85 (Eighty Five) children aged below 12 regularly visiting to paediatric OPD of Vedanta Institute of Medical Sciences, Palgarh, Maharashtra were studied.

2.1. Inclusive criteria

Children between the age group of 6 months to 12 years with pallor clinically diagnosed as anaemia were selected for study.

2.2. Exclusion criteria

Children more than 12 years, less than 6 months, children with congenital heart disease, immune compromised, tuberculosis and hepatitis were excluded from study.

2.3. Method

Routine blood examination for anaemia was morphologically based on peripheral smear (PS) findings packed cell volume (PCV), Mean corpuscular volume (MCV), Mean corpuscular haemoglobin (MCH), Mean corpuscular haemoglobin Concentration (MCHC) and red cell distribution width (RDW) were determined by automated cell counter Haemoglobin was estimated by sahil's method and expressed in gm%, PS (peripheral smear) was stained by Leishman's stain. Reticulocyte count was done by Brilliant crystal stain method; serum iron determination was done by Romany's bipyridyl method. Total iron binding capacity was determined by Ramsay's method, serum vitamin B12 and folic acid was determined by architect method.

The duration of study was June-2020 to July-2021.

2.4. Statistical analysis

Various findings including different types of anaemia, clinical manifestations, grades of anaemia were classified with percentage. The statistical analysis was done in the SPSS software. The ratio of male and female was 2:1.

2.5. Observation and Results

Table 1: Age wise distribution of paediatric patients – 13(15.2%) patients were between 6 months to 1 year, 35(41.1%) were 1 to 5 years, 27(51.7%) was between 6 to 9 years, 10(11.7%) were 10 to 12 years of age.

Table 2: Prevalence of different types of Anaemia 50(58.8%) Iron deficiency, 12(14.1%) thalassamia, 8(9.4%) megaloblastic anaemia, 7(8.23%) anaemia of acute haemorrhage, 5(5.88%) sickle cell anaemia, 2(2.55%) were aplastic anaemia, 1(1.4%) leukaemia.

Table 3: Clinical manifestations of Anaemic patients 85(100%) had pallor, 71(83.5%) weakness and fatigability, 32(37.6%) fever, 23(27.05%) Icterus, 17(20%) shortness

of breath, 15(17.6%) hepatomegaly, 12(14.1%) cough, 11(12.9%) History of pica, 11(12.9%) splenomegaly, 9(10.5%) petechiae, 8(9.4%) nausea/vomiting, 8(9.4%) koilonychia, 4(4.7%) hyper pigmentation, 5(5.8%) tremors

Table 4: Grades of Anaemia on the basis of haemoglobin level – Mild 13 male, 5 female, total 18(21.1%), Moderate 28 male, 11 female, total 39(45.8%) severe 18 male, 10 female, total 28(32.9%).

Table 1: Age wise distribution of patients of anaemia

Age in years	No. of patients	Percentage %
6 month to 1 year	13	15.2
1 to 5 years	35	41.1
6 to 9 years	27	31.7
10 to 12 years	10	11.7
Total	85	

Table 2: Prevalence of different types of anaemia

Diseases	No. of patients	Percentage %
Iron deficiency Anaemia	50	58.8
Thalassaemia	12	14.1
Megaloblastic anaemia	8	9.41
Anaemia of acute haemorrhage	7	8.23
Sickle cell anaemia	5	5.8
Aplastic anaemia	2	2.35
Leukaemia	1	1.17

Table 3: Clinical manifestations in anaemia patients

Clinical Manifestations	No. of Patients	Percentage %
Pallor	85	100
Weakness and fatigability	71	83.5
Fever	32	37.6
Icterus	23	27.05
Shortness of breath	17	20
Hepatomegaly	15	17.6
Cough	12	14.1
History of pica	11	12.9
Splenomegaly	11	12.9
Petechiae	9	10.5
Nausea/Vomiting	8	9.4
Koilonychia	8	9.4
Hyper pigmentation	4	4.7
Tremors	5	5.8

3. Discussion

In the present study of prevalence of anaemia and haematological parameters in children below 12 years – 13(15.2%) children were between 6 months to 1 year,

Table 4: Grades of anaemia on the basis of haemoglobin level

Grades of anaemia	Male	Female	Total	Percentage %
Mild	13	5	18	21.1
Moderate	28	11	39	45.8
Severe	18	10	28	32.9
Total	59	26	85	100

35(41.1%) were between 1 to 8 years, 27(31.7%) were aged between 6 to 9 years, and 10(11.7%) were between 10 to 12 years (Table-1). Types of anaemia were 50(58.8%) had iron deficiency 12(14.1%) had thalassemia, 8(9.4%) had megaloblastic anaemia, 7(8.23%) had anaemia of acute haemorrhage, 5 (5.88%) had sickle cell anaemia, 2(2.35%) had aplastic anaemia, 1(1.17%) had leukaemia (Table 2). The clinical manifestation included 85 (100%) pallor, 71(83.5%) weakness and fatigability, 32(37.6%) had fever, 23(27.05%) Icterus, 17(20%) had shortness of breath, 15(17.6%) hepatomegaly, 12(14.1%) had cough, 11(12.9%) had history of pica, 11(12.9%) had splenomegaly, 9(10.9%) had petechiae, 8(9.4%) had nausea/vomiting, 8(9.4%) koilonychia, 4(4.7%) hyper-pigmentation, 5(5.8%) had tremors (Table 3). 18(21.1%) had mild, 39(45.8%) moderate, 28(32.9%) had severe grade of anaemia (Table 4). These findings are more or less in agreement with previous studies.⁵⁻⁷

During the study of anaemia in paediatric patients included pertinent issues, related to the history, physical examination and initial laboratory investigations, Hematocrit, (HCT) is the fractional volume of whole blood sample occupied by RBC expressed as percentage. As an example the normal HCT in children from 6 months to 12 years is approximately 40% and Haemoglobin in children of 6 months to 12 years is approximately 13.5 g/dL (135 g/L).

Characterising the symptoms help to elucidate the severity and chronicity of anaemia and may identify patients with blood loss or haemolytic aetiologies. Common symptoms of anaemia include lethargy, tachycardia and pallor.⁸ Infants with anaemia may present with irritability and poor oral intake. Changes in urine colour, scleral icterus or jaundice may indicate the presence of haemolytic disorders, such as G6PD (glucose 6-phosphate dehydrogenase) deficiency. Bleeding from GIT (gastro-intestinal tract) includes changes in stool colour; identification of blood in stool, history of bleeding symptoms must be reviewed. Severe or chronic epistaxis also may result in anaemia from blood loss and iron deficiency.

Previous medical history also plays vital role to find out the cause of anaemia in children, duration of gestational age at birth, hospitalisation and history of jaundice and / or anaemia in the new born period. Travel to / from areas

of endemic infections (Eg. Malaria, hepatitis, tuberculosis) should also be ruled out to evaluate the cause of anaemia. Moreover herbal or oxidant drugs may cause haemolysis particularly in patients with G6PD deficiency, possible environmental toxins exposure should be explored including lead exposure and nitrates in well water, family history of inherited haemolytic anaemia.⁹

Anaemia with high absolute reticulocyte count (ARC) reflects an increased erythropoietic response for haemolysis or loss of blood. Anaemia with a low or normal ARC reflects deficient production of RBC (i.e. reduced bone marrow response to anaemia). However, haemolysis or blood loss can be associated with low concurrent disorder that impairs RBC production. In some cases reticulocyte count depends on the phase of illness.¹⁰

Review of PS (peripheral smear) is an essential part of anaemia evaluation. Even if patients RBC indices are normal review of the blood smear may reveal abnormal cells that can help to identify the cause of anaemia.

The diagnostic approach to anaemia includes pancytopenia in leukaemia, thrombocytopenia indicate haemolytic uremic syndrome and thrombocytosis in iron deficiency, leukocytosis or elevated WBC count includes leukaemia and infections.

4. Summary and Conclusion

The present study of anaemia in children below 12 years of age is mainly related to malnutrition. The prevalence of high lymphocyte count in anaemia indicates viral infection. The prevalence of anaemia was higher in the lower age group is due to frequent infections. Girls of preschool age had a probable iron, vitamin B12 or foliate deficiency as indicated by high RDW values. Girls of adolescent age (11-12 years) were more anaemic indicating more nutritional requirement with onset of puberty. Overall children below 12 years boys were found to be suffering from higher level of hypochromic and microcytic anaemia. The present study recommends about pure drinking water, sanitation and nutritional counselling to the parents having low social economic status.

5. Limitation of study

Owing to tertiary location of present institution, limited number of patients, lack of latest (advanced) instruments we have limited findings.

6. Conflict of Interest

The authors declare no relevant conflicts of interest.

7. Source of Funding

None.

References

1. Slotzfus RJ, Dreyfus ML. Nutrition foundation; A draft document prepared for the international Nutritional Anaemia consultative group (INCAG) 17th July. Geneva; 1997.
2. World Health Organization. (2008). The global burden of disease : 2004 update. World Health Organization. Available from: <https://apps.who.int/iris/handle/10665/43942>.
3. National family health survey for Indian conducted by Mumbai, India. International Institute for population science. NFHS-3; 2006. Available from: <https://dhsprogram.com/pubs/pdf/frind3/frind3-vol1andvol2.pdf>.
4. National family health survey India International Institute for population science. NFHS-2; 1999.
5. da Silva D, Franceschini SCC, Priore SE, Ribeiro SMR, Szarfarc SC, Souza SB, et al. SC - Anaemia ferro private meriandasnarede publica de saude do municipio de vicasa minas Gerais. *Rev Nutr*. 2002;15(3):301–8.
6. Kumar S, Prakash J. Prevalence of anaemia under five years old children. A cross sectional hospital based study. *Int J Contemp Med*. 2018;6(2):52–7.
7. Piel FB, Patil AP. Global epidemiology of Sickle haemoglobin in neonates. *Lancet*. 2013;38(1):142–51.
8. Panigrahi S, Patra PK. The screening and morbidity pattern of anaemia in chattisgarh. *Ind J Haotol Blood Transfus*. 2015;31(1):104–9.
9. Madoori S, Ramya C. Clinico- haematological profile and outcome of anaemia in children at tertiary care hospital Karim Nagar. *Int J Res Med Sci*. 2015;3(12):3567–71.
10. Meshram SA. A hospital based study of anaemia in children of Adilabad a tribal district of Andhra Pradesh. *Int J Biol Med Res*. 2013;4(1):2894–7.

Author biography

Shivendra Varshney, Paediatrician

Nilesh T Shah, HOD

Umang Patel, Associate Professor

Cite this article: Varshney S, Shah NT, Patel U. Study of prevalence of anaemia and haematological parameters in children below 12 years. *IP Int J Med Paediatr Oncol* 2022;8(2):65-68.