



Original Research Article

Phytochemical, Antioxidant and pharmacological study of *Heterotis Rotundifolia* (SM.) JACQ.- FEL. (Melastomataceae)

Sylvestre Dossou Eteko¹, Kohue Christelle Chantal N’Gaman-Kouassi^{1,*},
Treyia Rodrigue Yao², Janat Akhanovna Mamyrbekova-Bekro¹,
Yves-Alain Bekro¹

¹Dept. of Basic and Applied Sciences Training and Research, Nangui Abrogoua University, Abidjan, Côte d’Ivoire

²Dept. of Animal Physiology, Félix HOuphouet-Boigny University, Côte d’Ivoire



ARTICLE INFO

Article history:

Received 10-05-2022

Accepted 26-05-2022

Available online 06-07-2022

Keywords:

Heterotis rotundifolia

Phytochemical screening

DPPH test

Writhing test

Antiinflammatory activity

ABSTRACT

In developing countries, particularly those located south of the Sahara, research in both phytochemistry and phytopharmacology is a means of valorization endogenous medicinal plants into accessible phytomedicines for the populations. Our study focused on the species *Heterotis rotundifolia*, a Melastomataceae growing in Côte d’Ivoire, and used in traditional medical care for various diseases. Qualitative (color reaction and TLC identification tests) and quantitative (elemental analysis) phytochemical tests as well as antioxidant and pharmacological evaluations on the species were carried out. All the results obtained showed that the study plant contains, on the one hand, mineral elements and secondary metabolites that condition its antioxidant, analgesic, and anti-inflammatory properties.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The Ivorian flora is a reservoir of plants of pharmacological interest. From this point of view, it represents a natural pharmacy from which new molecules for therapeutic purposes can be discovered in the expectation of obtaining drugs with reduced side effects.¹ In recent years, research on plants of therapeutic interest has experienced a meteoric rise in view of the many scientific data collected, mentioning their all-round benefits.² Many authors^{2,3} believed that the therapeutic effect of these plants for the treatment of various diseases is governed by their phytochemical compositions. The present study was carried out on *Heterotis rotundifolia* (Sm.) Jacq. -Fél. (Melastomataceae) used in traditional Ivorian medicine to treat measles and gestational diabetes. Apart from Eteko et al. work⁴, there is no scientific data

on the phytochemical composition of the Ivorian species. We recall, however, that works on the leaves of the plant have been published. Indeed, Agwu et al.⁵ showed that *H. rotundifolia* leaves reduce the osmotic fragility of human hemoglobin S (HbS) erythrocytes. Ogunka-Nnoka et al.⁶ demonstrated by GC-MS analysis of the fat extracted from the same organs of this plant, that it contains 83.79% saturated fatty acid and 9.72% unsaturated fatty acid. The overall aim of this study was to highlight the phytochemical composition, the antioxidant, peripheral analgesic and anti-inflammatory activities of *H. rotundifolia* growing in Côte d’Ivoire.

2. Materials and Methods

2.1. Plant material

The whole plant of *H. rotundifolia* was the matrix of the study carried out. It was harvested in May 2018 on

* Corresponding author.

E-mail address: kohuechristelle@yahoo.fr

(Kohue Christelle Chantal N’Gaman-Kouassi).

Table 1: Developers used for CCM

Selective fraction	Developer
Hexane	Hexane / AcOEt (5: 1.5 (v/v))
CHCl ₃	Hexane / AcOEt / CHCl ₃ / AcOH (3.5 / 3 / 2.5 / 0.5 (v/ v/ v/ v))
AcOEt	Hexane / AcOEt / CHCl ₃ / AcOH (2 / 1.5 / 4 / 0.5 (v/v/v/v)) + 5 drops of NH ₄ OH
n-BuOH	AcOEt / CHCl ₃ / AcOH / EtOH (2 / 4.5 / 0.6 / 0.5 (v/v/v/v))
H ₂ O	AcOEt / CHCl ₃ / H ₂ O / MeOH (2 / 4.5 / 0.6 / 0.5 (v/v/v/v))

AcOH: acetic acid, MeOH: methanol, EtOH: ethanol

Table 2: Phytochemical families present in *H. rotundifolia*

Phytochemical family	Type of Test	Results
Polyphenols	FeCl ₃ (2%)	Positive
Flavonoids	Shinoda test	Positive
	Basic lead acetate	Positive
Coumarins	KOH (2%)	Positive
Tannins	Stiasny	Positive
Alkaloids	Dragendorff	Negative
Sterols, Polyterpenes	Acetic anhydride + H ₂ SO ₄	Negative
Reducing compounds	Fehling liquor	Positive
Saponins	Foam Index	130

Positive: Presence Negative: Absence

Table 3: Phytochemicals of *H. rotundifolia* identified by TLC

Selective fraction	Identified phytochemical [Rf] color
Hexane	Lupane triterpenes ^{c,d} : [0.05] Y ^{c2} -B ^{c1} ; [0.22 et 0.47] O ^{d2} Sterols ^d : [0.05] Y ^{d2} , [0.18] G ^{d1} ; [0.5] Y ^{d1} -G ^{d2} 1,8-dioxyanthracene ^b : [0.09; 0.15; 0.18; 0.22; 0.35; 0.40; 0.50; 0.60] R ^{b2} . [0.32; 0.45] Y ^{b1} -R ^{b2} . [0.65] G ^{b1} -R ^{b2}
CHCl ₃	Triterpenes ^d : [0.06; 0.21] P ^{d1} . [0.34] P ^{d1} -V ^{d2} . [0.45] B ^{d2} Coumarins ^{b,e} : [0.26; 0.29; 0.37; 0.45] B ^{b2} ; [0.27; 0.42; 0.81; 0.91] Y ^{b2} . [0.47] G ^{e2} Alkaloids ^g : [0.5; 0.77; 0.81; 0.87; 0.95] OY ^{g1}
AcOEt	Coumarins ^{e,b} : [0.15] B ^{e2} ; [0.22] B ^{c2} ; [0.47; 0.81] Y ^{b1} -B ^{c,b2} NI: [0.70] B ^{f1} Flavonoids: [0.15; 0.85] B ^{h2} . [0.47; 0.87] Y ^{h1} Alkaloids ^g : [0.52; 0.72; 0.75; 0.89] O ^{g1}
n-BuOH	Coumarins ^{b,e} : [0.06] B ^{b2} . [0.87] Y ^{b2} . [0.12; 0.91] G ^{e2} Flavonoids ^h : [0.06; 0.12] G ^{h2} . [0.91] G ^{h1} . [0.69] O ^{h2}
H ₂ O	Coumarins ^{b,e} : [0.11; 0.34; 0.40; 0.07; 0.65; 0.75] B ^{b2} -G ^{e2}

OY: orange-yellow; Y: yellow; G: green; R: red; B: blue; O: orange; P: purple; a: without developer; b: KOH; c: sulfuric vanillin; d: Liebermann Burchard; e: Basic lead acetate; g: Dragendorff; j: FeCl₃; h: AlCl₃; k: NH₃; _NI: Not identified; 1: visible 2: UV/365.



Table 4: Antioxidant parameters of *H. rotundifolia*

Sample	CR ₅₀ (mg/ml)	IE	EA
<i>H. rotundifolia</i>	0.5867	19.56	1.704 10 ⁻³
Vitamin C	0.095	3.17	7.886 10 ⁻²
Rutin	0.220	7.33	9.095 10 ⁻²

Table 5: Percentages of inhibition of abdominal cramps induced by the injection of acetic acid by aceclofenac and *H. rotundifolia*

Batch	Number	% inhibition
Witness	30 ± 1.69	-
Aceclofenac (75 mg/kg of CM)	14.7 ± 4.35**	55.11
(150 mg/kg of CM)	3.17 ± 2.23***	89.44
(300 mg/kg of CM)	7.17 ± 3.55***	76.11
	3.50 ± 1.75***	88.83

***P < 0.001: significant difference of the treated batches compared to that of the control batch; n = 6

**Fig. 1:** Quantitative mineral composition in macroelement (A) and trace elements (B) of *H. rotundifolia***Fig. 2:** Inhibition percentage of carrageenan-induced edema as a function of time by *H. rotundifolia* and indomethacin

the premises of University NANGUI ABROGOUA (UNA) located in Abidjan (5° 23' 21" north, 4° 01' 09" west). The species was authenticated at the National Floristic Center (CNF) in Abidjan in accordance with the available herbarium (*Malan 552 reference*). After harvesting and cleaning with water, the plant was dried during 5 days in an air-conditioned room (16°C). It was then reduced to powder for the tests.

2.2. Methods

2.2.1. Phytochemical tests

2.2.1.1. Determination of mineral composition. The elemental analysis of *H. rotundifolia* leaves, stems and roots was carried out at the Analysis and Research Center of the PETROCI laboratory (Abidjan-Côte d'Ivoire) using an X-ray fluorescence spectrometer (XRF).

Using a hydraulic press, 4 g of plant powder, previously incinerated, are pressed (pressure of 10 tons) to make pellets. These pellets were introduced into the spectrometer and bombarded by photons. The primary radiation is absorbed, the atoms of the sample ionized and bridged by internal electrons emitted (beam of secondary X-rays). Electronic relaxation releases energy (X-ray fluorescence) in the form of characteristic photons for each atomic element. These photons detected by a counter make it possible to identify the atom according to its energy and the flux of photons received by the analyzer makes it possible to deduce the mass concentration. The results of the quantification of mineral elements are expressed as a percentage (%).

2.2.1.2. Identification tests by color reaction and foam index (Im). The phytochemical screening was carried out according to the methodologies of reported works.⁷⁻⁹

2.2.1.3. TLC identification test. TLC identification test was carried out on the selective extracts in accordance with the methodologies from the literature.⁷⁻¹¹ The aqueous macerate of the vegetable powder (10 ml) was successively exhausted with 5 ml of hexane, chloroform (CHCl₃), ethyl acetate (AcOEt) and n-butanol (n-BuOH). The selective fractions and the residual aqueous extract were analyzed by TLC on Silufol chromatoplate (aluminum support, silica gel 60 F₂₅₄, thickness 0.2 mm; Merck). For the migration of the phytochemicals, various developers (solvent systems) were used (Table 1). The molecular fingerprints of the phytochemicals

were revealed using Liebermann-Bürchard reagent and sulfuric vanillin (5%, m/v) (terpenes and sterols), methanolic KOH solution (5%, m/v) and basic lead acetate (coumarins), NH₃ vapors (coumarins, anthocyanins, flavonoids), FeCl₃ solution (2%, m/v) (polyphenols, tannins), AlCl₃ solution (1%, m/v) (flavonoids) and Dragendorff reagent (alkaloids).

2.2.2. Antioxidant evaluation: DPPH test

The evaluation of the antioxidant activity by spectrophotometry was carried out.^{4,12,13} The reference antioxidants were the solutions of vitamin C and rutin prepared under the same conditions. To 1 ml of methanolic extract of the plant powder (0.0032 to 2 mg/ml) was added 2.5 ml to a methanolic solution of DPPH (0.03 mg/ml). The absorbance of the reaction mixture was read every 3 min for 30 min at 517 nm on a UV-visible spectrophotometer (Aqualytic A1800). The reduction percentage (PR); the efficiency index (IE) and the antiradical efficiency (EA) were determined from equations (1), (2) and (3):

$$PR (\%) = (1 - (\frac{\text{Absorbance extract}}{\text{Absorbance contrle}})) \times 100 \quad (1)$$

$$IE (\text{mg extract} / \text{mg DPPH}) = \frac{CR_{50}}{(DPPH)_{t=0}} \quad (2) \quad EA =$$

$$\frac{1}{CR_{50} \times TCR_{50}} \quad (3)$$

CR₅₀: Concentration that reduces 50% of the initial DPPH concentration

(DPPH) t = 0: Initial concentration of the DPPH solution

TCR₅₀: Time required reducing 50% of the initial concentration of DPPH

2.2.3. Pharmacological tests

For the pharmacological tests, we prepared samples with 20 g of vegetable powder macerated in 300 ml of distilled water under constant stirring for 24 h. After filtration, the macerate is concentrated at 80°C using a rotary evaporator (BÜCHI (Wather Bathb-480)). The concentrate obtained was maintained at 45°C in an oven for 48 hours.

2.2.3.1. Evaluation of peripheral analgesic activity: Writhing test. The methodology used is similar to that of Collier et al.¹⁴ and Kouakou-Siransy et al.¹⁵ The analgesic effect was achieved on five homogeneous batches each containing five Swiss albino mice (3 months), 22 g male and female, from the animal facility of the Laboratory of Animal Physiology of UNA:

1. Batch 1: control treated with distilled water before the intraperitoneal injection of acetic acid (1%);
2. Batch 2: reference treated with aceclofenac (100 mg/kg Body Mass (BM)), a non-steroidal antiinflammatory drug;
3. Batch 3: test with extract at 75 mg/kg of BM by gavage;
4. Batch 4: test treated with the extract at 150 mg/kg of BM by gavage;

5. Batch 5: test treated with the extract at 300 mg/kg of BM by gavage.

Intraperitoneal injection of acetic acid (1%) in mice causes a painful syndrome resulting in characteristic contortions (stretching movements) of the hind legs and the dorso-ventral muscle. After the administration of the samples, an intraperitoneal injection of acetic acid was made to the mice. For each dose, the percentage inhibition (% Inh) of cramps is determined according to expression:

$$\% \text{ Inh} = ((\text{Nb control} - \text{Nb treated}) / \text{Nb control}) \times 100 \quad (4)$$

2.2.3.2. Assessment of anti-inflammatory activity. The evaluation of the anti-inflammatory activity was carried out according to the method reported by Winter and Nuss.¹⁶ The intraperitoneal injection of carrageenan (1%) administered to the hind paw of the rat causes the appearance of an edema, the size of which was assessed by measuring the circumference of the paw. The rats used are young Wistar albino females (6 months) weighing 100 g from the animal facility of the Normal Superior School (ENS). Inflammation was reduced by the application of anti-inflammatories. Several treatments were administered to five batches of rats:

1. batch 1: Control treated with distilled water;
2. batch 2: Reference treated with indomethacin at 10 mg/kg BM (NSAID);
3. batch 3: Test treated with the extract at 75 mg/kg of BM by gavage;
4. batch 4: Test treated with the extract at 150 mg/kg of BM by gavage;
5. batch 5: Test treated with 300 mg/kg of BM by gavage from which an inflammation was induced 1 hour before.

The percentage inhibition of inflammation (% Inh) was calculated by comparing the mean of the increase in inflammation with that of the control group according to the expression (5):

$$\% \text{ Inh} = \frac{C-C_1}{C} \times 100 \quad (5)$$

C: average percentage increase in the circumference of the oedematous paw of the control batch.

C₁: average percentage increase in the circumference of the oedematous paw of the test group

2.3. Statistical analyzes

One-way analysis of variance (ANOVA) tests were used for analgesic and anti-inflammatory activities using Graph Pad Prism software version 5. Multiple comparisons of means by Tukey's test was made when a significant difference is noted. The significance of the test is determined by comparing the probability (P) associated with the test statistic to the theoretical value $\alpha = 0.05$. If $P \geq 0.05$, the

difference between the means is insignificant. If $P < 0.05$, the difference is significant.

3. Results and Discussions

3.1. Mineral composition

The results of the quantitative elemental analysis of the leaves, stems and roots of *H. rotundifolia* highlighted the variable contents of macroelements (Ca, K, Si, Mg, Na, S, P) and trace elements (Al, Fe, Ti, Mn) (Figure 1). Minerals have variable and diverse actions. They are essential to our body, in particular to the constitution of the skeleton, the teeth and the hair, and enter into many cellular reactions. Calcium (Ca) is the most abundant mineral element. Ca and phosphorus (P) are involved in building and maintaining tissues. The minerals sodium (Na), potassium (K), P, magnesium (Mg), iron (Fe) and manganese (Mn) are useful in regulating metabolism.¹⁷ K is present in the intracellular medium. In addition, it is also essential for maintaining muscle and nerve cell function.¹⁸ Among other main trace elements for plants, Fe is present in the analysed plant organs. This trace element plays a major role in the manufacture and functioning of haemoglobin and myoglobin.¹⁹ Ca and Mg contribute to good muscle function and are involved in cell division and differentiation processes.²⁰ Aluminum (Al) is a non-essential trace element, but classified as a nutrient. It seems to be the most prevalent in the environment, but plays no physiological role in metabolic processes.²¹ However, the accumulation of Al is a principal primitive characteristic of certain tropical families including the Melastomataceae.²²

3.2. Composition of secondary metabolites

The secondary metabolite composition of *H. rotundifolia* was determined (Table 2). It reveals the presence of seven phytochemical families: polyphenols, flavonoids, coumarins, tannins, reducing compounds and saponins. Shinoda test (Mg in concentrated HCl) demonstrated the presence of flavanones and dihydroflavonols among the flavonoids present.²³ Furthermore, another source from the bibliography²⁴ indicates that Shinoda test also shows the presence of flavones. Indeed, their presence in a sample was confirmed by their reduction to anthocyanidins. The conjugated system in the flavonoids produces a yellow color, while the conjugation in the anthocyanidin gives a red coloration. The color variation is the indicator of the presence of flavones. The moss index ($Im = 130$) confirms the detection of saponins. Polyphenols, flavonoids and coumarins have various biological properties²⁵, and are known for their inhibitory action on enzymes responsible for the production of free radicals.²⁶ Tannins also have the ability to inhibit the growth of many microorganisms,²⁷ and saponins have antimicrobial, antibacterial and analgesic activities.²⁸ Thus, there is every reason to believe

that *H. rotundifolia* use as a traditional medicinal treatment is governed by the co-presence of the bioactive phytoconstituents contained in the plant.

Phytochemical screening by TLC of the selective fractions was carried out (Table 3), on the one hand, in order to have a more extensive overview of the phytochemical composition of *H. rotundifolia*, and on the other hand, to isolate the suspected constituents responsible for its therapeutic properties. The results obtained confirm those obtained by colored reactions, unlike the absence of alkaloids and sterols and polyterpenes (Table 2).

Thanks to TLC screening, the presence of lupane, oleanane and ursanne type triterpenes,^{23,29} anthracene derivatives (anthracene; 1,8-dioxanthracene),³⁰ phenolic acids,²⁹ coumarin derivatives (angelicin or coumestrol; 7, 8-dihydroxycoumarin; 7,8-dihydroxy-6-methoxycoumarin)³¹ and alkaloids³² was confirmed with regard to the specificity of the molecular fingerprint differences. *H. rotundifolia* species from Nigeria does not contain alkaloids.³³ Polyphenols and flavonoids were demonstrated by quantification in the species from Benin.³⁴

3.3. Antioxidant activity

H. rotundifolia antioxidant potential against the DPPH radical was evaluated by comparison with vitamin C and rutin used as reference antioxidants. The results were shown in Table 4. The reduction percentages of DPPH (Table 4) made it possible to graphically determine the median concentrations (CR_{50}), reducing 50% of the DPPH. This parameter reflects the efficiency of the sample.³⁵ The lower its value, the more significant the antioxidant activity. The efficacy indices (IE) and the antiradical efficacy (EA) were also determined. The species *H. rotundifolia* presents an antioxidant potential with respect to the DPPH radical with a CR_{50} and an IE of values 0.5867 mg/ml and 19.56 respectively. Considering the method for EA determining and according to Sanchez-Moreno et al.³⁶ classification, *H. rotundifolia* has a slower DPPH-reducing action ($1.704 \cdot 10^{-3}$) than vitamin C ($7.886 \cdot 10^{-2}$) and rutin ($9.095 \cdot 10^{-2}$). This seems to depend on the ability to quickly give up hydrogen.^{37,38} We recall that the DPPH test makes it possible to determine the reducing capacity of an extract by hydrogen transfer.^{37,39} The overt antioxidant activity of *H. rotundifolia* appears to be confirmed by phytochemical screening, which reveals the presence of antioxidant phytochemical families. Published works^{34,40} have shown that *H. rotundifolia* species is antioxidant.

3.4. Analgesic activity

The analgesic activity of the plant extract at different concentrations (75; 150 and 300 mg/kg) was evaluated. The results obtained are recorded in Table 5. The pain inhibition by extract was compared to that induced by

aceclofenac widely used and/or associated in the treatment of a number of serious cases of clinical hepatotoxicity.⁴¹ The percentage inhibition of abdominal cramps induced by acetic acid injection of all doses received by mice (from 76.11% to 89.44%) is higher and more significant than that of aceclofenac (55.11%). The analgesic activity of *H. rotundifolia* could be explained by the presence of bioactive phytochemicals possessing analgesic properties among which flavonoids⁴² and coumarins.⁴³ It should also be noted that studies carried out on the chemical composition of the essential oils of the leaves of *H. rotundifolia* revealed that the leaves contain bioactive compounds. Studies⁶ showed that neophytadiene, a major constituent (35.8%) of the essential oil extracted from the leaves of *H. rotundifolia*, has antipyretic, anti-inflammatory, analgesic, antimicrobial properties and antioxidant.

3.5. Anti-inflammatory activity

The results of the evaluation of the anti-inflammatory activity of the aqueous extract of *H. rotundifolia* are recorded in Figure 2. After administration of carrageenan (1%) by intraperitoneal injection to the right hind paw of the rat, edema and inflammation was observed which increase with time. Indeed, the injection of carrageenan causes an inflammatory process accompanied by an edema evolving in 2 phases: phase 1 (1h) synthesis and release of chemical mediators (histamine and serotonin which maintain inflammation) and phase 2 (3h) formation of prostaglandins and leukotrienes due to activation of cyclooxygenase and lipoxygenase.⁴⁴ It is during the second phase that the inflammatory phenomena can be antagonized by natural anti-inflammatories, steroidal or not.^{45,46}

After injection of indomethacin, a gradual inhibition of edema was observed from 58.89% (2h) to 31.70% (5h). The plant extract also induces a significant inhibition at the concentrations 75 mg/kg (85.13% at 5 hours) and 150 mg/kg (88.62% at 3 hours). However, this inhibition varies slightly at 300 mg/kg (from 18.36% to 38.46%). This anti-inflammatory activity is comparable to that of indomethacin. At 150 mg/kg of BM, there is a significant difference (at the threshold $P < 0.01$) at 3h ($88.62 \pm 0.67^{**}$), 4h ($83.76 \pm 0.63^{**}$) and 5h ($81.15 \pm 1.63^{**}$). This inhibition is greater than that of indomethacin at all times. In view of the results obtained, it is well founded to affirm that *H. rotundifolia* is more active at doses of 75 and 150 mg/kg of BM. This seems to justify the use of the species in traditional medicine for the treatment of hemorrhoids⁴⁷ and measles.⁴⁸ There is no doubt that the anti-inflammatory activity of *H. rotundifolia* depends on its phytochemical composition. Indeed, phenols, saponins, tannins, triterpenes and flavonoids also possess anti-inflammatory properties.^{42,49,50}

4. Conclusion

The presence of mineral elements (macro and trace elements) essential for humans, as well as families of bioactive phytochemicals with antioxidant, analgesic and anti-inflammatory properties, was highlighted in *Heterotis rotundifolia*. The results obtained can justify the use of the plant in medicinal practices in Côte d'Ivoire.

5. Source of Funding

None.

6. Conflict of Interest

None.

Acknowledgments


We would like to thank the Department of Animal Physiology for allowing us to carry out this work.


References

- Grubben GJH, Denton OA. Plant Resources of Tropical Africa 11(1) : Medicinal plants 1; 2013. Available from: https://www.researchgate.net/publication/317510018_Plant_Resources_of_Tropical_Africa_111_Medicinal_plants_1.
- Arya OP. Ethnobotany and nutritional importance of four selected medicinal plants from Eastern Himalaya, Arunachal Pradesh. *J Med Plants Stud.* 2017;5(1):45–9.
- Devi KN. Estimation of essential and trace elements in some medicinal plants by PIXE and PIGE techniques. *Nucl Instrum Methods Phys Res.* 2008;266(8):1605–15. doi:10.1016/j.nimb.2007.12.004.
- Eteko DS. Antioxidant profiles of alcoholic tinctures from *Heterotis rotundifolia* (Sm.) Jacq. -Fel. (Melastomaceae) by DPPH• radical trapping. *European J Biomed Pharm Sci.* 2018;5(10):39–45.
- Agwu JA. Effect of *Heterotis rotundifolia* leaf extract on Hemoglobin-S (HbS) Polymerization, Osmotic fragility and Fe2+ /Fe3+ ratio. *Int J Sci Eng Res.* 2018;9(5):1872–7.
- Nnoka CUO. Nutrient and essential oil compositions of *Heterotis rotundifolia* leaves. *Amer J Biosci.* 2020;8(2):28–36.
- Ladiguina EY. Chemical analysis of medicinal plants. Moskva edition Vishaya Chkola. vol. 347; 1983.
- Dohou N. Phytochemical screening of an Ibero-Moroccan endemic *Thymelaea lithoides*. *Bull Soc Pharm.* 2003;4(2):61–78. doi:10.1016/S2222-1808(14)60324-8.
- Békro YA. Ethnobotanical study and phytochemical screening of *Caesalpinia benthamiana* (Baill) Herend. and Zarucchi (Caesalpinaceae). *Science et. Nature.* 2007;4(2):217–42. doi:10.4314/scinat.v4i2.42146.
- Brou KG. Sur la composition phytochimique qualitative des extraits bruts hydrométhanoliques des feuilles de 6 cultivars de *Manihot esculenta* Crantz de Côte d'Ivoire. *Eur J Sci Res.* 2010;45(2):200–11.
- N'gaman K. Étude phytochimique et effet d'extraits de *Gmelina arborea* Roxb. (Verbenaceae) de Côte d'Ivoire sur la stabilité osmotique d'érythrocytes. *J Appl Biosci.* 2013;144:14747–54. doi:10.35759/JABS.144.3.
- Espin JC. Characterization of the total free radical scavenger capacity of vegetable oils and oil fractions using 2,2-diphenyl-1-picrylhydrazyl radical. *J Agric Food Chem.* 2000;48(3):648–56.
- Sladjana MS. Kinetic behavior of DPPH radical scavenging activity of tomato waste extracts. *J Serb Chem Soc.* 2012;77(10):1381–9.
- Collier J. The abdominal contraction response and its suppression by antinociceptive drugs in the mouse. *Brit J Pharm Chem.* 1968;32(2):295–310. doi:10.1111/j.1476-5381.1968.tb00973.x.

15. Siransy GK. Study of the analgesic activity of the methanolic extract of *Gossypium* leaves *hirsutum* Linn. (Malvaceae). *J Pharm Biol Sci*. 2010;11(1):6–12.
16. Winter CA. Carrageenin induced oedema in hand paw of the rat as assays anti-inflammatory drugs. *Proc Soc Exp Biol Med*. 1962;111(62):544–7.
17. Kouassi AD. Phytochemical investigation of *Cissus aralioides* stems from Ivory Coast. *Amer J PharmTech Res*. 2021;11(3):1–14.
18. Stone SM, Martyn L, Weaver CM. Potassium intake, bioavailability, hypertension, and glucose control. *Nutrients*. 2016;8(7):1–13. doi:10.3390/nu8070444.
19. Camaschella C. Iron metabolism and iron disorders revisited in the hepcidin era. *Hematologica*. 2020;105(2):260–72. doi:10.3324/haematol.2019.232124.
20. Kouassi AD. Mineral and phenolic compositions, antioxidant activity and GC-MS analysis of the leaves of *Anchomanes difformis* (Blume) Engl. from Ivory Coast. *GSC Adv Res Rev*. 2022;10(1):145–55.
21. Igbokwe IO. Aluminum toxicosis: a review of toxic actions and effects. *Interdis Toxicol*. 2019;12(2):45–70.
22. Jansen S. Aluminum accumulation in leaves of 127 species in Melastomataceae with comments on the order Myrtales. *Ann Botany*. 2002;90(1):53–64. doi:10.1093/aob/mcf142.
23. Bruneton J. Pharmacognosie Phytochimie Plantes médicinales; 2016. Available from: https://www.amazon.in/Pharmacognosie-phytochimie-plantes-medicinales-ed/dp/2743011882#detailBullets_feature_div.
24. Chimie de base du test de Shinoda pour les flavonoïdes ?; 2021. Available from: <https://wikimho.com/fr/q/chemistry/13891>.
25. Gulcin I. Radical scavenging and antioxidant activity of tannic acid. *Arab J Chem*. 2010;3(1):43–53.
26. Dicarlo G. Flavonoids: old and new aspects of a class of natural therapeutic drugs. *Rev Life Sci*. 1999;65(4):337–53. doi:10.1016/S0024-3205(99)00120-4.
27. Sepúlveda L. Ellagic acid: Biological properties and biotechnological development for production processes. *Afric J Biotech*. 2011;10(22):4518–41.
28. Hirata T. Identification and physiological evaluation of the components from Citrus fruits as potential drugs for anti-corpulence and anticancer. *Bioorg Med Chem*. 2009;17(1):25–8. doi:10.1016/j.bmc.2008.11.039.
29. Lagnika L. Étude phytochimique et activité biologique de substances naturelles isolées de plantes Béninoises; 2005. p. 267.
30. Georgievsky BP. The bioactive compounds of medicinal plants; 1990. Available from: <https://www.theses.fr/2005STR13169>.
31. Glowniak K, Widelski J, Coumarins. Analysis by TLC. *Cazes J Encyclopedia of Chromatography*. 2010;533:511–3. doi:10.1016/B978-0-12-420067-8.00024-6.
32. Wagner H, Bladt S. Plant drug analysis in thin layer chromatography; 1996. p. 303–24.
33. Kutama RM. Phytochemical compositions in some Nigerian medicinal plants and their pharmacological properties. *J Anesthes*. 2018;6(1):15–25.
34. Dougnon VT. Chemical composition, cytotoxicity and antibacterial activity of selected extracts of *Euphorbia hirta*, *Citrus aurantifolia* and *Heterotis rotundifolia* on enteropathogenic Bacteria. *EC Microbiol*. 2017;12(4):180–95. doi:10.1016/j.jep.2020.113662.
35. Falleh H. Antioxidant activity and polyphenol content in the different organs of the *Cynara* wild artichoke *cardunculus*. *Arid Regions Review*. 2006;61:341–4. doi:10.1016/j.indcrop.2014.07.042.
36. Moreno CS. A procedure to measure the antiradical efficiency of polyphenols. *J Sci Food Agric*. 1998;76(2):270–6. doi:10.1002/(SICI)1097-0010(199802)76:2<270::AID-JSFA945>3.0.CO;2-9.
37. Cendrowski A. L-ascorbic acid, and antioxidant activity in wines from rose fruits (*Rosa rugosa*). *Molecules*. 2021;26(9):1–17. doi:10.3390/molecules26092561.
38. Yin X. Chemical stability of ascorbic acid integrated into commercial products: A review on bioactivity and delivery technology. *Antioxidants*. 2022;11(153):1–20.
39. Huang D. The chemistry behind antioxidant capacity assays. *J Agric Food Chem*. 2005;53(6):1841–56. doi:10.1021/jf030723c.
40. Abere TA. Antidiarrheal and toxicological evaluation leaf extract of *Dissotis rotundifolia* Triana (Melastomataceae). *BMC Comp Altern Med*. 2010;10(71):1–7.
41. Gupta A, Pandey AK. Aceclofenac-induced hepatotoxicity: An ameliorative effect of *Terminalia bellirica* fruit and ellagic acid. *World J Hepatol*. 2020;12(11):949–64.
42. Sahu R, Saxena J. Screening of total phenolic and flavonoid content in conventional and non-conventional species of curcuma. *J Pharm Phytochem*. 2013;2(1):176–9.
43. Reddy NS. Novel coumarin-3-(N-aryl) carboxamides arrest breast cancer cell growth by inhibiting ErbB-2 and ERK1. *Bioorg Med Chem*. 2005;13(9):3141–88. doi:10.1016/j.bmc.2005.02.051.
44. Rosa MD. Biological properties of carrageenan. *J Pharm Pharmacol*. 1972;24(2):89–102. doi:10.1111/j.2042-7158.1972.tb08940.x.
45. Alcaraz MJ, Jiménez MJ. Flavonoid and anti-inflammatory agent. *Fitoterapia*. 1988;59(1):25–38.
46. Adepo AA. Evaluation of the anti-inflammatory and antioxidant activities of the hydro-ethanolic extract of the root bark of *Dichrostachys cinerea* L. Wight and Arn. (Fabaceae); 2018.
47. Issali AE. Floristic study of the protected area from yacolidabou, located at soubré. *Octa J Env Res*. 2015;3(2):196–203.
48. Ajibesin KK. Ethnobotanical survey of plants used for skin diseases and related ailments in Akwa Ibom State. *Niger Ethno Res Appl*. 2012;10:463–522.
49. Nunes C. Plants as Sources of Anti-Inflammatory Agents. *Molecules*. 2020;2020(3726):1–22.
50. Yahfoufi N. The immunomodulatory and anti-inflammatory role of polyphenols. *Nutrients*. 2018;10(11):1–23. doi:10.3390/nu10111618.

Author biography

Sylvestre Dossou Eteko, Ph D Student  <https://orcid.org/0000-0001-5598-2011>

Kohue Christelle Chantal N’Gaman-Kouassi, Assistant Master  <https://orcid.org/0000-0002-2552-4471>

Treya Rodrigue Yao, -  <https://orcid.org/0000-0002-5165-3680>

Janat Akhanovna Mamyrbekova-Bekro, Professor  <https://orcid.org/0000-0003-1598-6020>

Yves-Alain Bekro, Professor  <https://orcid.org/0000-0002-6267-3791>

Cite this article: Sylvestre Dossou Eteko, Kohue Christelle Chantal N’Gaman-Kouassi, Treya Rodrigue Yao, Janat Akhanovna Mamyrbekova-Bekro, Yves-Alain Bekro. Phytochemical, Antioxidant and pharmacological study of *Heterotis Rotundifolia* (SM.) JACQ.- FEL. (Melastomataceae). *J Pharm Biol Sci* 2022;10(1):37-43.