

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Obstetrics and Gynecology Research

Journal homepage: www.ijogr.org

Case Report

Accessory fallopian tube –A rare anomaly

Manish R Pandya^{1,2,*}, Ghanshyam K Gorvadiya¹, Jeel A Modesara¹

¹Dept. of Obstetrics & Gynaecology, Scientific Research Institute, Surendranagar, Gujarat, India

²Dept. of Obstetrics and Gynecology, Nootan Medical Collage and Research Center, Visnagar, Gujarat, India



ARTICLE INFO

Article history:

Received 01-02-2023

Accepted 13-02-2023

Available online 18-02-2023

Keywords:

Accessory fallopian tube

Mullerian duct

Infertility

Ectopic pregnancy

Pyosalpinx

ABSTRACT

Accessory fallopian tube is a rare, congenital and developmental mullerian duct anomaly.¹ The documented incidence is around 6-10% in women seeking for infertility treatment.² We observed accessory fallopian tube in a patient during routine checking of operative field, ovaries and fallopian tube during caesarean section. Accessory fallopian tube is congenital anomaly which is attached with ampullary part of main fallopian tube. Accessory fallopian tube is common site for pyosalpinx, hydrosalpinx, cystic swelling and torsion which can lead to infertility and other complications. The ovum released by ovary may also captured by accessory fallopian tube which can lead to infertility or ectopic pregnancy. During the embryological development of female urogenital system, development of fallopian tube occurs from cranial unfused portion of mullerian ducts. This duct is derived from mesoderm, the middle layer of germ cell layers in the embryo. Bifurcation of cranial end of mullerian duct results in accessory fallopian tube.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

The paired paramesonephric (müllerian) ducts derive from the coelomic epithelium (mesoderm) of the posterior abdominal wall during the 5th–6th weeks of gestation. The cranial portions of the ducts become precursors of the FT, and the caudal portions grow caudally and medially to reach the pelvis and urogenital sinus, where they fuse to form the uterus (Figure 1). The cranial end of the FT is fimbriated and open to the coelomic cavity, allowing communication between the peritoneum and FT. The caudal end maintains communication with the uterine cornua. The uterine tubes (or fallopian tubes) are muscular ‘J-shaped’ tubes, found in the female reproductive tract. They lie in the upper border of the broad ligament, extending laterally from the uterus, opening into the abdominal cavity, near the ovaries. Fallopian tube is 10-12 cm long in size. The main function

of fallopian tube is to transport ovum from ovary to uterus and fertilization. The accessory fallopian tube sometimes pick up the ovum, which can lead to ectopic pregnancy. Accessory fallopian tube also plays an important role in infertility. Hence, all patients of infertility or PID should be screened for accessory fallopian tube. And if found, it should be removed.

2. Case Report

We found duplication of fallopian tube on right side in a 28-year old female patient. She underwent an emergency caesarean section (Figure 2). We found it during routine checking of operative field, ovaries and fallopian tube. We found a small tubular structure arising from the ampullary part of main fallopian tube, around 3-4 cm (Figure 3). The lateral end showed presence of fimbria similar to main fallopian tube, and the other end is obliterated. It was unilateral and contralateral tube was normal (Figure 4). We

* Corresponding author.

E-mail address: drmanish.pandya@gmail.com (M. R. Pandya).




Fig. 1: Embryologic morphology of fallopian tube



Fig. 4: Unilateral accessory tube

performed surgical excision of accessory fallopian tube and sent it to laboratory for histopathological evaluation. In this case, both ovaries and uterus were normal. No any other anomalies found. Histopathological examination confirmed accessory fallopian tube with presence of lumen (Figure 5).



Fig. 2: Accessory fallopian tube at CS



Fig. 3: Accessory tube arising from ampullary part

3. Discussion

The incidence of accessory fallopian tube was first described by Krossman et al. in 1894.³ Its incidence is about 5%. Based on the distance from the fimbria, they are termed terminal (<1 cm) and ampullary (>1 cm). The distance does not make any difference in the presentations usually⁴




Fig. 5: Histopathology report

Appearance may be stem-like, single or multiple, mostly extending from the ampulla of main fallopian tube and about 1-3 cm in length, may be unilateral or bilateral with umbrella shaped end. It is generally not connected with lumen of normal fallopian tube and has blind end attachment. Beyth and Kopolovic have postulated that the coelomic epithelium, which invaginates into the müllerian duct to form the ostia, is surrounded by one or many secondary invaginations. If they do not reach till the lumen, they persist as accessory tubes.⁵ Most of the patients with this anomaly are usually asymptomatic and are diagnosed incidentally on laparoscopy or laparotomy for some unrelated purpose. In our case, it was diagnosed during routine operative field checking of tubes and ovaries at caesarean section.

These accessory fallopian tubes contribute to infertility and ectopic pregnancy as fimbria are capable of capturing ova even if they are not in their normal anatomical position and occlusion of segment of the fallopian tube also does

not interfere with fimbrial ovum pickup function. This may cause decreased chance of intrauterine pregnancy in case of accessory fallopian tube. Hence, all patients of infertility should be screened for accessory fallopian tube and if found should be removed to prevent emergency and life threatening conditions. Transperitoneal migration of sperm through normal fallopian tube may migrate to fimbria of accessory tube and fertilize ovum captured by it, causing ectopic pregnancy. Medico-legal issue is another aspect as it may cause potential failure of postpartum sterilization procedure.⁶ Previous many studies suggest about such complications associated with accessory fallopian tube like, pyosalpinx, hydrosalpinx, cystic swelling, infertility, torsion and ectopic pregnancy. Torsion of accessory Right fallopian tube is more common than left, because the left tube is fixed in the left hemipelvis by sigmoid colon and mesentery.⁷ Ectopic pregnancy in accessory tube can lead to massive haemorrhage, and immediate surgical intervention is must.⁸

This anomaly may be associated with other mullerian duct anomalies and renal malformations like agenesis, ectopia, hypoplasia, fusion, malrotation, and duplication.⁹ In our case no such abnormalities were found. Early identification by pelvic ultrasound, hysterosalpingography or laparoscopy and it's preventive removal microsurgically followed by histopathological confirmation should be considered to avoid any dangerous consequences.^{6,10} Visualization of the tubes by laparoscopy or laparotomy is the only confirmatory modality.

Surgical management is with resection of the accessory stump after putting purse string sutures at the base or using energy sources by electrocoagulation of tubal stump. However, this should be performed after a thorough discussion about the pros and cons of the issue with the parents. Because this excision may itself predispose to adhesions or tubal blockade. Hence, maximum care should be taken while handling the tube.

4. Conclusion

Because of the rarity of this congenital variation, low suspicion index for its existence, and diagnostic subtleties associated with the condition, the diagnosis of accessory tube is often missed and cases under-reported. Nevertheless, this entity should always be kept in mind especially while dealing with the infertility cases. A high index of suspicion and meticulous systematic examination of the

fallopian tubes during abdominal surgeries is important for its early recognition and to prevent many gynecological complications. Preventive removal of such findings should be considered but weighed against its possible surgical complications like adhesions.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Rezvani M, Shaaban AM. FT disease in the Non pregnant Patient. *Radiographics*. 2011;31:527-48.
2. Brosens I, Gordon A. Tubal Infertility. In: The abnormal fallopian tube. Philadelphia: JB Lippincott Company; 1990. p. 30-2.
3. Beyth Y, Kopolovic J. Accessory tubes: A possible contributing factor in infertility. *Fertil Steril*. 1982;38(3):382-3.
4. Zheng X, Han H, Guan J. Clinical features of fallopian tube accessory ostium and outcomes after laparoscopic treatment. *Int J Gynaecol Obstet*. 2015;129(3):260-3.
5. Yablonski M, Sarge T, Wild RA. Subtle variations in tubal anatomy in infertile women. *Fertil Steril*. 1990;54(3):455-8.
6. Narayanan R, Rajeev M. Duplication of fallopian tube. *J Hum Reprod Sci*. 2008;1(1):35-6.
7. Coddington CC, Chandler PE, Smit GW. Accessory fallopian tube a case report. *J Reprod Med*. 1990;35(4):420-1.
8. Macnaughton-Jones H. Pyosalpinx in an Accessory fallopian tube. *Proc R Soc Med*. 1914;7:1-4.
9. Behr SC, Courtier JL, Qayyum A. Imaging of Mullerian duct anomalies. *Radiographics*. 2012;32(6):233-50.
10. Rottenstreich M, Smorgick N, Pansky M, Vaknin Z. Isolated torsion of accessory fallopian tube in a young adolescent. *J Pediatr Adolesc Gynecol*. 2016;29(4):57-8.

Author biography

Manish R Pandya, Professor & HOD  <https://orcid.org/0000-0003-4665-3069>

Ghanshyam K Gorvadiya, Senior Resident

Jeel A Modesara, Senior Resident

Cite this article: Pandya MR, Gorvadiya GK, Modesara JA. Accessory fallopian tube –A rare anomaly. *Indian J Obstet Gynecol Res* 2023;10(1):88-90.