

Content available at: <https://www.ipinnovative.com/open-access-journals>

IP Journal of Otorhinolaryngology and Allied Science

Journal homepage: <https://www.joas.co.in/>

Case Report

Unusual presentation of lung carcinoma as nasal tip metastasis: A review of literature

Jagat Singh¹, Usha Poonia^{2,*}, Manisha Kumari³, Sukriti Bansal³, Aman¹

¹Dept. of ENT, Pandit Bhagwat Dayal Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana, India

²Dept. of Obstetrics and Gynaecology, Pandit Bhagwat Dayal Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana, India

³Dept. of Anesthesia, Pandit Bhagwat Dayal Sharma Post Graduate Institute of Medical Sciences, Rohtak, Haryana, India



ARTICLE INFO

Article history:

Received 13-08-2021

Accepted 16-09-2021

Available online 01-11-2021

Keywords:

Hemangioma

Rhinophyma

Chemoradiation

ABSTRACT

Lung carcinoma is the most common type of carcinoma seen in males and 4th most common in females.¹ Skin metastasis from lung carcinoma is frequent with an incidence of 1%-12%, with most common site being the anterior chest wall.² Skin metastasis from lung carcinoma to the tip of nose is very rare with only 12 cases being reported in literature (Table 1). It can be confused with other benign and malignant conditions, such as infection, lymphoma, hemangioma, rhinophyma, sarcoidosis, tuberculosis and carcinomas, making its diagnosis difficult.³ Less than 20 cases of cutaneous nasal tip metastasis are reported in literature out of which 12 are from malignancies of lung. Sometimes it can appear earlier than the primary lesion and thus delays the diagnosis of primary lesion.⁴ In this case report, we report a case of nasal tip cutaneous metastasis from squamous cell carcinoma of lung.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Case Report

A 70-year-old male visited ENT opd with complaint of a lesion over nose since 3 months (Fig. 1), which was gradually progressive. Patient did not have any other symptom. It was about 5*5 cm in dimensions, ulcerated, erythematous and vascular on inspection. It was firm and tender on touch. No cervical lymph nodes were palpated. On magnetic resonance imaging of face, a well defined lobulated lesion of size about 3.7*2.5*3.6 cm was seen arising from the tip of nose with external surface bulge extending into the left nasal cavity (Fig. 2a, 2b). Punch biopsy was planned. All blood investigations were normal. Chest x-ray was done for pre anaesthetic check up and it showed homogenous opacity in the left upper lobe. (Fig.3) Punch biopsy was taken from nose which

on histopathological examination showed squamous cell carcinoma. Contrast enhanced computed tomography of chest showed large soft tissue mass in left upper lobe, abutting chest wall and showing contrast enhancement with left sided pleural effusion (Fig.4a,4b). Ultrasound guided fine needle aspiration cytology of left lung mass was done which showed squamous cell carcinoma of lung with inflammatory changes. He was referred to radiotherapy department where he was given palliative chemoradiation. Patient died after 2 months of cutaneous diagnosis.

2. Discussion

Skin metastasis are common in lung carcinoma with anterior chest wall being the most common site followed by the abdomen, extremities, neck, back and; head and neck.⁵ Metastasis to the tip of nose is however a very rare condition with only about 24 cases reported in

* Corresponding author.

E-mail address: amanjakhar03022000@gmail.com (U. Poonia).

Table 1:

Authors	Age/sex	Country	Metastasis	Primary	Treatment	Survival time from diagnosis of cutaneous metastasis
Ledderose GJ,Englhard. 2015	59yr/F	Germany	Small nodule on upper border of right nostril with extension towards tip and columella	Squamous cell carcinoma of dorsal thoracic esophagus	Palliative chemoradiotherapy for 1 yr with resection of painful soft tissue metastasis & palliative chemoradiation	Survived 18 months
Dong A, et. al, 2015	74yr/ F	china	Nodule on nasal tip	Squamous cell carcinoma esophagus	*	*
Kataoka A,et. Al. 1997	69yr/F	Japan	Nasal tip	Squamous cell carcinoma of uterine cervix	Radical hysterectomy followed by radiation	14 months
Itin PH, Heitzmann F, Stamm B. 1999	48yr/F	Switzerland	Nasal tip Radiation	Adenocarcinoma cervix	Preoperative radiotherapy followed by surgery with adnexectomy on right side with removal of urinary vesicle	Followed up for 5 months, patient was well till then
Antonio AM. 2015	58yr/F	Portugal	Dorsum of nose	Bilateral ovarian cystadenocarcinoma with malignant ascites nd hepatic nodule	Palliative chemotherapy	3 months
Gainers M, et. al. 2007	51yr/ M	United states of america	Erythematous indurated plaque nd papules over dorsum of nose (involving tip also)	Chronic lymphocytic leukemia	Chemotherapy & local radiation	Died
Pengfei C et. al. 2019	86yr/ M	china	Nose tip	Renal cell carcinoma(clear cell type)	Chemotherapy	Stable before submission of paper
Rajasekharan P.2004	55yr/ M	India Mumbai	Nasal mass	Renal cell carcinoma	*	*
Kohli SP.	54yr/ M	India Punjab	Nasal tip mass	Anaplastic thyroid malignancy	Palliative chemoradiotherapy	2 months
Isa NM, et. Al.	81/M	Malaysia	Left nasal ala mass	Hepatocellular carcinoma	Not described	Not described
Chau C, Siu W, Li MK.	54 /M	Hong kong	Nasal tip	Squamous cell carcinoma of esophagus	Not described	Not described
Cesinaro AM, Maiorana A, Annessi G, Collina G. 1995	40/M	United states of america	Small, hard nodule on right side dorsum of nose	Chordoma of sacrum	Not described	Not described

Table 1 Cont..

Gault DT, Subbuswamy SG. 1985	76/F	England	Blue black spherical mass over tip of nose	Oat cell carcinoma of lung	Surgical excision of nasal mass with palliative radiotherapy	Died in 3 months
Gault DT, Subbuswamy SG. 1985	67/M	England	Lump overlying the cartilage of left side of nasal tip	Squamous cell carcinoma of lung	X-ray therapy to chest	Not described
Nesi R, Lynfield Y. 1996	71/M	United states of america	Rhinophyma like lesion over nose	Squamous cell carcinoma of lung	*	Died with in a week
GAL TJ, KERSCHNER JE. 1997	65/M	United states of america	Tip of nose	Adenosquamous carcinoma of lung	Radiotherapy	Died several weeks later
De Simoni I et. al	64/M	SPAIN	Tip of nose	Squamous cell carcinoma of lung	Palliative radiotherapy	Died after 1 month
Viera Mota A et. Al	63/M	Portugal	Tip of nose	Anaplastic carcinoma of lung		Died after 2 weeks
Hammert WC, Champagne L, Heckler FR. 1999	59/M	United states	Tip of nose	Squamous cell carcinoma of lung	Chemoradiation with surgical resection of nasal mass	Not described
Rubinstein RY et. Al	64/M	United states	Mass over nose	Large cell undifferentiated carcinoma	*	*
Chun SM et. Al	62/M	Korea	Tip of nose	Squamous cell carcinoma of lung	Chemoradiation for primary	Died after 5 months
Chun SM et. Al	76/M	Korea	Tip of nose	Squamous cell carcinoma of lung	Palliative radiation therapy	Died after 1 month
Chun SM et. Al	57/M	Korea	Tip of nose	Squamous cell carcinoma of lung	chemoradiation	Died after 9 months



Fig. 1:




Fig. 2:

the literature till now including our case. Cutaneous metastasis from lung carcinoma may appear earlier than the diagnosis of primary malignancy. Renal cell carcinoma is considered the most common malignancy metastasizing



Fig. 3:




Fig. 4:

to the nose and paranasal sinuses. However on reviewing literature, lung carcinoma was seen to be the most common malignancy metastasizing to the nose tip.⁶ Among lung carcinomas, adenocarcinoma of lung is the most common type metastasizing to skin, followed by squamous cell carcinoma.⁷ The upper lobe tumours have more tendency of metastasizing to the skin.⁸ Squamous cell carcinoma was the most common malignancy of lung metastasizing to the tip of nose.⁹ It may appear as firm to rubbery with or without ulceration, blue to reddish in colour and may be single or multiple.^{10,11} Different theories are explained for its pathogenesis. Its spread may occur via the pulmonary vascular and lymphatic routes, the vertebral plexus, the arterial embolus and the venous sinuses of the skull which on communication with the vertebral system metastasize the tumour cells to the nose.^{12,13} Patient of internal malignancies presenting with cutaneous metastasis have poor prognosis with life expectancy of about 3-5 months.^[14-16]^{14,15} Nasal tip skin metastasis are also reported in literature from malignancies of thyroid, hypopharynx, esophagus, chordoma and leukemia.¹² There are different opinions regarding its management, ranging from wide surgical excision with reconstruction to chemoradiation. Due to very short life span of these patients, wide surgical excision with reconstruction doesn't seem

to be feasible. The tumour mass can be excised and raw area can be left for healing by secondary intention or patient can be given chemoradiation for it. On reviewing literature for nasal tip metastasis, we found that the age of cutaneous nasal metastasis varied from 40 years with chordoma to 86 years with renal cell carcinoma with a mean age of 63.45 ± 10.8 years. Out of 24 cases, only 6 were females(25%) and 18 were males(75%). Female dominance was seen in extrapulmonary primaries(45%) than pulmonary primaries(9%). Tip of nose was the most common site of metastasis in both pulmonary and extrapulmonary primaries. Histologically squamous cell carcinoma was the most common primary that metastasized to nasal tip. Survival time ranged from less than a week to 18 months with a mean life of 4.57 ± 5.45 months with mean survival life of metastasis from pulmonary primaries less than that of extrapulmonary.

3. Conclusion

Nose is a part of facial skeleton and due to its aesthetic value patients usually presents with lesions of nose while neglecting symptoms of internal malignancies. This usually delays the diagnosis and also lead to misdiagnosis in some cases. So it is the part of clinicians to examine internal systems in patients presenting with isolated malignancy of nose. Considering the low life expectancy of these patients, conservative approach should be taken for its management.

4. Source of Funding

None.

5. Conflict of Interest

None.

References

1. Rosen T. Cutaneous metastases. *Med Clin North Am.* 1980;64:885–900.
2. Brownstein MH, Helwig EB. Metastatic tumors of the skin. *Cancer.* 1972;29(5):1298–1307. doi:10.1002/1097-0142(197205)29:5<1298::aid-cnrcr2820290526>3.0.co;2-6.
3. Ledderose GJ, Reu S, Enghard AS, Krause E. Endonasal resection of early stage squamous cell carcinoma of the nasal vestibule. *Eur Arch Otorhinolaryngol.* 2014;271(5):1051–6. doi:10.1007/s00405-013-2660-4.
4. Lookingbill DP, Spangler N, Helm KF. Cutaneous metastases in patients with metastatic carcinoma: a retrospective study of 4020 patients. *J Am Acad Dermatol.* 1993;29(2):228–64. doi:10.1016/0190-9622(93)70173-q.
5. Krathen RA, Orengo IF, Rosen T. Cutaneous metastasis: a meta-analysis of data. *South Med J.* 2003;96(2):164–71. doi:10.1097/01.SMJ.0000053676.73249.E5.
6. Shindo M, Yoshida Y, Tominaga K, Yamamoto O. Skin Metastasis of Hypopharyngeal Carcinoma to the Nasal Tip. *Yonago Acta medica.* 2013;56(2):57–65.
7. Mollet TW, Garcia CA, Koester G. Skin metastases from lung cancer. *Dermatol Online J.* 2009;15(5):1.

8. Coslett LM, Katlic MR. Lung cancer with skin metastasis. *Chest*. 1990;97(3):757–66. doi:10.1378/chest.97.3.757.
9. Chun SM, Kim YC, Lee JB, Kim SJ. Nasal Tip Cutaneous Metastases Secondary to Lung Carcinoma: Three Case Reports and a Review of the Literature. *Acta Derm Venereol*. 2013;93(5):569–72. doi:10.2340/00015555-1529.
10. Dreizen S, Dhingra H, Chiuten D, Umsawasdi T, Valdivieso M. Cutaneous and subcutaneous metastases of lung cancer. *Postgrad Med*. 1986;80(8):111–7. doi:10.1080/00325481.1986.11699635.
11. Justribo MV, Seuma C, Portero JM, Egado L. Skin metastasis as the 1st manifestation of bronchogenic carcinoma. *Arch Bronconeumol*. 1994;30(6):314–20.
12. Itin PH, Heitzmann F, Stamm B. Metastasis to the nasal tip from a cervical carcinoma. *Dermatology*. 1999;199(2):171–4. doi:10.1159/000018231.
13. Kataoka A, Nishida T, Tomioka Y, Hirai N, Ohbuchi M, Yakushiji M. A metastasis to the nasal tip from a cervical carcinoma- a case report. *Kurume Med J*. 1998;45(1):127–58. doi:10.2739/kurumemedj.45.127.
14. Coslett L, Katlic M. Lung cancer with skin metastasis. *Chest*. 1990;97(3):757–9. doi:10.1378/chest.97.3.757.
15. Hammert WC, Champagne L, Heckler FR. Metastatic squamous cell carcinoma of the nasal tip: a case report. *J Oral Maxillofac Surg*;57(2):186–9. doi:10.1016/s0278-2391(99)90237-9.

Author biography

Jagat Singh, Professor

Usha Poonia, Senior Resident

Manisha Kumari, Senior Resident

Sukriti Bansal, Junior Resident

Aman, Consultant ENT Head

Cite this article: Singh J, Poonia U, Kumari M, Bansal S, Aman. Unusual presentation of lung carcinoma as nasal tip metastasis: A review of literature. *IP J Otorhinolaryngol Allied Sci* 2021;4(3):122-126.