



Original article

Knowledge, attitude and practice of Libyan medical students about vitamin D deficiency

Mustafa A. Alssageer^{1*}  , Naser M. Alaasswad² , Amma I. Jebril¹, Hajir A. Ahmed¹ and Roqia S. Almahdi¹

¹Department of Pharmaceutical care, Faculty of Pharmacy, Sebha University, Sebha, Libya

²Department of Clinical Biochemistry, Faculty of Medical Technology, Sebha University, Sebha, Libya

*Author to whom correspondence should be addressed

Received: 09-09-2022, Revised: 22-09-2022, Accepted: 27-09-2022, Published: 30-09-2022

Copyright © 2022 Alssageer et al. This is an open access article distributed under the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

HOW TO CITE THIS

Alssageer et al. (2022) Knowledge, attitude and practice of Libyan medical students about vitamin D deficiency.
Mediterr J Pharm Pharm Sci. 2 (3): 46 - 56. <https://doi.org/10.5281/zenodo.7115292>

Keywords: Attitude, knowledge, Libya, medical student, vitamin D deficiency

Abstract: The prevalence rate of vitamin D deficiency over the last decades has grown up rapidly worldwide among young adults. Therefore, the aim of this study is to explore knowledge, attitude and practice of Libyan medical students of Sebha University regarding vitamin D deficiency. Out of a total of 200 questionnaires, 148 completed questionnaires were returned (74.0% response rate). The majority of respondents did not measure their vitamin D levels (62.1%). Participants who indicated they exposing to sunlight for less than 15 min were 47.2% while 74.9% exposing to less than 30 min with highly significant difference in gender ($p < 0.001$). All the males did not use sunscreen creams compared with females (43.5%). Nearly two-thirds of the participants (61.4%) denied drink milk almost daily with significant differences based on the gender ($p < 0.005$) while 56.0 % of the participants reported eat fish, 60.8% of the students denied eating fast food and 74.3% eating eggs daily or weekly. However, 49.3% of the students are usually drinking of tea and coffee after meals, and do not usually practice exercise (53.3%), however, disagreed that their food has a low amount of calcium (30.4%). Most of the students agreed with the benefits of vitamin D (72.0%). Based on the assessment of awareness of the participants about vitamin D information, the average of correct answers percentage of the respondents was 47.1%. The most frequent sources of vitamin D information were the internet which accounted for 49.3%, followed by relatives and friends (31.7%). Some of the respondents (58.7%) reported that unawareness is the main cause of vitamin D deficiency. This study highlights the lack of awareness about the importance of vitamin D among medical students. This finding provides the institutions of medical education in Libya with an evidence base for a deficit knowledge regarding young health which could be utilized it to draw a strategy to correct health behaviors of students and community through medical education programs.

Introduction

Vitamin D is essential for intestinal calcium absorption and it plays a central role in maintaining calcium homeostasis and skeletal integrity [1].

Vitamin D is unique because it can be made in the skin from exposure to the sunlight [2]. Deficiency of vitamin D, or hypovitaminosis most commonly occurs in people when they have inadequate

sunlight exposure (particular sunlight with adequate ultraviolet B rays, UVB, and reducing in taking of foods with rich in vitamin D [3, 4]. Vitamin D deficiency has several complications which they associate with increases in mortality and adverse effects such as cardiovascular diseases, type 2 diabetes mellitus, obesity, metabolic syndrome, immune disorders and various types of cancer and mortality [5 - 7]. Individuals such as the elderly may have a high risk of vitamin D deficiency since they have concurrent other factors related to vitamin D deficiency, including decreased sunlight exposure, reduced intake of vitamin D in diet and decreased vitamin D absorption from sunlight due to decreasing in their skin thickness [8]. However, the prevalence rate of hypovitaminosis D is not restricted to the elderly and hospitalized population. Modernization has also brought the changes in the lifestyle and food habits of the young generation which may contribute to low dietary intake of vitamin D. The prevalence rate of vitamin D deficiency has grown up rapidly worldwide in adults during the past decades [9, 10]. In addition, some studies have reported a high prevalence rate of vitamin D deficiency among healthy, young adults [9 - 11]. Thus, in a Boston study, nearly two thirds of healthy, young adults were found to be vitamin D insufficient at the end of winter [11].

In Libya due to modernization and sudden rise in temperature, the numbers of hours spent outdoors have been decreased and sedentary way of working is most appreciated among urban people. Among Libyan population including young students may have a high risk of vitamin D deficiency. Recently, Alaasswad and others examined vitamin D status among pharmacy students of Sebha University which conducted concurrent with this study and some participant involved in this study [12]. The majority of pharmacy students in male and female blood donors have low vitamin D levels which represent 87.0% [12]. Inadequate awareness regarding vitamin D deficiency or non-adherence to the daily practice of the acquired knowledge or inappropriate exposure to the sun may also be the influential causes of deficiency of vitamin D. Therefore, this study was undertaken to explore

awareness, knowledge, attitude and practice of healthy medical students toward vitamin D deficiency in Libya.

Materials and methods

This is a cross-sectional study designed to determine vitamin D status among healthy young medical students at Sebha University, Sebha, Libya. It was carried out between January and April, 2020. All the medical students attending Sebha medical school were asked to participate voluntarily in the study ($n = 200$). The questionnaire was based on previous studies [13 - 15]. The survey contained 34 item questions that required varied response types. Most of the questions required specific categorical answers but some questions included other box to allow for more expressive answers. The questionnaire was translated to Arabic language form in order to clearly express for the participants. Verbal and written explanation of the study was provided when needed to the students for more detail expression. All the students were informed about the study and were required to read informative brochure to explain the purpose of the survey and the research. The investigator requested participant's verbal and written consent. They have also been informed that the questionnaire was anonymous and should be deposited in a box located in front of the classroom, in order to guarantee confidentiality. An ethics approval action for the study was obtained from the Sebha University Research Ethics Committee (2/2020).

Statistical analysis: All analysis were carried out by using Microsoft Office excel 2013 and SPSS statistical package the generated data presented into percentage, variant increase and decrease, mean and standard deviations were calculated. Paired *t*-test was used to compare between the individual groups. A probability value of less than 0.05 was considered as a level of significance difference.

Results

In this study, the questionnaires were handed to all the students who attended first years of pharmacy,

medicine and dentistry faculties. The returned questionnaires were 148 out of 200 which represents a response rate of 74.0% with notably more females than males, 83.7%, 16.3%, respectively. The majority of participants were female and aged between 18 and 20 years old accounted for 96.6%. Most of the participants denied to have diabetes mellitus (97.2%). A great

proportion of the participants reported that bone weakness is their family history (males: 29.1% and females: 45.1%). 62.1% did not measure their vitamin D levels compared with 35.1% who declared did. Only 02.7% did not answer the question. An equal rate of male and female students who reported did not measure their vitamin D levels, 62.5% and 62.4%, respectively.

Table 1: Data about responses of the medical students for factors related to vitamin D					
	Drinking of milk				Chi-square
	Yes	No		Total	P - value
Male	12 (50.0%)	12 (50.0%)	0	024	0.209
Female	45 (36.2%)	79 (63.7%)	0	124	
Total	57 (38.5%)	91 (61.4%)	0	148	
	Using sunscreen cream				
Male	00 (00.0%)	24 (100%)	0	024	0.000
Female	54 (43.5%)	70 (56.4%)	0	124	
Total	54 (36.4%)	94 (63.5%)	0	148	
	Eating fast foods				
Male	11 (45.8%)	13 (54.1%)	00 (00.0%)	024	0.327
Female	45 (36.2%)	77 (62.0%)	02 (01.6%)	124	
Total	56 (37.8%)	90 (60.8%)	02 (01.3%)	148	
	Eating fish				
Male	16 (66.6%)	07 (29.1%)	1	024	0.459
Female	67 (54.0%)	05 (04.0%)	0	124	
Total	83 (56.0%)	64 (43.2%)	1	148	
	Eating eggs daily or weekly				
Male	22 (91.6%)	02 (08.3%)	00 (00.0%)	024	0.035
Female	88 (70.9%)	34 (27.4%)	02 (01.6%)	124	
Total	110 (74.3%)	36 (24.3%)	02 (01.3%)	148	
	Drinking tea and coffee after meals				
Male	16 (66.6%)	08 (33.3%)	0	024	0.060
Female	57 (45.9%)	67 (54.0%)	0	124	
Total	73 (49.3%)	75 (50.6%)	0	148	
	Doing physical exercise				
Male	18 (75.0%)	06 (25.0%)	00 (00.0%)	024	0.001
Female	48 (38.7%)	73 (58.8%)	03 (02.4%)	124	
Total	66 (44.5%)	79 (53.3%)	03 (02.0%)	148	
	Diabetes mellitus				
Male	01 (04.1%)	023 (95.8%)	0	024	0.099
Female	01 (00.8%)	121 (97.5%)	0	124	
Total	02 (02.7%)	144 (97.2%)	0	148	
	Poverty of food with calcium source				
Male	05 (20.8%)	13 (54.1%)	06 (25.0%)	024	0.009
Female	71 (57.2%)	32 (25.8%)	21 (16.9%)	124	
Total	76 (51.3%)	45 (30.4%)	27 (18.2%)	148	
	Family history of bone weakness				
Male	07 (29.1%)	16 (66.6%)	01 (04.1%)	024	0.542
Female	56 (45.1%)	54 (43.5%)	14 (11.2%)	124	
Total	63 (42.5%)	70 (47.2%)	15 (10.1%)	148	
	Family history of vitamin D deficiency				
Male	11 (45.8%)	09 (37.5%)	04 (16.6%)	024	0.471
Female	70 (56.4%)	35 (28.2%)	19 (15.3%)	124	
Total	81 (54.7%)	44 (29.7%)	23 (15.5%)	148	

Almost half of the participants (47.2%) have exposure to sunlight less than 15 mints per day and 74.9% have exposure to less than 30 min. On the contrary, 24.9% reported to expose to light for more than 30 min. Thus, an analysis by Chi-square test revealed high significant difference between males and females to exposure to sunlight ($p < 0.01$). Half of the females declared to usually expose to sunlight (52.4%) for less than 15 minutes compared with males (20.8%). 36.4% of the total participants used sunscreen products. However, all the male students did not use sunscreen creams compared with 43.5% of female students who declared using sunscreen products (**Table 1**). In **Table 1**, 61.4% denied drink milk every day. Female students have a higher rate of drinking of milk than males, 63.7% and 50.0%, respectively. Females who denied drinking almost every day were as double as the males with no significant difference. 60.8% denied eating fast food. Female students were less likely to eat fast foods compared with males. In the same way, 45.8% of the males declared eating fast foods compared with females (37.8%). 56.0% eat fish and males eating more than females, 66.6% and 54.0%, respectively. The majority of male and female students usually eat

eggs whether daily or weekly (74.3%). This habit was more dominant in males (91.6%) than females (70.9%) with a significant difference of $p < 0.035$. About half of the students (49.3%) usually drinking tea and coffee after meals but 66.6% of the males reported yes and half of females denied do that regularly (54.0%).

In **Figure 1**, male students (54.1%) denied their food has a poor quantity of calcium compared with 25.1% of females. 30.4% disagreed that their food has a low amount of calcium. This difference is significant between genders ($p < 0.01$). **Table 1** shows that three quarters of male students do regular exercise. Thus, students do not usually practice physical exercise (53.3%) with highly significant difference ($p < 0.001$). Family history of vitamin D deficiency was found to be 54.7%. Females were more likely to have a family deficiency of vitamin D than males (56.4% and 45.8%, respectively). Most of the students' answers that agreed that vitamin D is used for the treatment of bone disorders and rickets (71.6%) and only 02.7% disagreed, but 25.6% do not know in their answers.




Figure 1: Libyan Medical participant's responses to benefits of vitamin D

Figure 1 shows the respondents agree that vitamin D is important of save the level of calcium and phosphate. However, only 03.37% disagreed and all of them were solitary females. Three quarters of the respondents agreed that vitamin D can protect the bone and teeth compared with 09.4% who disagreed (female students). 69.5% agreed that

vitamin D can enhance the body's immunity. 70.9% of females slightly more agreed on the statement compared with 62.5% males with 23.0% could not answer this question. Also, in **Figure 1**, 66.2% agreed that vitamin D can improve muscle strength compared to who disagreed (05.4%) and 28.3% could not answer this question.

Table 2: Knowledge of the Libyan participants about vitamin D deficiency

Questions	Yes	No	IDN
Do you agree that the incidence of vitamin D deficiency in Libya more than other countries	42 (28.3%)	21 (14.1%)	85 (57.4%)
Does the sun is the main source of vitamin D	130 (87.8%)	09 (06.0%)	09 (06.0%)
Does the Osteoporosis is one of the symptoms of vitamin D deficiency?	111 (75.0%)	13 (08.7%)	24 (16.2%)
Can vitamin D deficiency be replenished by eating dairy products?	52 (35.1%)	41 (27.7%)	55 (37.1%)
Does the vitamin D deficiency cause depression?	86 (58.1%)	11 (07.4%)	51 (34.4%)
Dose the level of vitamin D in the body differ by change in time and seasons?	64 (43.2%)	08 (05.4%)	76 (51.3%)
Does the vitamin D sources are found in animals but not in Vegetables and fruits?	15 (10.1%)	83 (56.0%)	50 (33.8%)
Are the people who live in cloudy areas more susceptible to vitamin D deficiency?	87 (58.7%)	13 (08.7%)	48 (32.4%)
Does frequent exposure to the sun not lead to vitamin D toxicity?	27 (18.2%)	43 (29.0%)	78 (52.7%)
Could the use of sunscreen be a cause of vitamin D deficiency?	34 (22.9%)	41 (27.7%)	73 (49.3%)
Could a fat-free diet be a cause of vitamin D deficiency?	46 (31.0%)	31 (20.9%)	71 (47.9%)
Dark skin more susceptible to vitamin D deficiency than light skin?	18 (12.1%)	54 (36.4%)	76 (51.3%)
Can vitamin D toxicity cause hypercalcemia?	29 (19.5%)	17 (11.4%)	102 (68.9%)
In the case of vitamin D deficiency, taking pharmaceutical preparations containing vitamin D is more effective compared to foods, diet and exposure to sunlight.	75 (50.6%)	37 (25.0%)	36 (24.3%)
Taking vitamin D supplements (medications) is wrong unless prescribed by a doctor.	107 (72.2%)	20 (13.5%)	21 (14.1%)
Taking vitamin D supplements is necessary to treat vitamin D deficiency, but not to prevent it.	90 (60.8%)	19 (12.8%)	39 (26.3%)
Average percentage of correct answers of the respondents is 47.1%.			

In **Table 2**, 57.4% did not know the incidence of vitamin D deficiency in Libya more than in other countries while 28.3% agreed with this statement compared with 14.1% who disagreed. 87.8% believed that sun is the main source of vitamin D compared with 06.0% in each of those who disagreed with the statement and those who could not answer the question. Also, in **Table 2**, the majority agreed that osteoporosis is one of the symptoms of vitamin D deficiency compared only 08.7% who disagreed. However, 16.2% chose they do not know. Furthermore, 35.1% agreed with the statement that vitamin D can be replenished by eating dairy products compared with 27.7% who disagreed. 58.1% agreed with the statement vitamin D deficiency can cause depression

compared with only 07.4% who disagreed. Over half of the respondents could not answer the question that levels of vitamin D differ by time and seasons changes. 05.4% disagreed with the statement. 10.1% agreed with the statement that vitamin D is found in animal sources rather than vegetable or fruit sources while 56.0% contradict with this statement, but 33.8% could not answer the question. 58.7% agreed with the statement that people who live in cloudy areas are more susceptible to vitamin D deficiency compared with 08.7% who disagreed and 32.4% could not answer this question. Only 12.1% agreed with the statement that dark skin is more susceptible to vitamin D deficiency than light skin compared with 36.4% who disagreed with the same statement and

36.4% could not answer this question. As shown in **Table 2**, 50.6% agreed on the question that taking pharmaceutical preparation is more effective than foods, diet and exposure to sunlight and they represent as double as respondents (25.0%) who disagreed with this statement with 24.3% could not answer this question. Furthermore, 72.2% agreed with the statement that vitamin D supplements is wrong unless prescribed by a physician while a minority of respondent disagreed and could not answer the question represented by 13.5% and 14.1%, respectively. 60.8% agreed that a vitamin D supplement is necessary to treat vitamin D deficiency rather than to prevent. 76.3% knew the disease of vitamin D deficiency while 16.2% did not have knowledge about this disease. Females (77.4%) were more declared that have knowledge about the disease than males (70.8%) and 07.4% were unable to answer this question. The highest

rate of source of information among respondents was the internet which accounted for 49.3% followed by relatives and friends (31.7%). To less extent, respondents chose medical providers and media by 16.2% and 10.1%, respectively. 12.1% would not receive information about vitamin D compared with 82.4% who would like to get this information. Also, in **Figure 2**, respondents who disagreed in males more than females (25.0% and 09.6%, respectively). The highest rate of causes reported by respondents (58.7%) was unawareness of the risk of vitamin D deficiency followed by medication (53.3%). To less extent, imbalance in nutrition was represented by 34.4%. Thus, **Figure 2** shows respondents reported a lifestyle and less exposure to sun rays, concomitant disease and others reported by 11.5%, 07.4% and 10.1%, respectively.




Figure 2: Causes of prevalence of vitamin D deficiency among Libyan participants

Discussion

Vitamin D can be made in the skin from exposure to sunlight [16]. The major source of vitamin D is due to synthesis of vitamin D from skin exposure to ultraviolet B radiation which leads to the conversion of 7-dehydrocholesterol to pre-vitamin D3 and rapidly converted to vitamin D3 [17]. Regarding its role, it suggests a low prevalence of vitamin D deficiency in Mediterranean Sea countries. This study reports the medical students

have exposure to sunlight for less than 30 minutes. Males had a higher average compared with females exposing themselves to sunlight. In a Chinese study, 62.3% declared do not like going in the sun [15]. Exposure peripherals as legs and arms for 15 min twice a week was reported to be enough for an adequate sun-induced cutaneous vitamin D synthesis in adolescents [17]. In veiled Arab women, vitamin D deficiency is a result of a combination of limitations in sunlight exposure and

low oral intake of vitamin D [18]. The current evidence showed wearing concealing clothes is associated with vitamin D deficiency irrespective of race or strength of solar radiation [18, 19]. The influence of clothing style on vitamin D status was the subject of several previous studies [20-22]. All Libyan females wearing hijabs. Thus, cover most of the body when women get outdoor activities, can decrease skin opportunity to exposure to direct sunlight. Hypovitaminosis of vitamin D less prevalence among individuals who staying indoor more time, wearing clothes cover wide area of the skins and their skin pigmentation is darker [23]. In Sudan, one third of female medical students used sunscreen [24]. Using of sunscreens could result in vitamin D deficiency since they have a protective effect against ultraviolet B. Using sunscreen with sun protection factor of 30 reduces vitamin D synthesis in the skin [3, 25]. This differs from China's study which reveals 85.0% of the participants used some sun protection and 33.0% of males used sunscreen compared with females [26]. Reason behind this variation could be related to the difference in geographical, socioeconomic and educational factors. A finding of study of medical students in Pakistan was in accord with our study, in which the female students have greater proportion using sunscreen than males [14]. This could be elucidated that males believe that sunscreen is not for masculine and it is a product intended for feminine since they are not highly concerned with skincare as remarkable with females [12]. Healthy young adults may develop vitamin D deficiency when vitamin D intake is below the recommended intake [28]. The intake of diet rich in vitamin D is important for maintaining body vitamin D hemostasis. Dietary supplements are useful to prevent and treat this deficiency [29, 30]. The present study revealed the participants' consumption frequency of some foods rich in vitamin D was low and participants denied usually drink milk. The UK retail study showed vitamin D₃ and 25(OH)D₃ concentrations of eggs were significantly different depending on egg production systems [31]. Evidence demonstrated that vitamin D in eggs is increased in birds exposed to ultraviolet radiation [31]. Performing physical

activity in outdoor environment with sun exposure would provide benefits from the physical work itself and from vitamin D synthesis and action in the body [32]. This finding is slightly higher in Libyan study in which reported that 44% of the adults do not get sufficient exercise [18]. Women are inclined to follow a sedentary lifestyle compared with men where they spend much time at home [33]. Physical inactivity among other factors such as obesity, low vitamin D dietary and supplement intake are major modifiable predictors of low vitamin D status [34].

Vitamin D deficiency relates to diseases as diabetes mellitus [35]. A meta-analysis study indicated risk of developing diabetes mellitus decreases by 04.0% with each 10 nmol/L increase in 25(OH) D [36]. Nutrition examination survey reported an inverse relationship between 25(OH) D and glycated hemoglobin (HbA1c) levels aged people (35 - 74 years) with no history of diabetes mellitus [37]. Previous studies have revealed genetic variants related to vitamin D hemostasis and strongly depend on endogenous vitamin D production, which influenced by genetic determinants [38]. Twin and family studies have shown high vitamin D heritability [39, 40]. Ethnic differences in prevalence of common genetic polymorphisms are another likely explanation for low vitamin D in African Americans [41]. Among the ethnic groups in Malaysia, Malay and Indian females were six fold times more to have vitamin D deficiency compared to Chinese females [38]. The majority of students agreed that vitamin D use for treatment of bone disorders and rickets, importance of save the level of calcium and phosphate, protect the bone and teeth, enhance the body immunity and improve the muscle strength. This is in good line with Babelghaith et al. [42]. The high awareness toward benefits vitamin D deficiency could be related to university students and particularly medical students which might behave higher education in health subjects. Brand et al. [43] reported older age appeared to be associated with a lower perception of risk of vitamin D deficiency. Many health-care professionals are not fully aware of the benefits of vitamin D to public health [44]. A similar trend is

reported in Pakistan and Australia studies [45, 46]. This awareness about benefits of vitamin D is important to encourage people and university students to influence their behaviors toward vitamin D practice. Over three quarters of the participants knew vitamin D deficiency disease which is similar to previously published study [47]. This could be elucidated that the respondents received some information but were still unsatisfied with the amount of information they have. This aligned with study conducted in Bahrein on the adult population [48]. The majority had heard about vitamin D despite the low level of awareness of sources and the role of vitamin D [15]. This poor knowledge could be due to conflicting information available from various sources about vitamin D. In contrast, 02.0% reported that internet is the source of their knowledge and information about calcium and vitamin D [49]. In UK, the most common source of vitamin D information is the media followed by friends and family [50]. In Saudi Arabia, information source was healthcare providers followed by friends [42]. It is not surprising that evidence also reported that healthcare workers lack a high knowledge of vitamin D deficiency [51]. The knowledge of medical students with regard to some direct and simple questions generally scored as good knowledge. For example, the majority believed that sun is the main source of vitamin D and recognize the influence of vitamin D deficiency can cause osteoporosis. This finding is in line with Indian study conducted on female medical students that the majority of the participants correctly pointed out the effect of vitamin D on bone health [13]. The statement that vitamin D is found in animal sources but not in vegetable or fruit sources was high in our study. However, fruits were the main source of vitamin D among university students followed by vegetables participants [45]. 20.0% of pharmacy students failed to mention at least one vitamin D rich product [49]. Over half of respondents agreed that taking pharmaceutical preparation is more effective than foods, diet and exposure to sunlight. However, participants had poor knowledge about

toxicities and pharmacokinetics of vitamin D management. Over quarter of respondents rejected the statement that exposure to the sun leads to the toxicity of vitamin D. A minority of respondents disagreed with the same statement can vitamin D toxicity causes hypercalcemia. Lack of awareness will increase this problem while adequate knowledge about vitamin D deficiency and its prevention can cause potential decrease in the disease burden. Similarly, in Sudan, half of female medical students stated sunscreen does not interfere with vitamin D synthesis in the skin [24]. The highest rate of causes of hypovitaminosis D among the Libyan population was unawareness of people about the risk of vitamin D deficiency. This consistency with finding that lack of awareness about importance of vitamin D, its health benefits and prevention of deficient states across populations as important factors for the spread of vitamin D deficiency [52, 53]. Lack of knowledge about vitamin D and negative attitude toward sunlight exposure is prevalent among university students [45, 54]. Ironically, in this study, a lifestyle and less exposure to sunlight can causes vitamin D deficiency which is in contrast to the fact that the worldwide pandemic of hypovitaminosis D mainly is attributed to lifestyle and environmental factors that reduce exposure to sunlight [55]. As consequence, the sedentary lifestyle of individuals who has fewer physical activities such as female gender due to sociocultural or indoor job be liable to exposure to sunlight at a lower rate since they have limited time for outdoor activities.

Conclusion: This study presents poor practice, inadequate knowledge and low awareness of the importance of vitamin D deficiency among medical students of the University. The majority have an adequate understanding of the benefits of vitamin D. This study provides Libyan institutions of medical education an evidence for a deficit knowledge which could be utilized as a strategy to correct the health behaviors through medical education programs.

Acknowledgments: The authors would like to thank all the participants for their corporation that facilitates this study.

Author contributions: All the authors substantially contributed to the conception, compilation of data, checking and approving the final version of the manuscript and agreed to be accountable for its content.

Author declarations: The authors confirm that all relevant ethical guidelines have been followed and any necessary IRB and/or ethics committee approvals have been obtained.

Conflict of interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Data availability statement: The raw data that support the findings of this article are available from the corresponding author upon reasonable request.

Ethical issues: Including plagiarism, informed consent, data fabrication or falsification and double publication or submission have completely been observed by authors.

References

1. Cashman KD (2007) Calcium and vitamin D. Novartis Foundation Symposium. 282:123-138, PMID: 17913228.
2. Lips P, Hosking D, Lippuner K, Norquist JM, Wehren L, Maalouf G, Ragi-Eis S, Chandler J (2006) The prevalence of vitamin D inadequacy amongst women with osteoporosis: an international epidemiological investigation. *Journal of Internal Medicine*. 260 (3): 245-254. doi:10.1111/j.1365-2796.2006.01685.x.
3. Holick MF, Chen TC (2008) Vitamin D deficiency: a worldwide problem with health consequences. *American Journal of Clinical Nutrition*. 87 (4): 1080S-1086S. doi:10.1093/ajcn/87.4.1080S.
4. Prentice A (2008) Vitamin D deficiency: a global perspective. *Nutrition Review*. 66 (10, 2S): S153-164. doi:10.1111/j.1753-4887.2008.00100.x.
5. Adams JS, Hewison M (2010) Update in vitamin D. *Journal of Clinical Endocrinology and Metabolism*. 95 (2): 471-478. doi:10.1210/jc.2009-1773.
6. Correale J, Ysraelit MC, Gaitán MI (2009) Immunomodulatory effects of vitamin D in multiple sclerosis. *Brain*. 132 (5):1146-1160. doi:10.1093/brain/awp033.
7. Bener A, Al-Hamaq AO, Saleh NM (2013) Association between vitamin D insufficiency and adverse pregnancy outcome: global comparisons. *International Journal of Womens Health*. 5: 523-531. doi:10.2147/IJWH.S51403.
8. Janssen HCJP, Samson MM, Verhaar HJJ (2002) Vitamin D deficiency, muscle function, and falls in elderly people. *American Journal of Clinical Nutrition*. 75 (4): 611-615. doi:10.1093/ajcn/75.4.611.
9. Bischoff-Ferrari HA, Giovannucci E, Willett WC, Dietrich T, Dawson-Hughes B (2006) Estimation of optimal serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *American Journal of Clinical Nutrition*. 84 (1): 18-28. doi:10.1093/ajcn/84.1.18.
10. Rajpathak SN, Rimm EB, Rosner B, Willett WC, Hu FB (2006) Calcium and dairy intakes in relation to long-term weight gain in US men. *American Journal of Clinical Nutrition*. 83 (3): 559-566. doi:10.1093/ajcn.83.3.559.
11. Tangpricha V, Pearce EN, Chen TC, Holick MF (2002) Vitamin D insufficiency among free-living healthy young adults. *American Journal of Medicine*. 112 (8): 659-662. doi:10.1016/s0002-9343(02)01091-4.
12. Alaasswad NM, Jebril AI, Ahmed HA, Almahdi RS, Alsageer MA (2022) Vitamin D deficiency and anemia among pharmacy students. *Mediterranean Journal of Pharmacy and Pharmaceutical Sciences*. 2 (2): 90-96. doi: 10.5281/zenodo.6780515.
13. Nowreen N, Hameed R (2019) Awareness regarding the importance of vitamin D and prevention of its deficiency among female undergraduate medical students. *International Journal Basic and Clinical Pharmacology*. 8 (5): 865-868. doi:10.18203/2319-2003.ijbcp20191563.
14. Juanid R, Feroz S, Mugh A (2019) Knowledge, attitude and practice of medical students regarding vitamin D. *Journal of Rawalpindi Medical College*. 23 (S-2): 98-103.
15. Kung AWC, Lee K-K (2006) Knowledge of vitamin D and perceptions and attitudes toward sunlight among Chinese middle-aged and elderly women: a population survey in Hong Kong. *BMC Public Health*. 6 (1): 226. doi:10.1186/1471-2458-6-226.
16. MacKeigan LD, Larson LN (1989) Development and validation of an instrument to measure patient satisfaction with pharmacy services. *Medical Care*. 27 (5): 522-536. doi:10.1097/00005650-198905000-00007.
17. Holick MF (2004) Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *American Journal of Clinical Nutrition*. 80 (6S): 1678S-88S. doi:10.1093/ajcn/80.6.1678S.

18. Glerup H, Mikkelsen K, Poulsen L, Hass L, Overbeck S, Thomsen J, Charles P, Eriksen EF (2000) Commonly recommended daily intake of vitamin D is not sufficient if sunlight exposure is limited. *Journal of Internal Medicine*. 247 (2): 260-268. doi.org/10.1046/j.1365-2796.2000.00595.x.
19. Gannagé-Yared M-H, Chemali R, Yaacoub N, Halaby G (2000) Hypovitaminosis D in a sunny country: relation to lifestyle and bone markers. *Journal of Bone and Mineral Research*. 15 (9): 1856-1862. doi.org/10.1359/jbmr.2000.15.9.1856.
20. Sedrani SH, Elidrissy AW, El Arabi KM (1983) Sunlight and vitamin D status in normal Saudi subjects. *American Journal of Clinical Nutrition*. 38 (1): 129-132. doi:10.1093/ajcn/38.1.129.
21. El-Sonbaty MR, Abdul-Ghaffar NU (1996) Vitamin D deficiency in veiled Kuwaiti women. *European Journal of Clinical Nutrition*. 50 (5): 315-318. PMID: 8735313.
22. Hashemipour S, Larijani B, Adibi H, Javadi E, Sedaghat M, Pajouchi M, Soltani A, Shafaei AR, Hamidi Z, Fard ARK, Hossein Nezhad A, Booya F (2004) Vitamin D deficiency and causative factors in the population of Tehran. *BMC Public Health*. 4: 38. doi:10.1186/1471-2458-4-38.
23. Holvik K, Brunvand L, Brustad M, Meyer HE (2008) Vitamin D status in the Norwegian population. *Solar Radiation and Human Health*. Oslo: The Norwegian Academy of Science and Letters, 2008.
24. Ali SM, Salih LMA, Saeed E (2019) Awareness of medical students about vitamin D deficiency at Ahfad University for women, Sudan. *Sudan Journal of Paediatrics*. 19 (2): 117-125. doi:10.24911/SJP.106-1574764595.
25. Matsuoka LY, Ide L, Wortsman J, MacLaughlin JA, Holick MF (1987) Sunscreens suppress cutaneous vitamin D3 synthesis. *Journal of Clinical Endocrinology and Metabolism*. 64 (6): 1165-1168. doi:10.1210/jcem-64-6-1165.
26. Zhou M, Zhuang W, Yuan Y, Li Z, Cai Y (2016) Investigation on vitamin D knowledge, attitude and practice of university students in Nanjing, China. *Public Health Nutrition*. 19 (1): 78-82. doi:10.1017/S1368980015000373.
27. Abroms L, Jorgensen CM, Southwell BG, Geller AC, Emmons KM (2003) Gender differences in young adults' beliefs about sunscreen use. *Public Health Education*. 30 (1): 29-43. doi:10.1177/1090198102239257.
28. Misra M, Tsai P, Anderson EJ, Hubbard JL, Gallagher K, Soyka LA, Miller KK, Herzog DB, Klibanski A (2006) Nutrient intake in community-dwelling adolescent girls with anorexia nervosa and in healthy adolescents. *American Journal of Clinical Nutrition*. 84 (4): 698-706. doi:10.1093/ajcn/84.4.698.
29. Stechschulte SA, Kirsner RS, Federman DG (2009) Vitamin D: bone and beyond, rationale and recommendations for supplementation. *American Journal of Medicine*. 122 (9): 793-802. doi.org/10.1016/j.amjmed.2009.02.029.
30. Henderson L, Gregory J, Swan G (2003) The national diet and nutrition survey: adults aged 19 to 64 years. *Vitamin Mineral intake. Urine Analysis*. 3: 1-8. ISBN: 0116215666
31. Guo J, Kliem KE, Lovegrove JA, Givens DI (2017) Effect of production system, supermarket and purchase date on the vitamin D content of eggs at retail. *Food Chemistry*. 221: 1021-1025. doi:10.1016/j.foodchem.2016.11.060.
32. Bjørgen K (2016) Physical activity in light of affordances in outdoor environments: qualitative observation studies of 3-5 years olds in kindergarten. *Springerplus*. 5 (1): 950. doi:10.1186/s40064-016-2565-y.
33. Altajori NN, ElshrekYM (2017) Risk factors for non-communicable diseases in Libya. *Egyptian Journal of Hospital Medicine*. 66 (1): 202-214. doi:10.12816/0034654.
34. Brock K, Huang W-Y, Fraser DR, Tseng M, Solomon RS, Peters U, Ahn J, Purdue M, Mason RS, McCarty C, Zeigler RG, Graubard B (2010) Low vitamin D status is associated with physical inactivity, obesity and low vitamin D intake in a large US sample of healthy middle-aged men and women. *Journal of Steroid Biochemistry and Molecular Biology*. 121 (1): 462-466. doi.org/10.1016/j.jsbmb.2010.03.091.
35. Ahmadieh H, Azar ST, Lakkis N, Arabi A (2013) Hypovitaminosis D in patients with type 2 diabetes mellitus: a relation to disease control and complications. *ISRN Endocrinology*. 2013: 641098. doi:10.1155/2013/641098.
36. Song Y, Wang L, Pittas AG, Del Gobbo LC, Zhang C, Manson JE, Hu FB (2013) Blood 25-hydroxy vitamin D levels and incident type 2 diabetes: a meta-analysis of prospective studies. *Diabetes Care*. 36 (5): 1422-1428. doi:10.2337/dc12-0962.
37. Kositsawat J, Freeman VL, Gerber BS, Geraci S (2010) Association of A1C levels with vitamin D status in U.S. adults: data from the National Health and Nutrition Examination Survey. *Diabetes Care*. 33 (6): 1236-1238. doi:10.2337/dc09-2150.
38. Wang TJ, Zhang F, Richards JB, Spector TD (2010) Common genetic determinants of vitamin D insufficiency: a genome-wide association study. *Lancet (London, England)*. 376 (9736): 180-188. doi:10.1016/S0140-6736(10)60588-0.
39. Karohl C, Su S, Kumari M, Tangpricha V, Veledar E, Vaccarino V, Raggi P (2010) Heritability and seasonal variability of vitamin D concentrations in male twins. *American Journal of Clinical Nutrition*. 92 (6): 1393-1398. doi:10.3945/ajcn.2010.30176.

40. Fuleihan GE, Bouillon R, Clarke B, Chakhtoura M, Cooper C, McClung M, Singh RJ (2015) Serum 25-Hydroxyvitamin D levels: variability, knowledge gaps, and the concept of a desirable range. *Journal of Bone and Mineral Research*. 30. doi:10.1002/jbmr.2536.
41. Powe CE, Evans MK, Wenger J, Zonderman AB, Berg AH, Nalls M, Tamez H, Zhang D, Bhan I, Karumanchi SA, Powe NR, Thadhani R (2013) Vitamin D-binding protein and vitamin D status of black Americans and white Americans. *New England Journal of Medicine*. 369 (21): 1991-2000. doi:10.1056/NEJMoA1306357.
42. Babelghaith SD, Wajid S, Al-Zaaqi MA, Al-Malki AS, Al-Amri FD, Alghadeer SA, Al-Arifi M (2017) Knowledge and practice of vitamin D deficiency among people lives in Riyadh, Saudi Arabia-A cross-sectional study. *An International Journal of Medical Sciences*. 28 (7): 3114-3118. Corpus ID: 28469126.
43. Brand CA, Abi HY, Couch D, Vindigni A, Wark JD (2008) Vitamin D deficiency: a study of community beliefs among dark skinned and veiled people. *International Journal of Rheumatic Diseases*. 11 (1): 15-23. doi.org/10.1111/j.1756-185X.2008.00323.x .
44. Khalsa S (2009) *Vitamin D revolution: how the power of this amazing vitamin can change your life*. (Inc. HH, ed.), pp-192. ISBN: 978-1-4019-2470-6. Publisher: Hay House, USA.
45. Tariq A, Khan SR, Basharat A (2020) Assessment of knowledge, attitudes and practice towards vitamin D among university students in Pakistan. *BMC Public Health*. 20 (1): 355. doi:10.1186/s12889-020-8453-y.
46. Vu LH, van der Pols JC, Whiteman DC, Kimlin MG, Neale RE (2010) Knowledge and attitudes about vitamin D and impact on sun protection practices among urban office workers in Brisbane, Australia. *Cancer Epidemiology, Biomarkers and Prevention*. 19 (7): 1784-1789. doi:10.1158/1055-9965.EPI-10-0127.
47. Iemu E, Varnam R (2012) Awareness of vitamin D deficiency among at-risk patients. *BMC Research Notes*. 5 (1): 17. doi:10.1186/1756-0500-5-17.
48. Sanad HM, Priya G, Mukhaimer JJ, Asokan GV, Belal S (2020) Knowledge and awareness of vitamin D deficiency among the general adult population in BAHRAIN: a cross-sectional study. *European Journal of Molecular and Clinical Medicine*. 7 (9): 180-191.
49. Uddin R, Huda NH, Jhanker YM, Jesmeen T, Imam MZ, Akter S (2013) Awareness regarding the importance of calcium and vitamin D among the undergraduate pharmacy students in Bangladesh. *BMC Research Notes*. 6: 134. doi:10.1186/1756-0500-6-134.
50. O'Connor C, Glatt D, White L, Iniesta RR (2018) Knowledge, attitudes and perceptions towards vitamin D in a UK adult population: A cross-sectional study. *International Journal of Environmental Research and Public Health*. 15 (11): 2387. doi:10.3390/ijerph15112387.
51. Al-Amri F, Gad A, Al-Habib D, Ibrahim AK (2017) Knowledge, attitude and practice regarding vitamin D among primary health care physicians in Riyadh city, Saudi Arabia, 2015. *World Journal of Food and Scientific Technology*. 1 (2): 47-55. doi: 10.11648/j.wjfst.20170102.13.
52. Mithal A, Wahl DA, Bonjour J-P, Burckhardt P, Dawson-Hughes B, Eisman JA, El-Hajj Fuleihan G, Josse RG, Lips P, Morales-Torres J (2009) Global vitamin D status and determinants of hypovitaminosis D. *Osteoporosis International*. 20 (11): 1807-1820. doi:10.1007/s00198-009-0954-6.
53. Hagenau T, Vest R, Gissel TN, Poulsen CS, Erlandsen M, Mosekilde L, Vestergaard P (2009) Global vitamin D levels in relation to age, gender, skin pigmentation and latitude: an ecologic meta-regression analysis. *Osteoporosis International*. 20 (1): 133-140. doi:10.1007/s00198-008-0626-y.
54. Arora H, Dixit V, Srivastava N (2016) Evaluation of knowledge, practices of vitamin D and attitude toward sunlight among Indian students. *Asian Journal of Pharmaceutical and Research*. 9 (1): 308-313.
55. Nair R, Maseeh A. Vitamin D (2012) The “sunshine” vitamin. *Journal of Pharmacology and Pharmacotherapy*. 3 (2): 118-126. doi:10.4103/0976-500X.95506.