

Diode laser or scalpel: A comparison of the two “Cutting Edge” surgical tools in maxillofacial surgery

¹Arunima Gupta, ²Ankur Mittal, Smiti Jassar Klaire ³Siddharth Sharma, ⁴Birsubhra Roy*, ⁵Md Tousif Alam⁶

^{1,6}PG Student, ²Professor and Head, ³Profesor ⁴Reader ⁵Senior Lecturer, ¹⁻⁶Dept of Oral & Maxillofacial Surgery, ¹⁻⁶D. J. College of Dental Sciences & Research, Modinagar, Uttar.Pradesh, India

Corresponding Author: Birsubhra Roy

Email-id: bir.ronie@gmail.com

Abstract

Context and Objective: Scalpel has been the gold standard tool for intraoral surgery. Since a few decades, lasers have also stepped up as operating instruments, be it cutting, ablation or photobiomodulation. In our research, we compared the efficacy of scalpel and diode laser in intraoral maxillofacial surgery.

Setting and Design: In the duration of 3 years, a total of 100 patients were included in our study which were divided into two groups of 50 patients each; 50 in the test group for scalpel and 50 in the control group for laser.

Materials and Methods: Inclusion criteria includes clinically diagnosed intraoral soft tissue lesions/growth indicative of tissue ablation, incision, excision, or biopsy. Exclusion criteria: Non consenting and/or non-cooperative patients. Presence of uncontrolled or advanced systemic diseases, immunocompromising diseases, porphyria or medications that cause photosensitivity (eg. Chloroquine, methotrexate etc) and serious eye defects.

Results: While laser provided a clean surgical field by facilitating haemostasis during bleeding, its healing time was slower as compared to laser during the first postoperative week. 2% Lignocaine HCl with 1:80000 adrenaline was used as a local anaesthetic agent. Swelling and redness were both found to be pronounced in scalpel as opposed to laser which produced a coagulum ring of eschar formation during cutting.

Discussion: Scalpel offers unmatched precision and speed during surgery. Its only drawback is bleeding which may be inconvenient for the surgeon. Laser not only clears the surgical field of the operator but also reduces the risk of bleeding from highly vascular areas like the tongue. The heat from the laser tip negates any additional measures required for cleaning.

Summary and Conclusions: The 980 nm diode laser promises versatility and surgical efficacy during intraoral maxillofacial procedures and can be a good alternative to the conventional cold scalpel, offering equal patient satisfaction.

Keywords: Immunocompromising diseases, Postoperative swelling, Maxillofacial surgery

Introduction

For millennia, surgical knives were single-piece instruments for which maintaining the sharpness of the blade was a constant problem. In 1910, Murphy began experimenting with a replaceable blade, but the process was cumbersome. In 1915, Morgan Parker, a 22-year-old engineer, patented a locking scalpel handle and blade system that was more secure and enabled an easier exchange of the blades.¹

Scalpel, also known as cold knife has been considered a gold standard cutting tool till now. Traditionally, it was considered the surgical cutting tool of choice because of its precision control, preservation of tissue integrity, and superior association with wound healing. But its main disadvantages like excessive bleeding rendered inadequate visibility in the operating field and non-sterilizing incision cut.²

Therefore, this necessitated the usage of other alternatives for surgery. A more recent surgical tool, the laser, an acronym for light amplification by stimulated emission of radiation, is a device for generating a high-intensity, ostensibly parallel beam of monochromatic (single wavelength) electromagnetic radiation. The possibility of stimulated emission was predicted by Einstein in 1917; based on the work of Gordon in 1955 and Shawlow and Townes in 1958, Maiman created the first operational laser in 1960, a ruby laser emitting a brilliant red beam of light.³ The research

paper by Maiman stated that when a ruby crystal of 1 cm dimensions was coated on two parallel faces with silver and irradiated by a high-power flash lamp, the emission spectrum obtained under these conditions showed the presence of negative temperatures and regenerative amplification via stimulated emission was ensued.⁴

A multitude of lasers with different wavelengths and lasing media are now used in surgical practice. Carbon dioxide (CO₂) laser (10,600 nm), Neodymium Yttrium Aluminium Garnet (Nd:YAG) laser (1064 nm), Erbium Yttrium Aluminium Garnet (Er:YAG) laser (2940 nm), Holmium Yttrium Aluminium Garnet (Ho:YAG) laser (2100 nm), Argon laser (514 nm), Potassium Titanyl Phosphate (KTP) laser (532 nm) and semiconductor Diode laser (808-980 nm) are commonly used laser systems.

Each machine delivers laser energy fibro-optically in continuous wave and gated pulsed modes and is used in contact with soft tissue for surgery or out of contact for deeper coagulation. The Diode laser is an excellent soft tissue surgical laser and is indicated for cutting and coagulating oral mucosa.⁶ D'Arcangelo et al observed that in minor oral surgical procedures, diode laser therapy certainly is less invasive and presents some indisputable advantages such as the elimination of bleeding and suturing, as well as minimal postsurgical pain and oedema. The use of low-level lasers started with Mester in 1971. Biomodulation by laser was

based on the principle that irradiation at a specific wavelength can alter cellular behaviour by acting on the mitochondrial chain or membrane calcium channels, promoting an increased cell metabolism and proliferation.⁷

Based on the studies of Ize-Iyamu et al, the laser was seen to expose unerupted teeth, reposition the maxillary frenum, and facilitate aesthetic recontouring of hyperplastic gingivae and remove granulomatous gingivae as a result of orthodontic movement. The use of lasers in soft tissue management maximized the health of the oral tissues.⁸

The efficacy of laser surgery has also been observed in the removal of mucocoeles¹⁰ in overcoming the limitations of surgical methods of treatment in oral submucous fibrosis patients with favourable results¹¹, reducing the incidence and severity of hematoma, swelling and pain during the first week after periosteal releasing incision during guided bone regeneration¹², improving the visibility, reducing the amount of anaesthesia required, offering a promising treatment for the ablation of oral leukoplakia¹³ and minimizing trauma in second stage implant surgery¹⁴ amongst its many other applications.

All in all, both scalpel and diode laser treatment modalities are effective, with the former preceding the latter in time taken for healing and the latter leading in terms of a clean surgical field, and minimal intra and postsurgical complications.

The main objective of this prospective study was to analyze the intraoperative and postoperative effects of both diode laser and scalpel in terms of pain, redness, bleeding swelling, and healing in the oral mucosa.

Materials and Methods

This study compared the properties of intraoral soft tissue diode laser of 970 ± 10 nm (GaAlAs) and cold scalpel (Bard-Parker No. 15) done during the tenure of 2020 to 2023 in the Department of Oral and Maxillofacial Surgery. We selected sample size of **100 patients** and divided equally into two groups, where **Group I** (50 patients; study group): was treated by intraoral soft tissue diode laser and **Group II** (50 patients; control group): were treated by Bard-Parker No. 15 blade and scalpel.

Operative technique: For the laser group, we applied 2W power in continuous mode and removed the tissues in slow, sweeping motion. Charring, if visible, was gently removed by a gauze piece soaked in saline solution. For the scalpel group, incisions were placed with a No. 15 blade in a firm continuous stroke on the mucosa. The rest of the surgical procedure is similar in both groups. After the placement of incision, a full thickness mucoperiosteal flap was reflected or exposed. For third molar extractions, Modified Ward's incision was applied and Coupland, Warwick James elevator and respective third molar forceps were used. For alveoloplasty, after crestal incision, we used bone file and rongeurs and for orthodontic exposures, semicircular incision was placed treatment was rendered up to mucoperiosteal tissue removal and the case was furthered by an orthodontist. After the procedure, copious irrigation was done using 0.9% saline 5% w/v povidone iodine solution and closure was done

3-0 black braided silk sutures. The patients were recalled on the 7th day for the removal of sutures.

Method of Data Collection

Our preliminary research comprised of patient demographic data like name, age, gender and residential address. We took detailed medical, dental, family, and personal history of the patient. A thorough intraoral examination of soft tissue was done to assess the treatment need and informed consent of the patients was taken for their voluntary participation in the study.

To fulfil our inclusion criteria, we selected healthy individuals without any uncontrolled disease/condition. Those who had clinically diagnosed intraoral soft tissue lesions/growth indicative of tissue ablation, incision, excision, or biopsy were sampled for this study.

We excluded patients who, abled or otherwise, did not consent to be a part of the study.

Patients with systemic diseases (uncontrolled diabetes mellitus and hypertension, respiratory or cardiovascular problems, chronic kidney disease, etc, serious eye defects, having porphyria or on medications that cause photosensitivity (eg. Chloroquine, methotrexate etc) and immunocompromised and non-cooperative patients were not involved in this study.

Results

In this clinical prospective study, 100 cases between 14 and 73 years (Male, n=44, Female, n=56) with a mean age of 36.37 years were evaluated. 50 cases were treated with the conventional scalpel and 50 cases with the 970 ± 10 nm diode laser. Out of the scalpel cases, 40 cases were treated for impacted third molar extractions and 10 were treated by minor oral surgery, for cases like alveoloplasty (n=4), surgical extraction (n=3), orthodontic exposure (n=2) and implant placement (n=1). In the laser group, 26 cases were treated for impacted third molar extractions and 24 cases underwent minor oral surgery, for alveoloplasty (n=10), surgical extraction (n=3), orthodontic exposure (n=9) and implant placement (n=2). The swelling percentage preoperatively for scalpel cases is 2.90 ± 1.59 and the laser cases are 1.82 ± 0.75 in the third molar group. The P value is significant between the scalpel and laser cases in the third molar (P=0.001). In the minor oral surgical group, the swelling percentage is 2.23 ± 1.03 for the scalpel and 1.98 ± 0.89 for laser cases respectively. The P value is significant between the scalpel and laser cases in the minor surgical groups (P=0.042). The mean values of the mean bleeding scores in the third molar group are 3.394 in the scalpel and 1.291 in the laser cases respectively. The P value is significant between the scalpel and laser cases (P=0.001) In the minor surgical group, the mean value of scalpel cases is 3.51 and the laser cases is 1.07. The P value is significant between the scalpel and laser cases (P=0.001). On the first postoperative day, the mean values of the mean pain scores in the third molar group are 1.925 in the scalpel and 1.846 in the laser cases respectively. The P value is non-significant between the scalpel and laser cases on the first postoperative day (P=0.623). On the seventh

postoperative day, the mean values of the mean pain scores in the third molar group are 0.6500 in the scalpel and 0.2308 in the laser cases respectively. The P value is significant between the scalpel and laser cases on the first postoperative day ($P=0.001$). On the first postoperative day, the mean values of the mean pain scores in the minor surgical molar group are 1.93 in the scalpel and 1.66 in the laser cases respectively. The P value is non-significant between the scalpel and laser cases on the seventh postoperative day ($P=0.315$). On the seventh postoperative day, the mean values of the mean pain scores in the minor surgical group are 0.70 in the scalpel and 0.20 in the laser cases respectively. The P value is significant between the scalpel and laser cases on the seventh postoperative day ($P=0.001$). The mean values of the mean bleeding scores in the third molar group are 4.52 in the scalpel and 4.5385 in the laser cases respectively. The

P value is non-significant between the scalpel and laser cases ($P=0.754$). In the minor surgical group, the mean value of scalpel cases is 4.20 and the laser cases is 4.25. The P value is non-significant between the scalpel and laser cases ($P=0.915$). In the third molar group, 25% cases showed redness in the scalpel group and 75% did not. In the laser cases, 26.9% showed redness and 73.1% did not. P value was found to be non-significant in the third molar group ($P=1.000$). In the minor oral group, 30% cases showed redness in the scalpel group and 70% did not. In the laser cases, 25% showed redness and 75% did not. P value was found to be non-significant in the minor oral group ($P=1.000$).

Group I:




Figure 1: Intraoperative procedure in Group I Patients. (a) Local anaesthesia administration (b) Laser incision (c) Incised area (d) Surgical exposure done



Figure 2: Closure



Figure 3: Postoperative 3rd day redness



Figure 4 (a)-(b): Postoperative 7th day healing




Figure 5: Intraoperative procedure in Group II patients. (a) Local anaesthesia administration (b) Scalpel incision (c) Mucoperiosteal flap elevation (d) Luxation of the tooth (e) Extracted tooth



Figure 6: Closure




Figure 7: Postoperative 3rd day redness




Figure 8 (a)-(b): Postoperative 7th day healing

Table 1: Intergroup comparison of mean percentage change in swelling scores between the groups

		Pre Op	Day 3	% Swelling	P value	Significance
IIIrd Molar	Scalpel Group	13.33±0.81	13.72±0.83	2.90±1.59	0.001	Significant
	Laser Group	13.35±0.53	13.59±0.52	1.82±0.75		
Minor Surgical Procedure	Scalpel Group	13.25±0.72	13.55±0.76	2.23±1.03	0.042	Significant
	Laser Group	13.10±0.81	13.35±0.83	1.98±0.89		

Table 2: Intergroup comparison of mean bleeding scores between the groups

		Mean	SD	Std Error	P value	Significance
IIIrd Molar	Scalpel Group	3.394	0.887	0.140	0.001	Significant
	Laser Group	1.291	0.507	0.099		
Minor Surgical Procedure	Scalpel Group	3.51	0.893	0.282	0.001	Significant
	Laser Group	1.07	0.478	0.097		

Table 3: Intergroup comparison of mean pain scores between the groups in iiird molars

		Mean	SD	Std Error	P value	Significance
At 1 st day	Scalpel Group	1.925	0.655	0.103	0.623	Non-Significant
	Laser Group	1.846	0.612	0.120		
At 7 th day	Scalpel Group	.6500	.62224	.09838	0.001	Significant
	Laser Group	.2308	.42967	.08427		

Table 4: Intergroup comparison of mean pain scores between the groups in minor surgical procedures

		Mean	SD	Std Error	P value	Significance
At 1 st day	Scalpel Group	1.93	0.737	.23333	0.315	Non-Significant
	Laser Group	1.66	0.564	.11526		
At 7 th day	Scalpel Group	0.70	0.483	0.152	0.001	Significant
	Laser Group	0.20	0.414	0.084		

Table 5: Intergroup comparison of mean healing scores between the groups

		Mean	SD	Std Error	P value	Significance
IIIrd Molar	Scalpel Group	4.52	.93336	.14758	0.754	Non-Significant
	Laser Group	4.5385	.90469	.17742		
Minor Surgical Procedure	Scalpel Group	4.20	1.31656	.41633	0.915	Non-Significant
	Laser Group	4.25	.94409	.19271		

Table 6: Intergroup comparison of redness between the groups

		Absent	Present	P value	Significance
IIIrd Molar	Scalpel Group	30	10	1.000	Non-Significant
		75.0%	25.0%		
	Laser Group	19	7		
		73.1%	26.9%		
Minor Surgical Procedure	Scalpel Group	7	3	1.000	Non-Significant
		70.0%	30.0%		
	Laser Group	18	6		
		75.0%	25.0%		

Statistical analysis

The data for the present study was entered in the Microsoft Excel 2007 and analyzed using the SPSS statistical software 23.0 Version. The descriptive statistics included mean,

standard deviation frequency and percentage. The level of the significance for the present study was fixed at 5%.

The intergroup comparison was done using the independent t tests/Mann Whitney U test depending upon the normality of the data. The categorical variables were compared using the Chi Square test. The Shapiro–Wilk test was used to investigate the distribution of the data and Levene's test to explore the homogeneity of the variables.

Discussion

The knife was the most commonly used tool and the first one to be developed.²⁰ Evidence of knives used in medicine goes back as far as the Mesolithic period (the Middle Stone Age)—around 8000 BC when flint knives were used as scrapers to cut through the skull.²¹

Hippocrates was the first to describe the surgical knife. Which he named *macairion*, from *machaira*, an old Lacedaemonian sword, which had a broad cutting blade on a single edge and a sharp, straight point. Therefore, even in Hippocrates' time, the shape of the scalpel was much the same as it is today. The Romans used the Latin "*scallpellus*," from which the English "*scalpel*" is clearly derived.²⁰

The development of the modern scalpel with a disposable blade, as we know it today, was largely the result of Mr. King Gillette's invention of the safety razor (patented in 1904). In 1910, the eminent Dr. John B. Murphy of Chicago perfected special handles for both single- and double-edged razor blades, assuring the surgeon of ready access to a very sharp knife blade.²¹

Morgan Parker, in 1915, patented a new 2-piece scalpel, with a blade and handle held together by overlapping metal parts which provided rigidity and enabled the exchange of old blades for new ones after each use. Together with Charles Russell Bard, Parker formed the Bard-Parker Company and developed a method of cold sterilization that would not dull the blades, as did the heat.²⁰ The present design of the Rib-Back blade was introduced in 1936.²¹

In 1917, the development of lasers began when Albert Einstein proposed a theory of stimulated emission whose concept of this theory was based on the quantum theory, that described light being composed of small energy packets called quanta. Maiman implemented the use of Ruby laser, using the experience he gained while working with the army corps. A pulsed flashlamp was used to pump a ruby optically and the first LASER working model was developed. On May 16, 1960, the first laser was successfully tested, pioneered by Theodore H. Maiman, based on the study of masers by Schawlow and Townes.⁴

Stern and Sognnaes in 1964 investigated the possible use of a ruby laser to reduce subsurface demineralization.^{15,22-24} However, Adrian et al found that the ruby laser produced significant heat that caused pulpal damage.²⁵

In 1962 Nick Holonyak Jr. laid the foundation for semiconductor laser diode²⁶ responsible for the emission of visible light. Romanos et al¹⁶, in their study, have found that the incision margin using the diode laser is more precise compared to the other systems. Over the past few decades, diode lasers have proved to be versatile and reliable tools

based on their myriad wavelengths and photobiological properties.

Postoperative swelling is a common occurrence after any surgical procedure. In our study, we found that there was marked swelling on the third day, postoperatively, seen more in the scalpel treated group. Pirnat et al⁵ mentioned in their study that when using NIR (near infrared) lasers on soft tissue there is minimal or no bleeding due to a combination of the sealing of small vessels through tissue protein denaturation and stimulation of Factor VII production in clotting. The heat build-up also allows for the sealing of small lymphatic vessels which results in reduced postoperative oedema. This observation has been further confirmed in the studies of Azma E et al²⁷, Elanchezhian et al.², Yasmeen et al.¹⁸

Kashyap et al have mentioned that after a laser excision, the associated lymphatics and blood vessels are sealed which results in insignificant extravasation of fluids and limited inflammatory reaction.¹⁸ However, in the wound following scalpel excision, there is continued extravasation of blood and lymph fluid, which is manifested as a greater degree of swelling and inflammatory reaction. This is the reason for the longer resolution period.⁵

It has also been found that an adjunct treatment modality of providing low-level laser therapy (LLLT) to patients has led to a marked reduction in swelling post-operatively. Aras et al.,²⁹ Kahraman et al.²⁹ and Hawkins et al.³⁰ have all vouched for a positive outcome of postoperative LLLT in their patients where they have found a significant reduction in swelling, pain, and trismus.

In 1988, Wong and Baker³² suggested various scales and charts record the degree of pain according to the patient perspective of young children. Although their study did not involve adult subjects, research made later included adult and geriatric patients and the Faces Pain Scale proved to be a valuable method for noting the magnitude of pain.

In our study, we found that in the intergroup comparison of mean pain scores between the groups on the first postoperative day scalpel cases showed more incidence of pain than lasers. On the seventh postoperative day, this difference was statistically non-significant. Singh et al³⁶ describe that in their study, the pain was significantly less for the diode laser. They say that the reason for less or no post-operative pain is due to the sealing of the sensory nerve endings with the heat of the laser beam. Lasers also bring about the sealing of lymphatics and fibrin clot is formed which protects it from external trauma.

Laser poses a "non-threatening" aesthetic to the patient with its slim and fine tip and therefore patients are more at ease during laser surgery than scalpel operations where patients are intimidated by the sharpness of the instrument and perceive the procedure as a painful one. Reduced postoperative pain has been observed in the laser-treated groups in a multitude of research such as the ones conducted by Ahad et al,⁹ Qafmolla et al,¹⁰ Gabric et al,¹⁹ Patel et al,¹⁷ Bhatsange et al² and Shahnaz et al.¹²

Kashyap et al¹⁸ have stated that although the mechanism of analgesic effects of laser therapy is not well understood, an increased pain threshold through the alteration of neuronal

stimulation and firing pattern, and the inhibition of the medullary reflexes are thought to be involved. In addition, the laser effect is seen in prostaglandin synthesis, resulting in an increased conversion of prostaglandin G2 and prostaglandin H2 into prostaglandin I2 (prostacyclin). Ize-Iyamu et al⁸ found that there was a significant association between treatment type and post-operative pain where laser cases had less pain than scalpel cases. Çayan et al³³ have noted that in their study, postoperative pain was reduced gradually from the 1st postoperative day to the 10th postoperative day and all patients reported either no pain or only mild pain on the 2nd postoperative day. Moreover, no significant difference in postoperative pain was found between the groups. Therefore, it may be considered that the perception of pain in patients is highly subjective and an individual point of view, as compared to a generalized opinion.

Intraoperative blood loss is almost inevitable during surgery. The gravimetric technique is an important tool to collect blood by weighing the swabs to record bleeding during surgery. In our research, we found a statistically significant outcome in the intergroup comparison of mean bleeding scores between the groups ($P=0.001$) where there was more bleeding in the scalpel groups than laser. It is evident that the scalpel wounds bleed and the laser does not because of the coagulation of proteins in the tissues. A similar outcome has been found in a study by Jayesh et al.³⁵ where haemostasis was compared by measuring the change in the weight of gauze pieces, and the swabs in the scalpel group weighed significantly more than that of the laser group. This was further confirmed in a recent study by Gundlapalle et al³⁴.

The heat and light from the laser beam interact with the tissues causing denaturation of protein and coagulation. Jayesh et al³⁵ have mentioned in their study that dental lasers use chromophores like haemoglobin, melanin, water and hydroxyapatite crystals with the diode laser used in the present study (980nm) the chromophores are melanin, haemoglobin (Hb), and oxyhaemoglobin. The oral cavity is highly vascularized and pigmented. The use of a diode laser offers significant advantages in cutting efficiency and getting a bloodless field. Azma E et al.⁹ stated that laser transmits energy to the cells causing warming, welding, coagulation, protein denaturation, drying, vaporization and carbonization. The power settings of the laser determine the size of the coagulum ring formed during the incision. A higher wattage results in a bigger coagulum ring formation during the surgery.

While visually assessing the redness, on the postoperative third day, we found a significant difference between the scalpel and the laser-treated groups ($P=0.001$) with scalpel cases showing more redness than laser ones. Tissue redness has a direct correlation with healing. In similar research done by Singh et al³⁶ where they checked for redness immediately after surgery, they found that the scalpel cases had more redness as compared to the laser cases. This is attributed to the presence and absence of bleeding caused by the scalpel and laser respectively. During our research, we noticed a whitish membrane-like appearance in the mucosa

of the surgical site. This may be due to the denaturation of protein in the tissues exposed by laser.

There is a controversy in the literature regarding the postoperative healing of lasers versus scalpels. Similar to our study, Ize-Iyamu et al⁸ found no significant difference in the healing of soft tissue in their study. This was confirmed by Patel et al¹⁷, who also observed that the wound healing on the 7th day and after 1 month for the scalpel and laser cases showed no statistical significance between groups. Singh et al³⁶ observed that improved wound healing was present in laser cases on both first and seventh days after surgery. They state that because laser-induced wounds, are definite and clean, they heal with secondary intention and there is no scar formation compared to scalpel incisions. This may be due to the minimal degree of wound contraction following laser irradiation which occurs through induction and formation of a smaller number of myofibroblasts and collagen. Jin et al.³⁷ observed that the thermal-induced damage in the laser has a proclivity towards producing more pronounced tissue change. Such changes are associated with increased inflammatory response and an initial delay in the healing response. The quantitative analysis of proinflammatory cytokines, such as tumour necrosis factor (TNF)- α , in wound extracts can contribute to the determination of vitality and wound age, in particular in the very early post-traumatic interval. Transforming growth factor (TGF)- β is composed of a family of multifunctional polypeptide growth factors involved in embryogenesis (cell growth and differentiation), inflammation, regulation of immune response, cell adhesion and migration, extracellular matrix formation, angiogenesis, and wound healing.

The laser-treated area exhibits a fibrinopurulent membrane within 72 hours, following the loss of the superficial necrotic layer of tissue. Epithelial growth is seen to commence at the edges and gradually covers the entire wound. The newly formed epithelium in laser biopsy specimens is seen to be thinner and parakeratotic when compared with the epithelium formed after scalpel excision.¹⁸ Post-laser wounds show an appreciably lesser quantity of myofibroblasts. This results clinically in a lesser degree of wound contraction and scarring and shows improved post-operative function, especially in critical areas of the tongue, the floor of the mouth, the soft palate, and the buccal mucosa.³⁸

In conclusion, we can state that within the confines of our study, we have found that a diode laser is an impeccable option when it comes to reduction in postoperative swelling, pain, tissue redness and bleeding. Its haemostatic property allows increased convenience for the surgeon and renders better patient comfort and satisfaction. It is an effective treatment modality for approaching highly vascularized areas. A thorough knowledge of the power setting of the laser system is imperative to cause minimal lateral tissue damage and promote better healing. The only drawback of laser is that it is expensive and at higher power settings, poses a risk of tissue necrosis. Thus, the diode laser is an effective and versatile surgical tool and can be used at par with the scalpel in intraoral procedures in maxillofacial surgery.

Source of Funding

None.

Conflict of Interest

None.

References

- Barbosa, R. R. (2022). Surgical scalpel handles and blades. *British Journal of Surgery*, 109(10), 916-920.
- Bhatsange A, Meshram EP, Waghmare A, Shiggaon L, Mehrete V, Shende A. A clinical and histological comparison of mucosal incisions produced by scalpel, electrocautery, and diode laser: A pilot study. *J Dent Lasers*. 2016;10:37-42.
- Clayman L, Kuo P, eds: *Lasers in Maxillofacial Surgery and Dentistry*. New York:1997
- Maiman, T. H. (1960). *Stimulated Optical Radiation in Ruby. Essentials of Lasers*, 134-136.
- Samo Pirnat. Versatility of an 810 nm Diode Laser in Dentistry: An Overview. *Journal of Laser and Health Academy* Vol. 2007; No. 4;
- Donald J. Coluzzi, Robert A. Convissar. *Lasers in Clinical Dentistry*. Dent Clin N Am 2004; 48: 751-77
- D'Arcangelo, C., Di Nardo Di Maio, F., Prosperi, G. D., Conte, E., Baldi, M., & Caputi, S. (2007). A preliminary study of healing of diode laser versus scalpel incisions in rat oral tissue: a comparison of clinical, histological, and immunohistochemical results. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*, 103(6), 764-773.
- Ize-Iyamu, I. N., Saheeb, B. D., & Edetanlen, B. E. (2013). Comparing the 810nm diode laser with conventional surgery in orthodontic soft tissue procedures. *Ghana medical journal*, 47(3), 107-111.
- Ahad, Abdul & Tandon, Shruti & Lamba, Arundee & Faraz, Farrukh. (2015). Minimally Invasive Management of Pericoronal Abscess using 810 nm GaAlAs Diode Laser. *International Journal of Laser Dentistry*. 4. 79-82. 10.5005/jp-journals-10022-1061.
- Qafmolla, A., Bardhoshi, M., Gutknecht, N., & Bardhoshi, E. (2014). Evaluation Of Early And Long Term Results Of The Treatment Of Mucocele Of The Lip Using 980 Nm Diode Laser. *European Scientific Journal, ESJ*, 10(6).
- Shah, A., Shah, R., Vyas, N., & Shah, N. (2017). Clinical trial to compare conventional incision technique and diode laser on the treatment of oral submucous fibrosis.
- Shahnaz, A., Jamali, R., Mohammadi, F., Khorsand, A., Moslemi, N., & Fekrazad, R. (2018). A preliminary randomized clinical trial comparing diode laser and scalpel periosteal incision during implant surgery: impact on postoperative morbidity and implant survival. *Lasers in medical science*, 33(1), 19-25.
- Yasmeen SA, Reddy BHS, Ramesh MV, Birajdar SS, Yarram A, Kumar NN. A Comparative Evaluation of Diode Laser Ablation versus Scalpel Excision for Management of Oral Leukoplakia. *Journal of Indian Academy of Oral Medicine & Radiology*. 2019;31(2):94-9.
- Kaur M, Sharma YP, Singh P, Sharma S, Wahi A (2018) Comparative evaluation of efficacy and soft tissue wound healing using diode laser (810 nm) versus conventional scalpel technique for second-stage implant surgery. *J Indian Soc Periodontol* 22:228-234
- Wigdor H, Walsh J, Visuri S, Fried D, Waldvogel J. Lasers in dentistry. *Lasers Surg Med* 1995;16:103-33.
- Romanos G. Clinical applications of the Nd:YAG laser in oral soft tissue surgery and periodontology. *J Clin Laser Med Surg* 1994;12:103-8.
- Patel R M, Varma S, Suragimath G, Abbayya K, Zope S A, Kale V. Comparison of labial frenectomy procedure with conventional surgical technique and diode laser . *J Dent Lasers* 2015;9:94-9
- Kashyap, S., Kumar, V., Prakash, P., Haidry, N., Ranganath, N., & D'souza, D. (2018). Comparison of Scalpel versus Soft Tissue Diode Laser for Biopsy of Oral Lesions.
- Gabrić, Dragana & Sušić, Mato & Katanec, Davor & Filipović-Zore, Irina & Vucicevic Boras, Vanja & Brailo, Vlaho & Vrdoljak, Danko. (2015). Comparison between Diode Laser and Conventional Technique for Soft Tissue Oral Surgery: A Pilot Study. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 6. 1913.
- Ochsner J. (2009). Surgical knife. *Texas Heart Institute journal*, 36(5), 441-443.
- Ochsner J. The surgical knife. *Bull Am Coll Surg* 1999;84(2): 27-37.
- Stern RH, Sognnaes RF. Laser beam effect on dental hard tissues. *J Dent Res* 1964; 43 (Suppl.):873 (Abs 307).
- Stern R, Renger HL, Howell FV. Laser effects on vital dental pulps. *Br Dent J* 1969; 26-28.
- Stern RH, Vahl J, Sognnaes RF. Laser enamel: Ultrastructural observations of pulsed carbon dioxide laser effects. *J Dent Res* 1972; 51:455-460.
- Adrian JC, Bernier JL, Sprague WG. Laser and the dental pulp. *JADA* 1971; 83:113-117.
- Holonyak Jr, N., & Bevacqua, S. F. (1962). Coherent (visible) light emission from Ga (AsI- x P x) junctions. *Applied Physics Letters*, 1(4), 82-83.
- Azma, E., & Safavi, N. (2013). Diode laser application in soft tissue oral surgery. *Journal of lasers in medical sciences*, 4(4), 206-211.
- Elanchezhian, S., Renukadevi, R., & Vennila, K. (2013). Comparison of diode laser-assisted surgery and conventional surgery in the management of hereditary ankyloglossia in siblings: a case report with scientific review. *Lasers in medical science*, 28(1), 7-12.
- Aras, M. H., & Güngörmüş, M. (2009). The effect of low-level laser therapy on trismus and facial swelling following surgical extraction of a lower third molar. *Photomedicine and laser surgery*, 27(1), 21-24.
- Kahraman, S. A., Cetiner, S., & Strauss, R. A. (2017). The effects of transcutaneous and intraoral low-level laser therapy after extraction of lower third molars: a randomized single blind, placebo controlled dual-center study. *Photomedicine and laser surgery*, 35(8), 401-407.
- Hawkins, D., Houreld, N., & Abrahamse, H. (2005). Low level laser therapy (LLLT) as an effective therapeutic modality for delayed wound healing. *Annals of the New York Academy of Sciences*, 1056(1), 486-493.
- Wong, D. L., & Baker, C. M. (1988). Pain in children: comparison of assessment scales. *Pediatr Nurs*, 14(1), 9-17.
- Çayan, T., Hasanoğlu Erbaşar, G. N., Akca, G., & Kahraman, S. (2019). Comparative Evaluation of Diode Laser and Scalpel Surgery in the Treatment of Inflammatory Fibrous Hyperplasia: A Split-Mouth Study. *Photobiomodulation, photomedicine, and laser surgery*, 37(2), 91-98.
- Gundlapalle, P., Nagappan, N., Ramesh, P., Ziauddin, S., Karthick, B. P., Paleti, V. S., & Kumar, B. V. (2022). Comparison of Oral Mucosal Biopsies Done Using Scalpel and Diode Lasers: A Vivo Study. *Journal of Pharmacy & Bioallied Sciences*, 14.
- Jayesh, Sibgatullah, Aklanta Kumar Gogoi, Vidhya Ningrajappa, Ningombam Gitson Meitei, et al. Surgical Scalpel vs Laser: A Comparative Clinical Study in Intra Oral Incisional Biopsy. *Mod Res Dent*. 6(1). MRD. 000630. 2021.
- Singh, P., & Nath, S. (2019). Diode laser versus conventional technique for frenectomy-A randomised controlled clinical trial. *Int J Appl Dent Sci*, 5(2), 04-07.

37. Jin, J. Y., Lee, S. H., & Yoon, H. J. (2010). A comparative study of wound healing following incision with a scalpel, diode laser or Er, Cr: YSGG laser in guinea pig oral mucosa: A histological and immunohistochemical analysis. *Acta Odontologica Scandinavica*, 68(4), 232-238.
38. Wlodawsky, R. N., & Strauss, R. A. (2004). Intraoral laser surgery. *Oral and Maxillofacial Surgery Clinics*, 16(2), 149-163.

<p>How to cite: Gupta A, Mittal A, Smiti Klaire SJ, Sharma S, Roy B, Tousif Alam MD, et al. An analogical study on antihypertensive effect of moxonidine versus clonidine in patients with renal failure. <i>Journal of Orofacial and Health Sciences (JOHS)</i>. 2024;11(1):28-37</p>
