

## Review Article

### Aripiprazole and Clozapine: A Review of Spectroscopic and Chromatographic Method

Noopur K. Gandhi, Darshil B. Shah\*, Dilip G. Maheshwari

Received on: 10-12-2016  
Accepted on: 25-12-2016  
Published on: 15-02-2017

#### Corresponding Author:

\* Darshil B. Shah,  
Department of Quality Assurance,  
L.J. Institute of Pharmacy,  
Ahmedabad. Gujarat  
India.



#### ABSTRACT

Aripiprazole and Clozapine are classified as an Atypical Antipsychotics. Aripiprazole primarily used in the treatment of schizophrenia and bipolar disorder. Clozapine works by changing the actions of chemicals in the brain. It is used to treat severe schizophrenia, or to reduce the risk of suicidal behavior in people with schizophrenia or similar disorders. It is also used in Parkinson's disease. The clinical and pharmaceutical analysis of these drugs requires effective analytical procedures for quality control and pharmacodynamic and pharmacokinetic studies as well as stability study. There are many analytical methods reported so far in the literature for the determination of Aripiprazole and Clozapine in Biological samples and pharmaceutical formulations. This article comprises reviews of analytical methods like Spectrophotometric methods, chromatographic methods.

\*Email Id- darshilshah89@yahoo.com

**Key-words:** Aripiprazole, Clozapine, Spectrophotometry, HPLC

#### Cite this article as:

Noopur K. Gandhi, Darshil B. Shah, Dilip G. Maheshwari, Prescription Pattern of Drugs In Pregnancy Induced Hypertension In A Tertiary Care Hospital, Asian Journal of Pharmaceutical Technology & Innovation, 05 (22); 27-39, 2017. [www.asianpharmtech.com](http://www.asianpharmtech.com)

**INTRODUCTION:****ARIPIPRAZOLE:**

Aripiprazole is an Atypical Antipsychotic drug. It is primarily used in the treatment of schizophrenia and bipolar disorder. Although it is used as an add-on treatment in major depressive disorder, tic disorders, and irritability associated with autism. Aripiprazole have partial agonistic activity at D2 receptor, also have partial agonist activity at 5-HT1A receptor, and have antagonist activity at 5-HT2A receptor.




Figure 1: Structure of Aripiprazole

**Table 1: Structural identification of Aripiprazole**

| Sr. no. | Class               | Identification   |
|---------|---------------------|--|
| 1       | Kingdom             | Organic compound   |
| 2       | Super Class         | Organohetrocyclic compound   |
| 3       | Class               | Diazinanes   |
| 4       | Sub class           | Piperazines  |
| 5       | Direct parent       | Phenylpiperazines  |
| 6       | Alternative parent  | N-arylpiperazines, Hydroquinolones, Substituted anilines, Dichlorobenzenes, Dialkylarylamines, N-alkylpiperazines, Alkyl aryl ethers, Aryl chlorides, Trialkyl amines, Secondary carboxylic acid amides, Azacyclic compounds, Organochlorides, Hydrocarbon derivatives, Carbonyl compounds |
| 7       | Molecular framework | Aromatic heterocyclic compound   |

**Table 2: Drug Profile**

| Sr. No. | Parameters       | Description   |
|---------|------------------|---|
| 1       | Category         | Antipsychotic agent   |
| 2       | Chemical Formula | C <sub>23</sub> H <sub>27</sub> Cl <sub>2</sub> N <sub>3</sub> O <sub>2</sub>       |
| 3       | IUPAC Name       | 7-{4-[4-(2,3-dichlorophenyl)piperazin-1-yl]butoxy}-1,2,3,4-tetrahydroquinolin-2-one |
| 4       | Molecular Weight | 448.38538 g/mol   |
| 5       | Characteristic   | Colorless, flake crystals from ethanol  |
| 6       | Solubility       | Soluble in Methanol, Ethanol and slightly soluble in water                          |

**Table 3: OFFICIAL METHODS FOR ESTIMATION OF ARIPIPRAZOLE**

Aripiprazole is official in Indian pharmacopoeia (IP 2014) and United State Pharmacopoeia (USP 2012).

| Sr. No. | Drug                   | Method                | Description  | Ref. No. |
|---------|------------------------|-----------------------|--|----------|
| 1       | Aripiprazole (IP 2014) | Liquid Chromatography | <b>Detection Wavelength:</b> 220 nm<br><b>Mobile Phase:</b> Potassium Dihydrogen orthophosphate: Triethylamine (pH 3): methanol: Acetonitrile (50:2:25:25 v/v/v/v)<br><b>Stationary Phase:</b> Stainless Steel Column 25 cm × 4.6 mm, packed with Octadecylsilane bonded to porous silica (5µm).<br><b>Flow Rate:</b> 1 ml/min | 7        |

|   |                                |                              |  |   |
|---|--------------------------------|------------------------------|--|---|
|   |                                |                              | <b>Injection Volume:</b> 20 µl   |   |
| 2 | Aripiprazole Tablet (USP 2012) | <b>Liquid Chromatography</b> | <b>Detection Wavelength:</b> 254 nm<br><b>Solution A:</b> 2.8 g/L of anhydrous sodium sulfate in water<br><b>Mobile Phase:</b> Acetonitrile: Methanol: Sloution A: Glacial Acetic Acid (33:11:56:1)<br><b>Stationary Phase:</b> Octylsilane Bonded to totally porous microsilica particles, 3 to 10 µm in diameter, 4.6 mm×15 cm<br><b>Flow Rate:</b> 1 ml/min<br><b>Injection Volume:</b> 20 µl | 8 |

**TABLE 4: REPORTED METHOD OF ARIPIPRAZOLE**

| Sr No. | Drug  | Method   | Description  | Ref. No. |
|--------|---|--|--|----------|
| 1      | Aripiprazole in Bulk and Pharmaceutical formulation | <b>UV Spectroscopy</b>                             | <b>Detection Wavelength:</b> 219 nm<br><b>Solvent:</b> Methanol<br><b>Linearity Range:</b> 2-10 µg/ml<br><b>Correlation coefficient:</b> 0.9998  | 9        |
| 2      | Aripiprazole in tablet dosage forms                 | <b>Visible Spectrophotometric Method</b>           | <b>Detection Wavelength:</b> 414 nm<br><b>Solvent:</b> HCl<br><b>Linearity Range:</b> 10-60 µg/ml<br><b>Coloring Agent:</b> Bromocresol green dye<br><b>Correlation coefficient:</b> 0.9968  | 10       |
| 3      | Aripiprazole in tablet dosage forms                 | <b>UV Spectrometry</b>                             | <b>Detection Wavelength:</b> 218 nm<br><b>Solvent:</b> 0.05 M Phosphoric Acid: Acetonitrile (40:60 v/v)<br><b>Linearity Range:</b> 2.5-20 µg/ml<br><b>Correlation coefficient:</b> 0.9992<br><b>LOD:</b> 0.01 µg/ml<br><b>LOQ:</b> 0.1 µg/ml   | 11       |
| 4      | Aripiprazole in tablet dosage form                  | <b>UV Spectrophotometer</b><br>A) Ion pair complex | <b>Detection wavelength:</b> 514 nm<br><b>Solvent:</b> dilute sulphuric acid<br><b>Linearity range:</b> 4-26 µg/ml<br><b>Coloring Agent:</b> Erichrome Black T (EBT)<br><b>Correlation coefficient:</b> 0.9999<br><b>LOD:</b> 0.9523 µg/ml<br><b>LOQ:</b> 3.1714 µg/ml   | 12       |
| 5      | Aripiprazole in Tablet Dosage form                  | <b>RP-HPLC Method</b>                              | <b>Detection wavelength:</b> 254 nm<br><b>Mobile Phase:</b> Acetonitrile: Sodium acetate buffer (55:45 v/v)<br><b>Stationary Phase:</b> Phenomenex Luna C18 column (250 mm length, 4.6 mm internal diameter, 5 mm particle size)<br><b>Flow Rate:</b> 1 ml/min<br><b>Linearity Range:</b> 2-12 µg/ml<br><b>Retention Time:</b> 6.84 min<br><b>Regression Coefficient:</b> 0.9995 | 13       |
| 6      | Aripiprazole in                                     | <b>UV</b>  | <b>Detection Wavelength:</b> 256 nm  | 14       |

|    |   |   |  |    |
|----|---|---|--|----|
|    | Pharmaceutical Dosage Form                              | <b>Spectrophotometry</b>                                  | <b>Solvent:</b> 95% ethanol<br><b>Linearity range:</b> 5-30 µg/ml<br><b>Correlational Coefficient:</b> 0.9995  |    |
| 7  | Aripiprazole in Tablet Dosage form                      | <b>Stability indicating liquid chromatographic column</b> | <b>Detection wavelength:</b> 254 nm<br><b>Mobile Phase:</b> Ammonium acetate buffer: Acetonitrile: Methanol (50:40:10 v/v)<br><b>Stationary Phase:</b> Zorbax 150mm×4.6mm, C18 column with 5µm particles<br><b>Flow Rate:</b> 1.5 ml/min<br><b>Linearity Range:</b> 100-800 µg/ml<br><b>Regression Coefficient:</b> 1  | 15 |
| 8  | Aripiprazole in Bulk and in pharmaceutical formulation  | <b>RP-HPLC Method</b>                                     | <b>Detection wavelength:</b> 255 nm<br><b>Mobile Phase:</b> buffer: acetonitrile: THF (30:60:10)<br><b>Stationary Phase:</b> waters spherisorb 5µ ODS (24.6mm×250mm) column<br><b>Flow Rate:</b> 1.5 ml/min<br><b>Linearity range:</b> 1-100 µg/ml<br><b>Correlation coefficient:</b> 0.999  | 16 |
| 9  | Aripiprazole in Bulk and in pharmaceutical formulation  | <b>RP-HPLC Method</b>                                     | <b>Detection wavelength:</b> 283 nm<br><b>Mobile Phase:</b> 0.02 M Sodium Dihydrogen Orthophosphate: Methanol (30:70 v/v)<br><b>Stationary phase:</b> INTERSIL C18 column (250×4.6 mm I.D., particle size 5 µm)<br><b>Flow Rate:</b> 0.8 ml/min<br><b>Linearity range:</b> 48-145 µg/ml<br><b>Correlation Coefficient:</b> 0.99988<br><b>LOD:</b> 0.22 µg/ml<br><b>LOQ:</b> 0.66 µg/ml | 17 |
| 10 | Aripiprazole in Human Serum                             | <b>LCMS/MS</b>  | <b>Stationary Phase:</b> API 3200 Triple quadrapole mass spectrometer using Chromolith, RP-18e column<br><b>Mobile Phase:</b> 5 mM Ammonium Acetate in Water: Acetonitrile (25:75 %v/v)<br><b>Linearity range:</b> 1-200 ng/ml   | 18 |
| 11 | Aripiprazole in Bulk and its Pharmaceutical formulation | <b>RP-HPLC Method</b>                                     | <b>Detection Wavelength:</b> 254 nm<br><b>Stationary Phase:</b> HSF5 C18 (4.6 mm × 250mm, 5µm) column<br><b>Mobile Phase:</b> Methanol: Acetonitrile: sodium sulphate buffer (25:50:50)<br><b>Linearity range:</b> 20-200 µg/ml<br><b>Flow rate:</b> 1.2 ml/min<br><b>Internal standard:</b> Caffiene  | 19 |
| 12 | Aripiprazole in   | <b>UV Spectroscopy</b>                                    | <b>A) Charge-transfer:</b>   | 20 |

|    |   |   |   |    |
|----|---|---|---|----|
|    | Tablet dosage Form                          | <p><b>a) Charge Transfer</b><br/> <b>b) Ion-Pair complexation</b></p> | <p><b>π acceptor:</b> 2,3-dichloro-5,6-dicyano-p-benzoquinone (DDQ)<br/> <b>σ acceptor:</b> iodine (I<sub>2</sub>)<br/> <b>Solvent &amp; Detection Wavelength:</b><br/>                     457 nm in acetonitrile<br/>                     364 nm in 1,2-dichloroethane<br/> <b>Correlation coefficient:</b><br/>                     DDQ: 0.9997<br/>                     I<sub>2</sub>: 0.9998<br/> <b>Linearity range:</b><br/>                     DDQ: 10-120 µg/ml<br/>                     I<sub>2</sub>: 2-28 µg/ml<br/> <b>LOD:</b><br/>                     DDQ: 2.44<br/>                     I<sub>2</sub>: 0.39<br/> <b>LOQ:</b><br/>                     DDQ: 8.12<br/>                     I<sub>2</sub>: 1.31<br/> <b>B) Ion-pair complexation:</b><br/> <b>Coloring Agent:</b> Bromocresol green (BCG), Bromocresol purple (BCP)<br/> <b>Solvent:</b> 1,2-dichloroethane<br/> <b>Detection Wavelength:</b><br/>                     BCG: 413 µg/ml<br/>                     BCP: 400 µg/ml<br/> <b>Correlation coefficient:</b><br/>                     BCG: 0.9997 µg/ml<br/>                     BCP: 0.9999 µg/ml<br/> <b>Linearity range:</b><br/>                     BCG: 2-24 µg/ml<br/>                     BCP: 2-20 µg/ml<br/> <b>LOD:</b><br/>                     BCG: 0.50 µg/ml<br/>                     BCP: 0.30 µg/ml<br/> <b>LOQ:</b><br/>                     BCG: 1.68 µg/ml<br/>                     BCP: 1 µg/ml</p> |    |
| 13 | Aripiprazole in pharmaceutical formulations | <b>UV-VIS Spectrometry</b>  | <p><b>Detection Wavelength:</b> 480 nm<br/> <b>Solvent:</b> Chloroform<br/> <b>Linearity range:</b> 2-12 µg/ml<br/> <b>Coloring Agent:</b> 3-Methyl-2-benzothiazolinone-hydrazone (MBTH), Ferric chloride<br/> <b>Correlation coefficient:</b> 0.9999</p>   | 21 |
| 14 | Aripiprazole in human plasma                | <b>SPE-UPLC-MS/MS</b>   | <p><b>Mobile phase:</b> Methanol: 10mM ammonium formate (85:15 v/v)<br/> <b>Stationary phase:</b> UPLC BEH C18 (50 mm×2.1 mm, 1.7 µm) column<br/> <b>Internal Standard:</b> aripiprazole-d8<br/> <b>Linearity range:</b> 0.05-80 ng/ml</p>  | 22 |

|    |  |   |   |    |
|----|--|---|---|----|
| 15 | Aripiprazole in pharmaceuticals and human plasma | <b>Chemiluminescence Method</b>   | <b>Chemiluminance</b> of tris(1,10-phenanthroline)-ruthenium(II), Ru(phen) <sub>3</sub> <sup>2+</sup><br><b>Linearity range:</b> 1.8-18 ng/ml<br><b>Correlation coefficient:</b> 0.9951<br><b>LOD:</b> 0.9 ng/ml  | 23 |
| 16 | Aripiprazole in bulk                             | <b>Stability indicating Spectrophotometric and TLC-Densitometric Method</b> | <b>A) Spectroscopic Method</b><br><b>Solvent:</b> Acetonitrile<br><b>Detection Wavelength:</b> 217.2 nm, 229 nm<br><b>Linearity range:</b> 1-6 µg/ml<br><b>B) TLC-Densitometric Method:</b><br><b>Standard solution:</b> 0.1 mg/ml<br><b>Detection Wavelength:</b> 255 nm<br><b>Mobile Phase:</b> Ethyl acetate: Methanol (11:4 v/v)<br><b>Stationary Phase:</b> TLC-Plate, 4 mm band length, 3 mm×0.45 mm slit dimension<br><b>Scanning Speed:</b> 20 mm/s | 24 |
| 17 | Aripiprazole in Human Plasma                     | <b>LC-ESI-MS Method</b>   | <b>Internal Standard:</b> Zolpidem tartrate<br><b>Stationary phase:</b> Grace Smart RP 184.6×100 mm, 3 µ column<br><b>Mobile phase:</b> Methanol: Ammonium acetate buffer (95:5 v/v)<br><b>Linearity range:</b> 0.2-60 ng/ml<br><b>Flow Rate:</b> 0.6 ml/min<br><b>M/Z:</b><br>Aripiprazole: 448.03→285.14<br>Zolpidem tartrate: 308.13→235.25  | 25 |
| 18 | Aripiprazole and dehydroaripiprazole             | <b>Capillary-Electrophoresis</b>  | <b>Detection Wavelength:</b> 214 nm<br><b>Solvent:</b> DMSO, Methanol<br><b>Capillary:</b> 60 cm length, 75 µm internal diameter<br><b>Running Buffer:</b> 2.5-20% MeOH-phosphate buffer, 1-10% DMSO-phosphate buffer, MeOH-DMSO-phosphate buffer<br><b>Linearity range:</b> 2-10 ng/ml   | 26 |
| 19 | Aripiprazole and dehydroaripiprazole             | <b>HPLC-UV Method</b>   | <b>Detection Wavelength:</b> 254 nm<br><b>Mobile phase:</b> Chloroform: n-heptane (3:7 v/v)<br><b>Stationary phase:</b> C18 STRODS-2, 5 µm<br><b>Internal Standard:</b> 7-[5-[4-(3-chloro-2-methylphenyl)-1-piperazinyl]pentyloxy]-3,4-dihydro-   | 27 |

|    |   |                           |  |    |
|----|---|---------------------------|--|----|
|    |   |                           | 2(1H)-quinolinone (OPC-14558)<br><b>Linearity range:</b><br>Aripiprazole: 2-600 ng/ml<br>Dehydroaripiprazole: 2-160 ng/ml  |    |
| 20 | Aripiprazole and escitalopram Oxalate                       | UV Spectrometry           | <b>Detection Wavelength:</b><br>Aripiprazole- 255 nm<br>Escitalopram Oxalate- 238 nm<br><b>Solvent:</b> Methanol<br><b>Linearity Range:</b><br>Aripiprazole: 5-25 µg/ml<br>Escitalopram Oxalate: 15-75 µg/ml<br><b>Correlation coefficient:</b><br>Aripiprazole: 0.999<br>Escitalopram:0.999<br><b>LOD:</b><br>Aripiprazole: 0.129 µg/ml<br>Escitalopram Oxalate: 0.223 µg/ml<br><b>LOQ:</b><br>Aripiprazole: 0.392 µg/ml<br>Escitalopram Oxalate: 0.677 µg/ml | 28 |
| 21 | Aripiprazole and divalproex sodium in combined dosage form. | RP-HPLC Method            | <b>Detection wavelength:</b> 210 nm<br><b>Mobile Phase:</b> Acetonitrile:0.32% KH <sub>2</sub> PO <sub>4</sub> (60:40 v/v)<br><b>Stationary Phase:</b> C18 column (250×4.6 mm) in isocratic mode<br><b>Flow Rate:</b> 1 ml/min   | 29 |
| 22 | Aripiprazole (ARP) and Tapentadol (TAP)                     | UV-VIS. Spectrophotometry | <b>Detection Wavelength:</b> 543 nm<br><b>Solvent:</b> Chloroform<br><b>Coloring Agent:</b> Chloranilic acid<br><b>Linearity range:</b><br>ARP: 80-400 µg/ml<br>TAP: 200-1000 µg/ml<br><b>Correlation Coefficient:</b> 0.9999<br><b>LOD:</b><br>ARP: 5.17<br>TAP: 82.5<br><b>LOQ:</b><br>ARP: 15.66<br>TAP: 250  | 30 |

**CLOZAPINE:**

Clozapine is an Atypical Antipsychotic drug. It works by changing the actions of chemicals in the brain. It is used to treat severe schizophrenia, or to reduce the risk of suicidal behavior in people with schizophrenia or similar disorders. It is also used in Parkinson’s disease.




Figure 1: Structure of Clozapine

**Table 5: Structural identification of Clozapine**

| Sr. no. | Class               | Identification  |
|---------|---------------------|---|
| 1       | Kingdom             | Organic compound  |
| 2       | Super Class         | Organoheterocyclic compound   |
| 3       | Class               | Benzodiazepines   |
| 4       | Sub class           | Diabenzodiazepines  |
| 5       | Direct parent       | Diabenzodiazepines  |
| 6       | Alternative parent  | 1,4-benzodiazepines, N-methylpiperazines, Chlorobenzines, Imidolactams, Aryl chlorides, Trialkylamines, Secondary amines, Propargyl-type 1,3-dipolar organic compounds, carboxamidines, Azacyclic compounds, Organochlorides, Hydrocarbon derivatives |
| 7       | Molecular framework | Aromatic heterocyclic compound  |

**Table 6: Drug Profile**

| Sr. No. | Parameters       | Description  |
|---------|------------------|--|
| 1       | Category         | Antipsychotic agent  |
| 2       | Chemical Formula | C <sub>18</sub> H <sub>19</sub> ClN <sub>4</sub>                   |
| 3       | IUPAC Name       | 8-chloro-11-(4-methylpiperazin-1-yl)-5H-dibenzo[b,e][1,4]diazepine |
| 4       | Molecular Weight | 326.823 g/mol  |
| 5       | Characteristic   | Yellow Crystal   |
| 6       | Solubility       | 0.1889 mg/ml in water  |

**TABLE 7: OFFICIAL METHOD ESTIMATION OF CLOZAPINE:**

Clozapine is Official in Indian pharmacopoeia (IP 2014), United State Pharmacopoeia (USP NF 2004), European Pharmacopoeia (EP 2014).

**Table 2.3: Official method estimation of clozapine**

| Sr. No. | Drug                       | Method                       | Description   | Ref. No. |
|---------|----------------------------|------------------------------|---|----------|
| 1       | Clozapine (IP 2014)        | <b>Titrimetric Method</b>    | 0.115 gm, dissolve in 70 ml of glacial acetic acid and titrate with 0.1 M Perchloric acid, determining the end-point potentiometrically. Carry Out a blank titration.<br>1 ml of 0.1 M Perchloric acid is equivalent to 0.01634 g of C <sub>18</sub> H <sub>19</sub> ClN <sub>4</sub> . | 7        |
| 2       | Clozapine Tablet (IP 2014) | <b>Liquid Chromatography</b> | <b>Detection Wavelength:</b> 257 nm<br><b>Mobile Phase:</b> Methanol: Water: Triethylamine ( 800:200:0.75 v/v/v)<br><b>Stationary Phase:</b> Stainless steel column 25 cm × 4.0 mm, packed with Octylsilane bonded to porous silica (5 µm)<br><b>Flow Rate:</b> 1 ml/min                | 7        |
| 3       | Clozapine (USP NF 2004)    | <b>Titrimetric Method</b>    | 115 mg, dissolve in 70 ml of glacial acetic acid and titrate with 0.1 M Perchloric acid, determining the end-point potentiometrically. Carry Out a blank titration.   | 31       |



|   |                     |                           |   |    |
|---|---------------------|---------------------------|---|----|
|   |                     |                           | 1 ml of 0.1 M Perchloric acid is equivalent to 16.34 mg of C <sub>18</sub> H <sub>19</sub> ClN <sub>4</sub> .   |    |
| 4 | Clozapine (EP 2014) | <b>Titrimetric Method</b> | 0.100 g in 50 ml of anhydrous acetic acid. Titrate with 0.1 M Perchloric acid, determining end-point potentiometrically.<br>1 ml of 0.1 M Perchloric acid is equivalent to 16.34 mg of C <sub>18</sub> H <sub>19</sub> ClN <sub>4</sub> . | 32 |

**TABLE 8: REPORTED METHOD OF CLOZAPINE**

| Sr. No. | Drug   | Method  | Description   | Ref. No. |
|---------|--|---|---|----------|
| 1       | Clozapine in Bulk & tablet dosage forms      | <b>RP-HPLC Method</b>   | <b>Detection Wavelength:</b> 290 nm<br><b>Mobile Phase:</b> Acetonitrile: Phosphate buffer (70:30% v/v)<br><b>Stationary Phase:</b> C <sub>18</sub> column (250×4 internal diameter in isocratic mode)<br><b>Linearity Range:</b> 1-5 µg/ml<br><b>Retention Time:</b> 3.06 min<br><b>Flow Rate:</b> 1 ml/min<br><b>% Recovery:</b> 97.85 to 101.45%<br><b>LOD:</b> 0.1 µg/ml<br><b>LOQ:</b> 0.5 µg/ml | 33       |
| 2       | Clozapine in Human plasma                    | <b>HPLC-UV Method</b>   | <b>Detection Wavelength:</b> 250 nm<br><b>Mobile Phase:</b> acetonitrile: Methanol: 0.5% triethylamine (40:10:50)<br><b>Stationary Phase:</b> ODS hyoersil (5µm) cartridge column (125×4mm I.D.)<br><b>Linearity Range:</b> Drug-free human plasma concentration- 25-2000 ng/ml<br>Drug- human plasma concentration- 75-1500 ng/ml<br><b>Flow Rate:</b> 1 ml/min<br><b>LOQ:</b> 25 mg/ml              | 34       |
| 3       | Clozapine in Human plasma                    | <b>HPLC-UV Method</b>   | <b>Detection Wavelength:</b> 250 nm<br><b>Internal Standard:</b> amino-acetophenone<br><b>Mobile Phase:</b> Acetonitrile (30%)- 29 mM Phosphate buffer<br><b>Stationary Phase:</b> CTO-10ASC18 column (25×4.6×5 µ)<br><b>Linearity Range:</b> 20-2000 ng/ml<br><b>Flow Rate:</b> 1 ml/min<br><b>Liquid-Liquid extraction:</b> Diethyl ether   | 35       |
| 4       | Clozapine in Tablet and in biological fluids | <b>UV-VIS Spectrophotometer Method using Ion-Pair Complex</b> | <b>Detection Wavelength:</b> 514 nm<br><b>Linearity Range:</b> 2-18 µg/ml<br><b>Coloring Agent:</b> Erichrome black T (EBT)<br><b>Solvent:</b><br>For Tablet:Sulphuric Acid<br>For Human Serum and Urine: Acetonitrile  | 36       |
| 5       | Clozapine in tablets and in Urine            | <b>Spectrophotometric and Fluorimetric Method</b>             | <b>A) Spectrophotometric Method:</b><br><b>σ-acceptor:</b> Iodine<br><b>π-acceptor:</b> 7,7,8,8-  | 37       |

|   |   |  |   |    |
|---|---|--|---|----|
|   |   |  | tetracyanoquinondimethane (TCNQ), 2,3-dichloro-5,6-dicyano-1,4-benzo-quinone (DDQ), tetracyanoethane (TCNE), p-chloranilic acid (pCA)<br><b>Detection Wavelength:</b><br>Iodine: 365 nm<br>TCNQ: 843 nm<br>DDQ: 460 nm<br>TCNE: 414 nm<br>pCA: 520 nm<br><b>Linearity Range:</b> 4-200 µg/ml<br><b>LOD:</b><br>Iodine: 1.12 µg/ml<br>TCNQ: 176 µg/ml<br>DDQ: 2.22 µg/ml<br>TCNE: 0.95 µg/ml<br>pCA: 13.26 µg/ml<br><b>B) Fluorimetric Method:</b><br><b>Detection Wavelength:</b><br>$\lambda_{\text{excitation}}$ -260 nm<br>$\lambda_{\text{emission}}$ - 355 nm<br><b>Solvent:</b> 1 M Suphuric Acid<br><b>Linearity Range:</b> 24-250 ng/ml<br><b>LOD:</b> 6.69 ng/ml |    |
| 6 | Clozapine, Norclozapine in various biological fluids                | <b>HPLC-UV Method</b>  | <b>Detection Wavelength:</b> 254 nm<br><b>Internal Standard:</b> Loxapine<br><b>Mobile Phase:</b> 10 mM Ammonium acetate: Acetonitrile: Methanol (5:3:2 v/v/v)<br><b>Stationary Phase:</b> C6 Phenyl column (3 µm, 2×150 mm)  | 38 |
| 7 | Clozapine, Desmethylclozapine and Clozapine N-oxide in Human Plasma | <b>HPLC-UV Method</b>  | <b>Detection Wavelength:</b> 254 nm<br><b>Internal Standard:</b> Triprolidine<br><b>Mobile Phase:</b> Acetonitrile: 0.06 M Phosphate buffer (48:52 v/v)<br><b>Stationary Phase:</b> 250×4.60 mm I.D. analytical column packed with 5 µ C6 silica particles<br><b>Flow rate:</b> 1 ml/min<br><b>Liquid-Liquid Extraction:</b> n-hexane: isoamyl alcohol (75:25 v/v)<br>Organic phase back extracted with 150 microl of 0.1 M dibasic phosphate   | 39 |
| 8 | Lamotrigine and Clozapine   | <b>UV Spectroscopic method</b><br><b>Q Absorbance Ratio Method</b> | <b>Detection Wavelength:</b><br>Lamotrigine: 307 nm<br>Clozapine: 360 nm<br><b>Solvent:</b> Methanol<br><b>Linearity Range:</b><br>Lamotrigine: 1-5 µg/ml<br>Clozapine: 6-30 µg/ml<br><b>Correlation coefficient:</b>   | 40 |

|   |  |                               |   |    |
|---|--|-------------------------------|---|----|
|   |  |                               | Lamotrigine: 0.9992<br>Clozapine: 0.993<br><b>LOD:</b><br>Lamotrigine: 0.259 µg/ml<br>Clozapine: 0.205 µg/ml<br><b>LOQ:</b><br>Lamotrigine: 0.786 µg/ml<br>Clozapine: 0.679 µg/ml   |    |
| 9 | Clozapine(CLZ),<br>Olanzapine(OLZ),<br>Risperidone(RIP),<br>Quetiapine(QTP) in<br>plasma | <b>HPLC-MS/ESI<br/>Method</b> | <b>Detection Wavelength:</b><br><b>Mobile Phase:</b> Formic Acid(2.70 mmol/l),<br>Ammonium Acetate(10 mmol/l) in water –<br>Acetonitrile (53:47)<br><b>Stationary Phase:</b> C <sub>18</sub> (2 mm×125 mm, 3<br>µm)<br><b>Linearity Range:</b><br>CLZ: 20-1000 ng/ml<br>OLZ: 1-50 ng/ml<br>RIP: 1-50 ng/ml<br>QTP: 20-1000 ng/ml<br><b>Retention Time:</b><br>CLZ: 0.9903<br>OLZ: 0.9993<br>RIP: 0.9979<br>QTP: 0.9985<br><b>Flow Rate:</b> 0.16 ml/min | 41 |

**Conclusion:**

This review depicts the reported Spectrophotometric and Chromatographic methods; developed and validated for estimation of Atypical Antipsychotics. According to the literature review it was concluded that for Clozapine and Aripiprazole different Spectroscopic & Chromatographic methods are available for Single component as well as for combination. This all methods found to be simple, accurate, economic, precise, and reproducible in nature. Most of Methods were of RP-HPLC and UV absorbance detection because these methods provided with best available reliability, repeatability, analysis time and sensitivity.

There is no reported method for Aripiprazole and Clozapine in synthetic mixture. So there will be a great scope for development of highly Precise, Accurate, Simple as well as rapid analytical methods for latest drugs such as Aripiprazole and Clozapine.

**Acknowledgement:** The authors are thankful to Dr. K. Pundarikakshudu, Director of L. J. Institute of Pharmacy, Ahmedabad, India for providing all the facilities and encouragement to carry out the work.

**Reference:**

1. Rang H., Dale M., Ritter J., Flower R. Rang Dale's pharmacology; 6<sup>th</sup> Edn; Elsevier Publication, 2007, pp 545-555.
2. Tripathi KD. Essentials of Medical pharmacology; 6<sup>th</sup> Edn; Jaypee Brothers Medical publishers Pvt Limited, New Delhi, 2010, pp 424-425, 432-433.
3. George MK., Praveen A. Medicine; 4<sup>th</sup> Edn; Elsevier Publication, 2012, pp 673-674.
4. Maryadele NJ. The Merck Index an Encyclopedia of chemicals drugs and biological; 14<sup>th</sup> Edn; Merck Research Laboratories, UK, 2006, pp 129, 408
5. Drugbank, "Aripiprazole(DB01238) Drug profile", Oct 2016  
<http://www.drugbank.ca/drugs/DB01238>

6. Drugbank, "Clozapine(DB000363) Drug profile", Oct 2016  
<http://www.drugbank.ca/drugs/DB00363>
7. Indian Pharmacopoeia. Government of India Ministry of Health and Family Welfare Published by Indian Pharmacopoeia Commission, Ghaziabad, 2014, Vol-II, pp 1081-1082, 1448-1450
8. United State pharmacopoeia and National formulary. Asian Edition, The United State Pharmacopoeial Convention, Rockville, June 2016, USP
9. Kalalchelvi R., Thangabalan B., Srinivasa R., Jayachandran E. "UV Spectrometric Determination of Aripiprazole in Bulk and Pharmaceutical Formulation", *E-Journal Of Chemistry*, **2009**,6(S1):587-590
10. Jain R., Kashaw SK., Rishab J., Mishra P., Kohli D.V. "Visible Spectrophotometric Method For Determination of Aripiprazole in Tablets", *IJPS* **2011 Jan-Feb**;73(1):74-76
11. Nagamallika J., Dr. Aruna M., "Development and Validation of Spectrophotometric estimation of Aripiprazole in Tablet Dosage Form", *Asian J. Pharm. Ana.* **2011**,1(3)
12. Akram M., Sameh M., Ismail I. "Extractive Spectrophotometric Method for the Determination of Some Antipsychotic Drugs Using Erichrome Black T" *Journal of Applied Pharmaceutical Science* **2015 June**;5(6):26-33
13. Sastri BS., Gananadhamu S., Devala Rao G. " RP-HPLC Determination of Aripiprazole In Pharmaceutical Formulation", *Asian journal Of Chemistry* **2009**;21(9):6643-6646
14. Simran D., Nitesh C., Dr. Malairajan P., Murugan R., Rakhal Chandra D., shafique A. "A Simple and Rapid Spectrophotometric Determination of Aripiprazole in Pharmaceutical Dosage Form", *IJDDR* **2011 Jan-March**;3(1):205-208
15. Srinivasa NR., Srinivas P., Venkataramana, Venkateswara R. " Analytical Method Development Report For Aripiprazole" *International Journal Of Advances In Pharmacy, Biology, Chemistry*, **July-sep 2012**; 1(3):377-380
16. Ashu M., Shikha P., Nagarajan k. "Development and Validation of Rapid HPLC Method for Determination of Aripiprazole in Bulk drug and pharmaceutical dosage forms", *Med.Ugal.*
17. Bhadru B., Srinath P., Kedarnath J. "Development, Estimation and Validation in bulk and its pharmaceutical formulation by HPLC Method", *Int.J. ChemTech Res.* **2012 Jan-March**;4(1):124-128
18. Kunal M., Dhamecha, Reena V., Satish K. "A Rugged and Economic Method for the Estimation of Aripiprazole in Human Serum By LCMS/MS Detection for Clinical trials", *International Journal of Pharmacy & Pharmaceutical Sciences* **2011 Oct**;3:121-123
19. Ravindra N., Indrajeet S., Kumaraswamy G. "New RP-HPLC Method for the Estimation of Aripiprazole in Bulk and in Pharmaceutical Dosage Forms", *IAJPR* **2014**;4(4):1842-1849
20. Ahmed G., Fatma M., Abdel G., Eman F.M., "Spectrophotometric Determination of Aripiprazole in Tablets by Charge-Transfer and Ion-Pair Complexation Reactions with Some Acceptors" *Asian J. Pharm. Ana.* **2012**;2(1):12-19
21. Subbayamma AV., Ramababu C. "Spectrophotometric Determination of aripiprazole in pharmaceutical formulations with MBTH and ferric chloride", *Oriental Journal of Chemistry* **2008**;24(2):677-680
22. Daxesh P., Primal S., Mallika S., Shrivastav PS. "SPE-UPLC-MS/MS method for sensitive and rapid determination of aripiprazole in human plasma to support a bioequivalence study" *Journal of chromatography. B, Analytical technologies in the biomedical and life sciences* **2013 Feb**; 925C:20-25
23. Ali M. "Sensitive Detrmination of Aripiprazole using chemiluminescence reaction of tris(1,10-phenanthroline)ruthenium(II) with acidic Ce(IV)", *Analytical Methods* **2014 Oct**;6:9588-9595
24. Ravinder S., Akula T., Mukkanti K., Raju D., Laxmi V., Dendhi C.R. "Development and Validation of an LC-MESI-MS Method for Quantitative Determination of Aripiprazole in human plasma and an Application to Pharmacokinetic Study" *Journal of Chromatographic Science* **2012 Oct**
25. Kareem M. "Stability Indicating Spectrophotometric and TLC-Densitometric Methoda for Determination of Aripiprazole in Bulk and Dosage Form", *IJPPS* **2014** 6(4):542-548
26. Chia-Jui T., Hsien-Jane C., El-Wui L., Jau-Tay W., Chin-Haung C., Tsuo-Haung L., "The quantitative detection of Aripiprazole and its main metabolite using capillary-electrophoresis", *Journal of the Chinese Medical Association* **2011 June**;74(6):267-271

27. Yumiko A., Norio YF., Midori K., Yoshimasa I., Tsukasa U. "A sensitive column-switching HPLC method for aripiprazole and dehydroaripiprazole and its application to human pharmacokinetic studies", *Journal of Separation science* **2010 Nov**; 33(21):3292-3298
28. Divya S., Hasumati R., Neelam P. "Development and Validation of Analytical Method for Aripiprazole and Escitalopram Oxalate by Simultaneous Equavation Spectroscopic Method" *Inventi Rapid: Pharm Analysis & Quality Assurance* **2016 April-June**; 2:1-5
29. Zia A., Fazal S., Saba A., Qazi R., Sagheer A., Muhhamad S., Farooq S. "Development Of fixed dose combination tablets of aripiprazole plus divalproex sodium and their simultaneous determination using HPLC-UV", *Drug DevInd Pharm*, **2016 Sep**; 42(9):1393-405
30. Sri Ramya N., Vijayalakshmi R., Dhanraju MD., "Spectrophotometric Determination of Aripiprazole and Tapentadol using chloranilic acid Reagent", *IJPSR* **2015 May**;6(5):2052-2055
31. United State pharmacopoeia and National formulary. Asian Edition, The United State Pharmacopoeial Convention, Rockville, 2004, USP 27 NF 22, 498-499.
32. European Pharmacopoeia. Council of Europe, 2005, Vol-5, pp 1337-1338
33. Uma P., Bijaya G. "Reverse phase Liquid chromatographic method for the Estimation of clozapine from tablet dosage forms", *IJPRIF* **2009 July-Sept**;1(3):733-736
34. Chokchai W., Wandee T., Sayam K., Siriluk S., Saowarunee S. "Determination of Clozapine in Human Plasma by HPLC with UV-VIS Detector", *CMU. J. Nat. Sci.* **2010**;9(1):29-37
35. Manoj GT. "Estimation of Clozapine in Human plasma by HPLC and detection by UV-VIS detector" *Asian Journal of Biochemical and Pharmaceutical Research* **2012**;2(2):49-53
36. Akram M., Sameh M., Ismail I. "Extractive Spectrophotometric Method for the Determination of Some Antipsychotic Drugs Using Erichrome Black T" *Journal of Applied Pharmaceutical Science* **2015 June**;5(6):26-33
37. Ibrahim D., Hanaa AW., Niveen AL. "Validated Spectrophotometric and Fluorimetric Methods for Analysis of Clozapine in Tablets and Urine" *Annali di Chimica* **2005**;95:345-356
38. Micheal R., Priscilla S., Ric P., Alasdair MB., Kishor MW., "Determination of clozapine and its Metabolite, Norclozapine in Various Biological Matrices Using HPLC" *Drug Development and Industrial Pharmacy* **2008 Sep**;33(10):1158-1166
39. Avenso A., Facciola G., Campo GM., Fazio A., Spina E. "Determination of clozapine, desmethylclozapine and clozapine N-oxide in human plasma by RPLC with UV Detection", *J. Chromatogr B Biomed Sci Appl.* **1998 Sep** 4;714(2):299-308
40. Priyanka PA., Hasumati AR., Vineet CJ. "Development and Validation of Analytical Method for Simultaneous estimation of Lamotrigine and Clozapine in Synthetic Mixture by absorbance Correction Method" *Pharmatutor* **2015**;3(6):41-48
41. Zhou Z, Li X, Li K, Xie Z, Cheng Z, Peng W, Wang F, Zhu R, Li H. "Simultaneous Determination Of Clozapine, Olanzapine, Risoeridone and Quetiapine in Plasma by HPLC- Electrospray Ionization Mass Spectrometry" *J Chromatogr B Analyt Technol Biomed Life Sci.* **2004 April** 5;802(2):257-62