

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Obstetrics and Gynecology Research

Journal homepage: www.ijogr.org

Case Report

A rare case of pedunculated vulval angiomfibroblastoma

Madhurya R M^{1,*}, Jayanthi Thammiah¹, Chethana Mannem²

¹Dept. of Obstetrics and Gynaecology, Kempegowda Institute of Medical Sciences, Bangalore, Karnataka, India

²Dept. of Pathology, Kempegowda Institute of Medical Sciences, Bangalore, Karnataka, India



ARTICLE INFO

Article history:

Received 06-09-2022

Accepted 27-03-2023

Available online 18-05-2023

Keywords:

Angiomfibroblastoma

Vulvovaginal area

Benign mass

Excision biopsy

Pedunculated

ABSTRACT

A rare benign mesenchymal tumour that predominantly occurs in lower genital tract, more frequently vulvo vaginal areas of young to middle-aged women, rarely occurs in retroperitoneum. It needs to be differentiated from other benign conditions like Gartner's duct cyst, epithelial inclusion cyst, fibroepithelial polyps. It grows more slowly and is non tender, non invasive, and have least tendency for local recurrence. The histologic findings of the tumors are abundant thin-walled blood vessels with hypocellular and hypercellular areas. Almost all tumor cells have immunoreactivity for both desmin and vimentin. It also has estrogen and/or progesterone receptors, but staining for cytokeratin is negative. Here is a case of AMFB of the vulva occurring in a 62-year-old woman, involving the left labia majora. The patient described that her vulva mass grew over few years. The dimension of the tumor was measured as 18*12*4 cms. Excision biopsy done and sent for histopathology.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

Angiomfibroblastoma is a benign mesenchymal tumour that occurs predominantly in subcutaneous tissue of vulvovaginal region of premenopausal women and very rarely in the inguinoscrotal region of men.¹ While majority of the tumors are encountered in the vulva, 10% to 15% of them occur in the vagina, occasionally in the perineum, inguinal region and retroperitoneum.² Differential diagnosis of mesenchymal neoplasms in the vulvovagina includes an array of lesions that are overlapping in both their morphology and immunohistochemistry like lipoma, epithelial inclusion cyst, fibroepithelial polyps and locally invasive aggressive angiomyxoma.³

2. Case Report

A 62-year old postmenopausal female, P₄L₄, presented with a painless vulval mass that started few years before, which had gradually grown to present size. She was a known hypertensive on medication. No significant past and family history.

Clinical examination showed a normal general physical examination and local examination showed a pedunculated mass, arising from lower part of left labia majora, soft to firm in consistency, measuring approximately 10*10 cms, pedicle measured approximately 8-10 cms.(as shown in Figure 1) Pelvic examination was normal.

2.1. Differential diagnosis

Vulval fibroma, vulval lipoma.

* Corresponding author.

E-mail address: snk.madhurya@gmail.com (Madhurya R M).

2.2. USG

Solid heterogenous lesion from left labia measuring 18*10*4 cms with volume of 490 – 500cc with significant internal vascularity, probably angiomyoibroblastoma or vulval lipoma or labial sarcoma.

After relevant investigations, mass was excised and sent for histopathology.

2.3. Gross findings

Well demarcated, soft tissue mass measuring approximately 18*13 cms. Firm in consistency.



Fig. 1: Benign Angiomyofibroblastoma

2.4. Cut section

Bright yellow homogenous fatty tissue with fine fibrous capsule and trabeculae (as shown in Figure 2). Specimen sent for histopathology, reported as Angiomyofibroblastoma

2.5. HPE

Benign lesion with alternating hypercellular and hypocellular areas admixed with blood vessels (Figure 3 b). Spindle cells and plump stromal cells are seen aggregating around the vessels.(Figure 3 a) Bundles of muscle fibres and mature adipose tissue can be seen.(as shown in Figure 3 c)



Fig. 2:




Fig. 3:

3. Discussion

1. AMF is a benign, soft-tissue tumor, its cause and pathogenesis are unknown at present. Since AMF is rare, no statistical data are available about its incidence in the general population. Most cases of AMF are benign, and only one case with sarcomatous transformation has ever been reported.⁴
2. On perineal USG, appear as a soft-tissue mass with inhomogeneous mixed echogenicity, which corresponds to the cellular inhomogeneity found on

histopathological examination.

3. Immunohistochemistry-tumors express estrogen and progesterone receptors, which suggests that it might have originated as a neoplastic proliferation of hormonally responsive mesenchymal cells, also positive to desmin, vimentin and smooth muscle actin.^{5,6}

4. Conclusion

Although the exact nosologic position of angiomfibroblastoma is still surrounded by some controversy and requires further elucidation. Simple excision appears sufficient in its management. Whereas wide excision after histological examination is needed for the management of the aggressive angiomfibroma which is associated with a high risk of local recurrence and infiltration.

5. Source of Funding

None.

6. Conflict of Interest

None.

References

1. Giannella L, Costantini M, Mfuta K, Cavazza A, Cerami LB, Gardini G, et al. Pedunculated angiomfibroblastoma of the vulva: case report and review of the literature. *Case Rep Med*. 2011;2011:893261. doi:10.1155/2011/893261.
2. Hutchens K, Angiomfibroblastoma, Pathologyoutlines.
3. Qiu P, Wang Z, Li Y, Cui G. Giant pelvic angiomfibroblastoma: Case report and literature review. *Diagn Pathol*. 2014;9:106. doi:10.1186/1746-1596-9-106.
4. Nielsen GP, Young RH, Dickersin GR, Rosenberg AE. Angiomfibroblastoma of the vulva with sarcomatous transformation ("angiomfibrosarcoma"). *Am J Surg Pathol*. 1997;21(9):1104–8.
5. Nasu K, Fujisawa K, Takai N, Miyakawa I. Angiomfibroblastoma of the vulva. *Int J Gynecol Cancer*. 2002;12:228–31.
6. Horiguchi H, Matsui-Horiguchi M, Fujiwara M, Kaketa M, Kawano M, Ohtsubo-Shimoyamada R, et al. Angiomfibroblastoma of the vulva: report of a case with immunohistochemical and molecular analysis. *Int J Gynecol Pathol*. 2003;22(3):277–84.

Author biography

Madhurya R M, -

Jayanthi Thammiah, Professor and HOD

Chethana Mannem, Professor

Cite this article: Madhurya R M, Thammiah J, Mannem C. A rare case of pedunculated vulval angiomfibroblastoma. *Indian J Obstet Gynecol Res* 2023;10(2):221-223.