

Content available at: <https://www.ipinnovative.com/open-access-journals>

Indian Journal of Clinical Anaesthesia

Journal homepage: www.ijca.in

Original Research Article

A randomized controlled study on efficacy and safety of air versus alkalinized 2% lignocaine for inflating endotracheal tube

Bhagyashri Tappe¹, Charushila Patkar¹, Sucheta Lokare¹, Deepak Phalgune^{2,*}¹Dept. of Anaesthesiology, Poona Hospital & Research Centre, Pune, Maharashtra, India²Dept. of Research, Poona Hospital & Research Centre, Pune, Maharashtra, India

ARTICLE INFO

Article history:

Received 25-01-2021

Accepted 28-02-2021

Available online 10-09-2021

Keywords:

Postoperative sore throat

Lidocaine

Air

Endotracheal tube

ABSTRACT

Introduction: The postoperative sore throat (POST) occurs in approximately $\geq 50\%$ of individuals after general anaesthesia. An attempt is made in the present research to evaluate the efficacy of alkalinized 2% lignocaine with conventional air as cuff inflating media, in preventing POST and coughing in patients undergoing a surgical procedure under general anaesthesia.

Materials and Methods: This randomised double-blinded controlled study was conducted in sixty patients. In Group A, the endotracheal tube (ETT) cuff was filled with air, whereas in Group B, it was filled with 2% lignocaine. Coughing and POST were recorded immediately, at 1 hour, 12 hours and 24 hours. We have compared coughing, POST following tracheal extubation, and volume of inflation medium and intra-cuff pressure at the start and at the end of surgery between the two groups.

Results: The mean volume at the end of the surgery was 7.9 mL and 4.5 mL in Group A and Group B respectively (p -value = 0.001). The mean intra-cuff pressure at the end of the surgery was 32.5 cm and 19.8 cm in Group A and Group B respectively (p -value = 0.001). The percentage change in the volume at the end of the surgery was + 49.7 and -12.7% in Group A and Group B respectively (p -value = 0.001). The percentage change in the intra-cuff pressure at the end of the surgery was + 62.5 and -1.0% in Group A and Group B respectively (p -value = 0.001). The incidence of coughing at one hour postoperative was 26.7% and 3.3% in Group A and Group B respectively (p -value = 0.026).

Conclusion: The intracuff alkalinized lidocaine is useful adjunct to endotracheal intubation.

This is an Open Access (OA) journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprint@ipinnovative.com

1. Introduction

There are many airway complications associated with tracheal intubation, or extubation after general anaesthesia.¹ In roughly $\geq 50\%$ of patients the postoperative sore throat (POST) is observed.¹ The occurrence of vigorous coughing, agitation, or restlessness, increase intracranial, intra-thoracic, or intra-abdominal pressure, bronchospasm, wound dehiscence and bleeding are also noted after coming out from general anaesthesia.¹ During postoperative care patients may develop complications such as hoarseness,

dysphonia, or dysphagia.¹ Lignocaine is one of the most commonly used drugs for preventing POST and its efficacy was evaluated.² When lignocaine is injected into the endotracheal tube (ETT) cuff, it spreads through the cuff membrane and induces anaesthetic action in the trachea.³ This increases airway tolerance to tracheal tubing. After tracheal extubation, the hemodynamic alterations are minimized resulting in reducing the incidence of coughing. Only the non-ionized base form of the drug diffuses across the hydrophobic polyvinyl chloride walls of the ETT cuff. Sodium bicarbonate increases the unionized form of local anaesthetics, which causes an increase in the diffusion of it through the cuff, thereby reducing the dosage of local

* Corresponding author.

E-mail address: dphalgune@gmail.com (D. Phalgune).

anaesthetics.³ An attempt is made in the present research to evaluate the efficacy of alkalinized 2% lignocaine with conventional air as cuff inflating media, in preventing POST and coughing in patients undergoing a surgical procedure under general anaesthesia.

2. Materials and Methods

We conducted this randomised double-blinded controlled study from April 2019 to October 2019 in operation theatre, postoperative care unit (PACU) and wards. The study was approved by the institutional ethics committee (Letter No: RECH/EC/2019-20/66). Before enrolling the patients for the study, we took written informed consent. The risks and benefits of the procedure were explained to the patients.

2.1. Inclusion criteria

Patients of either gender having age between 18 and 65 years, duration of surgery up to 90 minutes, Cormack-Lehane grade I-II and American Society of Anaesthesiologist (ASA) grades I and II.

2.2. Exclusion criteria

Patients with a laryngeal disease/ laryngeal surgery, a history of smoking and sore throat seven days previous to surgery, patients with a history of severe gastro-oesophageal reflux disease, patients who had difficult intubation/failed intubation

We initially included 75 patients as per the inclusion criteria. Fifteen patients were excluded. The remaining 60 patients were randomly divided into two equal groups of 30 each, with the help of www.randomizer.org (Fig 1). ETT cuff was filled with air and 2% lignocaine 2 mL (40 mg) made alkalinized with 1.5% sodium bicarbonate 3 mL (the sodium bicarbonate 7.5% was available, which was diluted five times) to prevent air leak during positive pressure ventilation guided with cuff manometer in Group A and Group B respectively. A senior anaesthesiologist supervised the aforesaid procedure. Patients and observer were blinded for the study. The presence or absence of coughing, and POST immediately, at 1 hour, 12 hours and 24 hours were recorded by the observer.

The detailed pre-anaesthesia check-up was conducted for fitness which included airway assessment as mentioned in the study proforma to look for any signs of difficult intubation which could contribute as an independent factor for the POST. In the operation theatre, adequate intravenous (IV) access was confirmed. Minimum mandatory monitors such as non-invasive blood pressure, pulse oximeter, and electrocardiography were attached. Surgery was performed under standard general anaesthesia protocols. Premedication was done by Inj. glycopyrrolate (4µg/kg) and Inj. fentanyl (2µg/kg). Patients were pre-oxygenated for three minutes. Induction was done with Inj. propofol (2

mg/kg). After confirming that the patient can be ventilated by the mask (100% oxygen given for 2-3 minutes) a long-acting muscle relaxant Inj. vecuronium (0.08-0.1 mg/kg) or Inj. atracurium (0.5 mg/kg) was administered.

Atraumatic direct laryngoscopy was performed. Insertion of appropriate sized Portex ETT was done under direct vision till cuff went beyond the vocal cords. For female patients, 7.5 mm ETT and for male patients 8.5 mm ETT was used. Confirmation of ETT placement was carried out by auscultation of the chest, chest rise after ventilation and capnography monitoring. The cuff pressure at the start of the surgery was approximately 20 cm of H₂O.

Anaesthesia was maintained with oxygen: nitrous oxide 50: 50 and sevoflurane with end-tidal concentration maintained between 1.5% - 1.8% (adjusted according to hemodynamic parameters) with controlled ventilation. End-tidal carbon dioxide was maintained between 30% - 35%. Muscle relaxant supplemental dose was given if required. After completion of the surgery, the patient was reversed with Inj. neostigmine (0.05 mg/kg), and Inj. glycopyrrolate (0.008 mg/kg) IV. After proper nasopharyngeal and oropharyngeal suctioning, ETT was removed during inspiration. The total duration of anaesthesia, volume and cuff pressure of both the groups were noted.

Coughing, POST and volume of inflation medium, and intra-cuff pressure at the start and at the end of surgery were the primary and secondary outcome measures respectively. Gaur P et al reported that the incidence of POST was 40.0% and 8.0% in air and lignocaine group respectively.⁴ The sample size was calculated by formula⁵ $N = \{2p_{av}(1-p_{av})(Z_{\alpha} + Z_{\beta})^2\} / \Delta^2$. We have taken Z_{α} a standard normal variate at 5% type I error (1.96) and Z_{β} the standard normal deviate for β power 80% at type II error (0.84). A sample size of 30 patients in each group was calculated by above method.

2.3. Statistical analysis

Data collected were entered in Excel 2007. Statistical Package for Social Sciences for Windows, Version 20.0 from IBM Corporation, Armonk, NY, USA was used for the analysis of the data. The comparison of categorical and continuous variables was done using Chi-Square test/Fisher's exact test and student's t-test respectively. The confidence limit for significance was fixed at 95% level with p-value < 0.05.

3. Results

Of 75 patients assessed for eligibility, 15 were excluded because patients with a history of sore throat seven days prior to surgery (2), smokers (5), patient with the laryngeal disease (1), severe gastro-oesophageal reflux disease (1) and difficult intubation/failed intubation (6) (Figure 1). The patients were randomized into two groups. ETT cuff was

Table 1: Baseline characteristics

Characteristics	Group A N = 30	Group B N = 30	Total	P value
Mean age in years \pm SD	41.8 \pm 11.9	43.6 \pm 14.0		0.593*
Gender (%)				
Male	12 (40.0)	16 (53.3)	28 (46.7)	0.301**
Female	18 (60.0)	14 (46.7)	32 (53.3)	
Mean BMI in Kg/m² \pm SD	26.1 \pm 2.7	25.4 \pm 1.9		0.282*
ASA grade (%)				
Grade I	21 (70.0)	20 (66.7)	41 (68.3%)	0.781**
Grade II	9 (30.0)	10 (33.3)	19 (31.7%)	
Mean duration of anaesthesia in min. \pm SD	138.4 \pm 9.0	140.3 \pm 12.2		0.502*

*Unpaired 't' test was used

**Chi square test was used

SD- Standard deviation

BMI- Body mass index

ASA - American Society of Anaesthesiologist

Table 2: Comparison of mean volume, and mean pressure of inflation medium

Variable	Group A (n=30)	Group B (n=30)	p-value
Mean volume in mL \pm SD			
At the start of surgery	5.3 \pm 0.5	5.2 \pm 0.4	0.549
At the end of surgery	7.9 \pm 0.9	4.5 \pm 0.5	0.001
% Change	49.7% (Rise)	-12.7% (Fall)	0.001
Mean pressure in cm \pm SD			
At the start of surgery	20.0 \pm 0.0	20.0 \pm 0.0	0.999
At the end of surgery	32.5 \pm 2.3	19.8 \pm 4.1	0.001
% Change	62.5% (Rise)	-1.00% (Fall)	0.001

Unpaired 't' test was used

SD –Standard deviation

filled with air and 2% lignocaine 2 mL (40 mg) made alkalized with 1.5% sodium bicarbonate 3 mL in Group A and Group B respectively.

Mean age, gender, mean BMI, ASA grades and mean duration of anaesthesia were comparable in both the groups with no statistically significant difference (Table 1).

The mean volume at the start of the surgery was 5.3 mL and 5.2 mL in Group A and Group B respectively (p-value = 0.549). The mean intra-cuff pressure at the start of the surgery was 20.0 cm in both the groups. The mean volume at the end of the surgery was 7.9 mL and 4.5 mL in Group A and Group B respectively (p-value = 0.001). The mean intra-cuff pressure at the end of the surgery was 32.5 cm and 19.8 cm in Group A and Group B respectively (p-value = 0.001). The percentage change in the volume at the end of the surgery was + 49.7 and -12.7% in Group A and Group B respectively (p-value = 0.001). The percentage change in the intra-cuff pressure at the end of the surgery was + 62.5 and -1.0% in Group A and Group B respectively (p-value = 0.001) (Table 2). The incidence of coughing immediately postoperative, 12 hours and 24 hours postoperatively did not differ significantly between the two study groups, whereas

the incidence of coughing at one hour postoperative was 26.7% and 3.3% in Group A and Group B respectively (p-value = 0.026) The incidence of POST at one hour, 12 hours and 24 hours postoperatively did not differ significantly between the two study groups (Table 3).

4. Discussion

Coughing and POST are the most bothersome postoperative complaints in patients undergoing general anaesthesia.^{1–3} An increased intra-cuff pressure is a major contributing factor for coughing and POST.³ Lignocaine is a local anaesthetic. Being liquid, the lignocaine prevents the transmission of N₂O into the cuff, it also permeates through the semipermeable membrane of polyvinylchloride cuff. Thus, it provides a calming effect on tracheal mucosa and reduces pressure-induced problems.^{2,3,6}


Jaichandran et al. concluded that maximum diffusion of lignocaine across the ETT cuff occurred at pH 7.4, and cough receptors are blocked in the tracheal mucosa. This technique can be used for surgeries of less than two hours duration.⁷ If nitrous oxide is used as a adjunct with other

Table 3: Comparison of incidence of postoperative coughing, and POST

Postoperative Coughing	Group A n=30 (%)	Group B n=30 (%)	p-value
Immediate			
Yes	11 (36.7)	4 (13.3)	0.072
No	19 (63.3)	26 (86.7)	
1 h			
Yes	8 (26.7)	1 (3.3)	0.026
No	22 (73.3)	29 (96.7)	
12 h			
Yes	3 (10.0)	0 (0.0)	0.237
No	27 (90.0)	30 (100.0)	
24 h			
Yes	1 (3.3)	0 (0.0)	0.999
No	29 (96.7)	30 (100.0)	
POST			
1 h			
Yes	6 (20.0)	2 (6.7)	0.254
No	24 (80.0)	28 (93.3)	
12 h			
Yes	5 (16.7)	0 (0.0)	0.050
No	25 (83.3)	30 (100.0)	
24 h			
Yes	2 (6.7)	0 (0.0)	0.492
No	28 (93.3)	30 (100.0)	

Fisher's exact test was used

POST- Postoperative sore throat

**Fig. 1:** Consort diagram

volatile anaesthetics, intra-cuff pressure is increased. At the beginning of the surgery, this diffuses into the cuff. When cuff pressure exceeds the tracheal mucosal capillary pressure (> 30 mm of Hg), tracheal erosion occurs and postoperative cough and sore throat are caused.⁸ Estebe et al. reported that the liquid volume removed from the cuff decreased significantly (5.9 ± 1.6 mL) for the alkalinized group (p -value < 0.001), whereas the air volume withdrawn at extubation time increased significantly in Air Group (11 ± 2.7 mL) [p -value < 0.0001].⁹ Estebe et al. reported that in the air-filled cuff with ETT, the incidence of cough was 95% whereas in the alkalinized lidocaine filled cuff it was 5%. In the air-filled cuff, the incidence of hoarseness was 80% whereas in the alkalinized lidocaine filled cuff it was 10%. They further reported that there was a decrease in the sore throat during the postoperative period when the cuff of an ETT was inflated with a low dose of alkalinized lidocaine as compared to the air. There was reduction in the sore throat during 24 hours postoperative period in alkalinized-lidocaine groups as compared to air group. (p -value < 0.0001).¹⁰ Estebe et al. concluded that due to alkalinization of intra-cuff lignocaine, the diffusion of nonionized neutral base across the hydrophobic structure of cuff membrane was increased from 1% to 65% within six hours.¹¹

It was observed that there was a significant decrease in dose requirement of lignocaine in the alkalinized form (20–40 mg) as compared to non-alkalinized lignocaine dose (200–500 mg).^{3,11,12} Therefore, in our study we used 40 mg of alkalinized lignocaine for more safety and efficacy. External lubrication in the form of lignocaine jelly over the cuff was used in the present study during intubation for the introduction of the cuff up to the vocal cords. At the time of coming out from anaesthesia, cuff lubrication in the form of jelly or topical spray was associated with unfavorable phenomena.^{2,3,13} Lignocaine spray contains additives such as l-methanol and ethanol. These additives are linked with causing POST and hoarseness.⁶

In our study, in group A, the mean percentage change in volume was 49.7%, whereas the mean percentage change in cuff pressure was 62.5%. In B group, the mean percentage change in volume was -12.7%, whereas the mean percentage change in cuff pressure was -1.00% (p -value = 0.001). In the lignocaine group the cuff volume and pressure did not change with time whereas in the air group they change considerably. Lignocaine is a liquid medium hence, hyperinflation of the cuff with N_2O is prevented during surgery.³ It was observed that lignocaine diffused through the cuff membrane in time and concentration-dependent manner and produced local anaesthetic action on tracheal mucosa, and increased ETT tolerance.¹⁴

Gaur et al. reported that the intra-cuff volume at the end of the surgery was significantly higher in the Air group as compared to lignocaine group (6.0 mL Vs 4.1 mL) [p -value < 0.001]. They further reported that the intra-cuff pressure

at the end of the surgery was significantly higher in the Air group as compared to lignocaine group (49.9 cm Vs 19.7 cm) [p -value < 0.001].⁴

In the present study, 2% lignocaine and 1.5% of sodium bicarbonate were used. Many studies have observed that variation in the concentration of sodium bicarbonate injected into the cuff had no effect on the diffusion of lignocaine. Lignocaine is known to be absorbed rapidly from tracheobronchial mucosa. For systemic lignocaine to be effective in reducing ETT discomfort, a very high plasma concentration of lignocaine is required (IV lignocaine 2 mg/Kg gives plasma lignocaine level $> 3 \mu\text{g/mL}$) than that attained in case of lignocaine diffusion with sodium bicarbonate ($< 0.08 \mu\text{g/mL}$) suggesting that improved ETT tolerance after intra-cuff alkalinized lignocaine was local rather than a systemic effect.

Gaur et al. reported that the occurrence of coughing and POST at immediately, 1 hour and 24 hours after operation was higher in the air group as compared to the lignocaine group (p -value < 0.05). In the lignocaine group, the impact of the duration of anaesthesia on the increase in cuff pressure was considerably lower.⁴

Lam et al. in a recent meta-analysis stated that the incidence of early and late POST, agitation, coughing, hoarseness and dysphonia like post-intubation emergence phenomena were lower in the lignocaine group as compared to the air group (p -value < 0.05).¹ Tanaka et al. stated that local and systemic lignocaine therapy reduced the occurrence and severity of the POST.²

POST and coughing are very stressful and turn out to be more disturbing than the surgery itself. Most frequent postoperative complications occur due to ETT intubation. Patient's bucking or coughing and friction between the tracheal mucosa/increase in ETT cuff pressure during general anaesthesia are the important causes of these morbidities. This has detrimental effects such as an increase intracranial, intrathoracic or intra-abdominal pressure, bronchospasm, wound dehiscence, bleeding, sore throat, hoarseness, or dysphonia. When the ETT cuff pressure is elevated, it compromises the blood supply of tracheal mucosa. This may lead to ciliary loss, inflammation, ulceration, hemorrhage, tracheal stenosis and trachea-esophageal fistula. It is very challenging to maintain the ideal cuff pressure during the entire surgery. ETT cuff filled lignocaine with alkalinized sodium bicarbonate may reduce these effects.

5. Limitation

Potential limitations of the study merit consideration. The sample size was small. Plasma alkalinized lignocaine levels were not measured. There was no practical way of assessing the amount of alkalinized lignocaine that diffused across the cuff. We didn't grade the severity of coughing and POST, but only noted the incidence. Further multi-centric studies

with a large sample size are needed to validate our results.

6. Conclusion

The mean volume at the start of the surgery was 5.3 mL and 5.2 mL in Group A and Group B respectively (p-value = 0.549). The mean volume at the end of the surgery was 7.9 mL and 4.5 mL in Group A and Group B respectively (p-value = 0.001). The mean intra-cuff pressure at the end of the surgery was 32.5 cm and 19.8 cm in Group A and Group B respectively (p-value = 0.001). The percentage change in the volume at the end of the surgery was +49.7 and -12.7% in Group A and Group B respectively (p-value = 0.001). The percentage change in the intra-cuff pressure at the end of the surgery was +62.5 and -1.0% in Group A and Group B respectively (p-value = 0.001). The incidence of coughing at one hour postoperative was 26.7% and 3.3% in Group A and Group B respectively (p-value = 0.026). The incidence of POST at one hour, 12 hours and 24 hours postoperatively did not differ significantly between the two study groups.

7. Source of Funding

None.

8. Conflict of Interest

None.

References


1. Lam F, Lin YC, Tsai HC, Chen TL, Tam KW, Chen CY. Effect of intracuff lidocaine on postoperative sore throat and the emergence phenomenon: A systematic review and meta-analysis of randomized controlled trials. *PLoS One*. 2015;10:e0136184.
2. Tanaka Y, Nakayama T, Nishimori M, Sato Y, Furuya H. Lidocaine for preventing postoperative sore throat. *Cochrane Database Syst Rev*. 2009;3:CD004081.
3. Navarro LH, Braz JR, Nakamura G, Lima RM, Fde PS, Módolo NS. Effectiveness and safety of endotracheal tube cuffs filled with air versus filled with alkalized lidocaine: A randomized clinical trial. *Sao Paulo Med J*. 2007;125:322–8.
4. Gaur P, Ubale P, Khadanga P. Efficacy and safety of using air versus alkalized 2% lignocaine for inflating endotracheal tube cuff and its pressure effect on incidence of postoperative coughing and sore throat. *Anesth Essays Res*. 2017;11:1057–63.
5. Moutulsky H. *Intuitive Biostatistics*. New York: Oxford University Press; 1995.
6. Fagan C, Frizelle HP, Laffey J, Hannon V, Carey M. The effects of intracuff lidocaine on endotracheal-tube-induced emergence phenomena after general anesthesia. *Anesth Analg*. 2000;91:201–206.
7. Jaichandran VV, Angayarkanni N, Karunakaran C, Bhanulakshmi IM, Jagadeesh V. Diffusion of lidocaine buffered to an optimal pH across the endotracheal tube cuff-an in-vitro study. *Indian J Anaesth*. 2008;52:536–40.
8. Manissery JJ, Shenoy V, Ambareesha M. Endotracheal tube cuff pressures during general anaesthesia while using air versus a 50% mixture of nitrous oxide and oxygen as inflating agents. *Indian J Anaesth*. 2007;51:24–7.
9. Estebe JP, Dollo G, LeCorre P, LeNaoures A, Chevanne F, LeVerge R, et al. Alkalinization of intracuff lidocaine improves endotracheal tube-induced emergence phenomena. *Anesth Analg*. 2002;94:227–30.
10. Estebe JP, Gentili M, LeCorre P, Dollo G, Chevanne F, Ecoffey C, et al. Alkalinization of intracuff lidocaine: Efficacy and safety. *Anesth Analg*. 2005;101:1536–41.
11. Estebe JP, Treggiari M, Richebe P, Joffe A, Chevanne F, LeCorre P. In vitro evaluation of diffusion of lidocaine and alkalized lidocaine through the polyurethane membrane of the endotracheal tube. *Ann Fr Anesth Reanim*. 2014;33:73–7.
12. Matias E. Effect of NaHCO₃ on the diffusion of lignocaine through the wall of endotracheal tube's cuff. *Br J Anaesth*. 1995;74:72.
13. Maruyama K, Sakai H, Miyazawa H, Toda N, Iinuma Y, Mochizuki N. Sore throat and hoarseness after total intravenous anaesthesia. *Br J Anaesth*. 2004;92:541–3.
14. Estebe JP, Delahaye S, LeCorre P, Dollo G, LeNaoures A, Chevanne F, et al. Alkalinization of intra-cuff lidocaine and use of gel lubrication protect against tracheal tube-induced emergence phenomena. *Br J Anaesth*. 2004;92:361–6.

Author biography

Bhagyashri Tappe, Senior Resident

Charushila Patkar, Consultant Anaesthesiologist

Sucheta Lokare, Consultant Anaesthesiologist

Deepak Phalgune, Research Consultant  <https://orcid.org/0000-0003-4225-8010>

Cite this article: Tappe B, Patkar C, Lokare S, Phalgune D. A randomized controlled study on efficacy and safety of air versus alkalized 2% lignocaine for inflating endotracheal tube. *Indian J Clin Anaesth* 2021;8(3):381–386.