

Study of serum malondialdehyde and vitamin E in chronic bronchitis patients

Nagaraj R Shetkar^{1*}, Prashant Talikoti²

¹Associate Professor, ²Assistant Professor, Dept. of Biochemistry, ¹Khaja Banda Nawaz Institute of Medical Sciences (KBNIMS), Kalaburagi, Karnataka, ²Employees State Insurance Corporation Medical College (ESICMC), Kalaburagi, Karnataka, India

***Corresponding Author: Nagaraj R Shetkar**

Email: nagarajshetkar@gmail.com

Received: 24th September, 2018

Accepted: 18th January, 2019

Abstract

Aim: The oxidative stress caused by free radicals and reactive oxygen species released during smoking can play a significant role in pathogenesis and development of Chronic Bronchitis. This study is conducted to evaluate serum levels of Malondialdehyde (MDA), a lipid peroxidation product as a marker of oxidative stress and vitamin E major antioxidant in body to evaluate antioxidant status in chronic bronchitis patients and healthy controls.

Materials and Methods: Study comprised of 60 subjects, 30 healthy controls and 30 chronic bronchitis cases were included. Kei Satoh method used for measuring MDA and Serum vitamin E was estimated by method of Baker and Frank.

Results: Serum MDA a lipid peroxidation product from cell membrane considered as marker of oxidative stress was elevated in chronic bronchitis patients when compared to healthy controls where as serum level of vitamin E major antioxidant significantly decreased in chronic bronchitis patients when compared to controls.

Conclusion: MDA maker is of oxidative injury significantly increased in chronic bronchitis cases which is likely to be associated with active smoking habit and also be due to systemic inflammation. The decrease in level of antioxidant capacity in chronic bronchitis patients appears likely consequence of higher oxidative stress and increased consumption of antioxidants.

Clinical Significance: By stopping smoking and advising antioxidant rich food may halt the further progression of chronic bronchitis disease.

Keywords: Chronic bronchitis, MDA, Vitamin E.

Introduction

Chronic Bronchitis is described by chronic inflammation of the lungs mainly in bronchi. It includes one of the two variants of Chronic Obstructive Pulmonary Disease (COPD). COPD patients are higher in areas where smoking habit is common. In India, consumption of cigarette and tobacco is high so chronic bronchitis is thought to be one of the major health concern.¹ Thoracic Society of America defines Chronic Obstructive Pulmonary Disease as “A disease state that characterized by the obstruction to airflow caused due to chronic bronchitis or emphysema; here airflow obstruction is progressive and accompanied hyper-reactivity of airways in lungs and may be reversible partially.”²

Chronic bronchitis is clinically diagnosed by increased bronchial mucus secretion and is featured by productive cough daily for continuous 3 months or more in at least two consecutive years.² A present hypothesis in pathogenesis of COPD is due to excess oxidative stress consequence of current active smoking practice and also due to spilling of reactive oxidative compounds from airspace WBCs which are not precisely counter balanced by respiratory antioxidant defense mechanism, leading to elevated oxidative stress in lungs. Increased formation of oxidative compounds lead to increased pro inflammatory gene expression and also tissue injury by oxidants proceed to COPD formation.³

Malondialdehyde with the chemical formula $\text{CH}_2(\text{CHO})_2$. Free radicals and other oxidants hydrolyse PUFA

molecules present in cell membrane leading to formation of Malondialdehyde. MDA is very reactive and is one of the electrophilic compound that can cause toxic damage in tissues. MDA induces damage to tissues by formation reactive compounds which are considered as ALE (advanced lipoxidation end products).⁴

On other side deficiency of antioxidants also leads increase in oxidative stress. Antioxidants protects body from highly reactive oxygen and nitrogen compounds and also brings changes in the inflammatory reactions that can cause a significant role in development of chronic bronchitis.⁵ Lipid soluble antioxidant vitamin E is body's major defense against oxidative stress in preventing cell membrane injury. Vitamin E functions by breaking the chain reaction of lipid peroxidation reactions.⁶ The trial is conducted to study serum levels Malondialdehyde and serum vitamin E in controls and in chronic bronchitis patients.

Materials and Methods

A cross sectional study of serum Malondialdehyde and serum vitamin E was carried out in controls and Chronic Bronchitis cases from medicine outpatient department, JJM medical college and Bapuji hospital Davangere for a period of one year from April 2009 to March 2010. Institutional ethics committee approval was taken. Study and informed consent was taken from participants.

Inclusion Criteria

- Cases: 30 chronic bronchitis cases which were diagnosed included (clinical and radiologically).
- Controls: 30 normal healthy individuals without any past history of smoking habit.

Exclusion Criteria

- Patients with chronic lung disease such as asthma
- Patients with history of recent surgical operation
- Patients with history of DM
- Patients with recent history of major illness

Collection of Blood Samples

About 5ml of venous blood collected under aseptic conditions in plain vial for biochemical tests at admission. Immediately after collecting sample serum is separated properly stored.

The following tests carried out in cases and controls.

- Serum Malondialdehyde(MDA)
- Serum Vitamin E

Estimation of Serum Malondialdehyde

Serum Malondialdehyde estimation carried out by method Kei Satoh.⁷ It is characterized by principle of auto-oxidation of PUFA molecules present in cell membrane leading to formation of semistable peroxides, these reactive compounds undergo a series chain reactions to form malondialdehyde (MDA). MDA undergo reaction with compound TBA(thiobarbituric) to produce pink colored substance. This resulting colored substance is extracted with n-butyl alcohol and the absorbance is measured at 530 nm.

Estimation of Serum Vitamin E


Serum vitamin E was measured by Baker and Frank method⁸ by reduction of ferric to ferrous ion which forms red colored complex with alpha bipyridyl.

Statistical Analysis

Results are expressed as mean \pm SD. The unpaired 't' test is used for comparing different biochemical parameters between cases and controls. The 'P' value of < 0.05 was considered as statistical significance.

Results

A total number of 60 subjects were included in the study of which 30 consist of cases and 30 were controls. Among 30 controls, 20 were male and 10 were females and their



mean age is 57.4 ± 7.4 years and among 30 chronic bronchitis cases, 23 were male and 7 were females and their mean age is 59.3 ± 7.8 years.

Table 1 shows biochemical characteristics of the study subjects. Serum mean level of MDA a marker of lipid peroxidation was significantly ($p < 0.001$) elevated in chronic bronchitis patients when compared to controls. Mean level of vitamin E major antioxidant was significantly ($p < 0.001$) decreased in chronic bronchitis patients than in controls.

Table 1: Comparison of Serum Malondialdehyde and vitamin E in Controls and chronic bronchitis cases

| Groups | | MDA (nmol/ml) | Vitamin E (mg/dl) |
|---------------------------------|-----------------|-----------------|-------------------|
| Controls | Mean \pm SD | 2.61 ± 0.51 | 1.42 ± 0.29 |
| Chronic bronchitis | Mean \pm SD | 4.84 ± 0.53 | 0.81 ± 0.15 |
| Controls vs. chronic bronchitis | Mean difference | 2.23 | 0.61 |
| | t* | 19.20 | 10.65 |
| | P | < 0.001 | < 0.001 |


* Unpaired t- test, P value < 0.001 considered as highly significant

Discussion

The findings from present trial observed that there is an elevated oxidative stress and decrease in antioxidant levels in chronic bronchitis cases when related to healthy controls. Oxidative stress caused by free radicals released from active smoking habit involved in tissue injury and development of COPD.⁹

Smoking is one of the major factor in pathogenesis and formation of chronic bronchitis. Smoking is directly associated with chronic bronchitis cases worldwide. In developing countries such as India 40% of chronic bronchitis cases are related with non smoking causes such as fuel burning examples include usage of wood and cow dung and burning crop wastages from farm which releases of reactive oxygen compounds into the atmospheric air.¹⁰

Airway obstruction is vital clinical finding in identification



of chronic bronchitis. In development of disease initially elastin tissue in lungs undergo hydrolysis leading to decrease in elastic recoiling capacity in the lungs, Flow of air in lungs mainly dependent on elastic recoil capacity, the injury to elastin tissue in chronic bronchitis causes airway obstruction air-trapping in lungs and fibrosis of the airways also causes airway narrowing. Further Consequences of these changes leading to development of chronic bronchitis disease.^{11, 12}

Chronic bronchitis is also associated with systemic inflammation. The systemic inflammation which happens in chronic bronchitis disease is due to release of inflammatory compounds from the lungs into the systemic circulation through the thin layered respiratory vasculature that can expose the other tissues of the body to inflammatory consequences.¹³

MDA is produced during oxidation of PUFA molecules in cell membrane by free radicals. Free radicals can damage all organic and inorganic compounds in body but PUFA present in cell membranes are the most susceptible. Cell membranes are rich in PUFA molecules which are attacked by free radicals and reactive oxygen species.¹⁴ Chronic bronchitis patients are associated with increased oxidative stress and higher level of MDA. When correlated with healthy controls, chronic bronchitis cases have significantly (P value < 0.001) higher level of MDA. Results here are in conferring with the research done by M.K. Daga et al,¹⁵ and Gamze kirkil et al.¹⁶

When associated to healthy controls chronic bronchitis cases have significantly reduced(P value <0.001) level of vitamin E. This is in conferring with the research done by Anita MR. et al¹⁷ and Rupali SP et al.⁶

Vitamin E lipid soluble non enzymatic antioxidant. It breaks chain reaction of lipid peroxidation reactions by reacting with more-reactive lipid product and make them to form stable products. In this study, we observed that reduced level of vitamin E in chronic bronchitis patients mainly appears to be consequences of increased utilization because of excess release of reactive free radicals by cigarette smoking and also due to inflammatory reaction present in chronic bronchitis disease.¹⁸

Tobacco smoking causes release of reactive free radicals which is directly associated in the pathogenesis and progression of Chronic Bronchitis disease.¹⁵ Increased MDA concentration and reduced level of antioxidant vitamin E in patients with Chronic bronchitis is due to excess formation of reactive oxygen species because of current active smoking and inflammatory reaction and decrease level of antioxidants due its excess utilization for neutralizing free radicals.

Conclusion

In present study, we have observed that there is increased oxidation of biomolecules in chronic bronchitis patients mainly consequence of current active smoking habit and decrease in antioxidant capacity when related to healthy controls. This study demonstrates the possible effects of imbalance in oxidative stress and antioxidant capacity in pathogenesis of chronic bronchitis.

Clinical Significance

By stopping tobacco smoking and consumption of food rich in antioxidants may prevent the further oxidative stress induced alveolar damage in chronic bronchitis patients.

Conflict of Interest: None.

References

1. Premanand R, Santhoshkumar PH, Mohan A. Study of thiobarbituric reactive substances and total reduced glutathione as indices of oxidative stress in chronic smokers with and without chronic obstructive pulmonary disease. *J Chest Dis Allied Sci* 2007;49:9-12.
2. Mephie SJ, Papadakis MA, Lawrence MT. Current medical diagnosis and treatment. 47th edition, McGraw Hill Medical Publisher, New Delhi, 2008; pp. 216-221.
3. Drost EM, Skwarski KM, Saulea J, Soler N, Roca J, Agusti A, et al. Oxidative stress and airway inflammation in severe exacerbations of COPD. *Thorax* 2005;60:293-300.
4. Malondialdehyde from Wikipedia, the free encyclopaedia (serial online) cited 27th march.2017. available from :<http://en.wikipedia.org/wiki/Malondialdehyde>.
5. Mac Nee W. Oxidants / Antioxidants and COPD. *Chest* 2000;117:303S-317S.
6. Rupali SP, Subhodhini AA, Borale P, Rahul L. Study of Correlation of pulmonary function test with the markers of oxidative stress and non-enzymatic antioxidants in chronic obstructive pulmonary disease (COPD) patients. *Br J Med Med Res* 2014;4(28):4710-22.
7. Satoh K. Serum lipid peroxide in cerebrovascular disorders determined by a new colorimetric method. *Clinica Chimica Acta* 1978;90:37-43.
8. Baker, Frank. Determination of serum tocopherol by colorimetric method. In Varley's practical clinical biochemistry, 6th edition: Heinmann professional publishing 1988;902.
9. Gerritsen WBM, Asin J, Zanen P, van den Bosch JMM, Haas FJLM. Markers of inflammation and oxidative stress in exacerbated chronic obstructive pulmonary disease patients. *Respir Med* 2005;99:84-90.
10. Bill B B, Rahul K. Risk Factors and Pathophysiology of Chronic Obstructive Pulmonary Disease (COPD). *JAPI* 2012;60.
11. O'Donnell DE, Laveneziana P. Physiology and consequences of lung hyperinflation in COPD. *Eur Respir Rev* 2006;15:61-7.
12. Shapiro S. The pathophysiology of COPD: What go wrong and why? *Proceedings. Adv Stud Med* 2003;3(2B): S91-S8.
13. Fabbri LM, Rabe KF. From COPD to chronic systemic inflammatory syndrome? *Lancet* 2007; 370:797-9.
14. Ozbay B, Dulger H. Lipid peroxidation and antioxidant enzymes in Turkish population: Relation to age, gender, exercise, and smoking. *Tohoku J Exp Med* 2002;197:119-24.
15. Daga MK, Chhabra R, Sharma B, Mishra TK. Effects of exogenous vitamin E supplementation on the levels of oxidants and antioxidants in chronic obstructive pulmonary disease. *J Biosci* 2003;28(1):7-11.
16. Kirkil G, Muz MH, Seckin D, Sahin K, Kucuk O. Antioxidant effect of zinc picolinate in patients with chronic obstructive pulmonary disease. *Respir Med* 2008;102:840-4.

17. Anita MR, Adinath NS, Dilip M. A study of oxidative stress, thiol proteins and role of vitamin E supplementation in chronic obstructive pulmonary disease (COPD). *Al Ameen J Med Sc* 2013; 6(2):134-7.
18. Jyoti B, Sudeep Kumar, Yogesh T and Ranjana Singh. Study of Pulmonary Function Test, Oxidative Stress Marker and Non-enzymatic Antioxidants in Chronic Obstructive Pulmonary Disease. *Sch J App Med Sci* 2016;4(4D):1371-4.

How to cite this article: Shetkar NR, Talikoti P. Study of serum malondialdehyde and vitamin E in chronic bronchitis patients. *Int J Clin Biochem Res* 2019;6(2):157-60.