



CODEN [USA]: IAJPBB

ISSN: 2349-7750

INDO AMERICAN JOURNAL OF PHARMACEUTICAL SCIENCES

http://doi.org/10.5281/zenodo.1245715

Available online at: <http://www.iajps.com>**Research Article**

A CROSS SECTIONAL STUDY; GERD AS A RISK FACTOR OF ACUTE EXACERBATION OF COPD

¹Humna Khalid, ²Ayesha Zahid, ³Muhammad Waseem Zahid

Bahawal Victoria hospital, Bahawalpur Pakistan

Abstract:

Objective: The study is based to assess the prevalence of the GERD in the patients of COPD.

Design and place: It is cross-sectional study done at the department of Pulmonology in the Bahawal Victoria Hospital, Bahawalpur.

Duration: July 2016-Feb 2017

Methodology: 103 patients were enrolled in the study fulfilling the inclusion criteria and established diagnosis of the disease i.e. patients with age more than 40, with >20 pack years of smoking history and FEV/FVC < 0.7

Results: In the study 35% of patients were in the age ranging 40-55 years and 65% were of age >55 years. The mean age of patients was 66.87 ± 8.71 , out of 103 patients 88% were males and 12% were females.

Conclusion: It was concluded from the study results that 42% of the patients of COPD suffered from the GERD as well. So, in order to put a control on the Exacerbations of the disease surveillance of the GERD should be done of the each and every COPD patient.

Key Words: Exacerbation, COPD, Disease Surveillance, Pulmonary, Vagal Nerve stimulation

Corresponding author:

Humna Khalid,

Bahawal Victoria hospital,

Bahawalpur,

Pakistan.

QR code



Please cite this article in press Humna Khalid et al., A Cross Sectional Study; GERD as A Risk Factor of Acute Exacerbation of COPD, Indo Am. J. P. Sci, 2018; 05(05).

INTRODUCTION:

COPD is one of the major health concerns of the developing and the developed countries. It is predicted to be the third leading cause of disease burden of the world by 2030. It is characterized by the persistent and progressive limitation of the air flow to lungs and abnormal inflammatory response of the tissue of respiratory system to gases or irritating/noxious Particles. The exacerbations determine the quality of life of a COPD patient. The exacerbation is characterized by acute increase in frequency of cough, dyspnea, severity, and change in the amount/color/smell of the sputum.

The exacerbations are highly morbid and limit the activity of the patient badly, affecting the quality of life and cause burden on the patient economically, physically and Psychologically. There are a number of risk factors of the acute exacerbation of COPD, one of them is Gastro esophageal Reflux disease. GERD is characterized by reflux of the stomach contents into the esophagus due to lower esophageal sphincter incapability. This may cause a number of pulmonary complications like aspiration pneumonia, chronic, cough, Bronchial asthma, fibrosis and Acute Exacerbation of COPD. Vagal Nerve stimulation (causing Broncho spasm) and Micro Aspirations Contribute to Pulmonary complications.

A study done in Lahore shows 39% of the COPD patient suffered from GERD as well. The percentage is quite high in a study done in Turkey and it showed that the patients who suffered from GERD have more frequent exacerbations as compared to Non-GERD patients and they consequently are on intensive therapy for the COPD. This study is focused to

explore and assess the frequency of GERD in COPD patients in order to highlight an important risk factors of the disease that may be helpful for the health care professionals to manage the disease appropriately and morbidity may be decreased.

OBJECTIVES: The objective of the study was to assess the frequency of GERD in COPD patients.

DEFINITIONS: GOLD's criteria were used for the patients to be enrolled in the study. Age > 40 years Smoking history more than 20 pack years.
FEV1/FVC < 0.7

GERD: The frequency of the GERD was evaluated by a preformed questionnaire. Four options were given ranging from none to always, to scale the frequency of the GERD. Scoring of the answers were done as 0 for none and 4 for always. And patients having score more than 8 were considered GERD positive.

INCLUSION CRITERIA: All patients fulfill GOLD criteria definition of the CORD were included in the study.

Exclusion criteria: Patients having asthma other pulmonary diseases and esophageal diseases like Achalasia, Gastric/Esophageal cancer and peptic ulcer.

RESULTS: 103 patients were enrolled for the study. Among them 35 % were 40-55 years of age and rest >55 years of age. 12% were females and 88% were males. 39% of the patients came positive with GERD and 61% had no GERD.

| GENDER | NO OF PATIENTS | % |
|--------|----------------|----|
| MALE | 91 | 88 |
| FEMALE | 12 | 12 |

| GERD | NO OF PATIENTS | % |
|------|----------------|----|
| YES | 41 | 39 |
| NO | 59 | 61 |

DISCUSSION:

COPD is major health concern of the world accounting more than 20% of the total burden of the diseases.

GERD is one of the risk factors of the acute exacerbations of the disease. A number of studies have been done on the topic and all show a correlation of the GERD and COPD. The data of the study not only shows the correlation but also that the GERD is the main cause of the exacerbations of disease causing morbidity and poor quality of life.

The study was based on highlighting the GERD as the major risk factor for acute exacerbation of COPD. So that it may help healthcare professionals manage the disease in a better way.

In our study it was explored that 39% of COPD patients suffered from GERD. The results are in accordance with another study done in Karachi showing that 41% of COPD patients suffered from GERD. The severity of the GERD increases as the severity of the COPD increases. There are more patients with severe COPD having severe GERD

symptoms and this frequency increases as pack year history of smoking increases. Another study done in Iran shows that there is more use of acid suppressive drugs among COPD patients than in control group.

CONCLUSION:

It is concluded that the frequency of GERD is directly related to COPD. So, every patient presenting with COPD should undergo surveillance and sorted out for GERD. So that, this treatable disease can be managed appropriately.

REFERENCES:

1. Pauwels RA, Buist AS, Calverley PM, Jenkins CR, Hurd SS. The GOLD Scientific Committee. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease. NHLBI/WHO Global Initiative for Chronic Obstructive Lung Disease (GOLD). Am J Respir Crit Care Med 2001; 163:1256–1276.
2. CrossRefPubMedGoogle Scholar
3. Locke GR, Talley NJ, Fett SL, Zinmeister AR, Melton LJ. Prevalence and clinical spectrum of gastroesophageal reflux: a population based study in Olsted County, Minnesota. Gastroenterology 1997; 112:1448–1456.
4. CrossRefPubMedWeb of ScienceGoogle Scholar
5. Lagergren J, Bergstrom R, Lindgren A, Nyren O. Symptomatic gastroesophageal reflux as a risk factor for esophageal adenocarcinoma. N Engl Med 1999; 340:825–831.
6. CrossRefPubMedWeb of ScienceGoogle Scholar
7. Harding SM, Richter JE, Guzzo MR, Schan CA, Alexander RW, Bradley LA. Asthma and gastroesophageal reflux: acid suppressive therapy improves asthma outcome. Am J Med 1996; 100:395–405.
8. CrossRefPubMedWeb of ScienceGoogle Scholar
9. Harding SM, Guzzo MR, Richter JE. The prevalence of gastroesophageal reflux in asthma patients without reflux symptoms. Am J Respir Crit Care Med 2000; 162:34–39.
10. PubMedWeb of ScienceGoogle Scholar
11. Ing AJ, Ngu MC, Breslin ABX. Chronic persistent cough and GERD. Thorax 1991; 46:479–483.
12. Abstract/FREE Full TextGoogle Scholar
13. Theodoropoulos DS, Ledford DK, Lockey RF, et al. Prevalence of upper respiratory symptoms in patients with symptomatic gastroesophageal reflux disease. Am J Respir Crit Care Med 2001; 164:72–76.
14. PubMedWeb of ScienceGoogle Scholar
15. Gislason T, Janson C, Vermeire P, et al. Respiratory symptoms and nocturnal gastroesophageal reflux: a population-based study of young adults in three European countries. Chest 2002; 121:158–163.
16. CrossRefPubMedWeb of ScienceGoogle Scholar
17. Tobin RW, Pope CE, Pellegrini CA, Emond MJ, Sillery J, Raghu G. Increased prevalence of gastroesophageal reflux in patients with idiopathic pulmonary fibrosis. Am J Respir Crit Care Med 1998; 158:1804–1808.
18. CrossRefPubMedWeb of ScienceGoogle Scholar
19. De Vault KR, Castell DO. For the Practice Parameters Committee of the American College of Gastroenterology. Guidelines for the diagnosis and treatment of the gastroesophageal reflux disease. Arch Intern Med 1995; 155:2165–2173.
20. CrossRefPubMedWeb of ScienceGoogle Scholar
21. Adhami T, Richter JE. Twenty-four-hour pH monitoring in the assessment of esophageal function. Semin Thorac Cardiovasc Surg 2001; 13:241–254.
22. CrossRefPubMedGoogle Scholar
23. David P, Denis P, Nouvet G, Pasquis P, Lefrancois R, Morére P. Fonction respiratoire et reflux gastro-cesophagien au cours de la bronchite chronique. [Respiratory function and gastro-esophageal reflux during chronic bronchitis]. Bull Eur Physiopathol Respir 1982; 18:81–86.
24. PubMedWeb of ScienceGoogle Scholar
25. Duculoné A, Vandevenne A, Jouin H, et al. Gastroesophageal reflux in patients with asthma and chronic bronchitis. Am Rev Respir Dis 1987; 135:327–332.
26. PubMedWeb of ScienceGoogle Scholar
27. Orr WC, Shamma-Othman Z, Allen M, Robinson MG. Esophageal function and gastroesophageal reflux during sleep and waking in patients with chronic obstructive pulmonary disease. Chest 1992; 101:1521–1525.
28. CrossRefPubMedWeb of ScienceGoogle Scholar
29. Mokhlesi B, Morris AL, Huang CHF, Curcio AJ, Barrett TA, Kamp DW. Increased prevalence of gastroesophageal reflux symptoms in patients with COPD. Chest 2001; 119:1043–1048.
30. CrossRefPubMedWeb of ScienceGoogle Scholar
31. Stein MR, Towner TG, Weber RW, et al. The effect of theophylline on the lower esophageal sphincter pressure. Ann Allergy 1980; 45:238–241.
32. PubMedWeb of ScienceGoogle Scholar
33. Crowell MD, Zayat EN, Lacy BE, Schettler-Duncan A, Liu MC. The effects of an inhaled β_2 -adrenergic agonist on lower esophageal function. Chest 2001; 120:1184–1189.
34. CrossRefPubMedWeb of ScienceGoogle Scholar
35. Vigneri S, Termini R, Leandro G, et al. A comparison of five maintenance therapies for reflux esophagitis. N Engl J Med 1995; 333:1106–1110.

36. CrossRefPubMedWeb of ScienceGoogle Scholar
37. De Meester TR, Johnson LF, Joseph GJ, Toscano MS, Hall AW, Skinner DB. Patterns of gastroesophageal reflux in health and disease. *Ann Surg* 1976; 184:459–469.
38. CrossRefPubMedWeb of ScienceGoogle Scholar
39. Jamieson JR, Stein HJ, Demeester TR, et al. Ambulatory 24-h esophageal pH monitoring; normal values, optimal thresholds, specificity, sensitivity, and reproducibility. *Am J Gastroenterol* 1992; 87:1102–1110.
40. PubMedWeb of ScienceGoogle Scholar
41. Mahler D, Weels C. Evaluation of clinical methods for rating dyspnoea. *Chest*. 1988; 93:580–586.
42. CrossRefPubMedWeb of ScienceGoogle Scholar
43. Ferrer M, Alonso J, Prieto L, et al. Validity and reliability of the St. George's Respiratory Questionnaire after adaptation to a different language and culture: The Spanish example. *Eur Respir J* 1996; 9:1160–1166.
44. AbstractGoogle Scholar
45. Field SK, Evans JA, Price LM. The effects of acid perfusion of the esophagus on ventilation and respiratory sensation. *Am J Respir Crit Care Med* 1998; 157:1058–1062.
46. PubMedWeb of ScienceGoogle Scholar
47. Cuttitta G, Cibella F, Visconti A, Scichilone N, Bellia V, Bonsignore G. Spontaneous gastroesophageal reflux and airway patency during the night in adult asthmatics. *Am J Respir Crit Care Med* 2000; 161:177–181.
48. PubMedWeb of ScienceGoogle Scholar
49. Alexander JA, Hunt LW, Patel AM. Prevalence, pathophysiology, and treatment of patients with asthma and gastroesophageal reflux disease. *Mayo Clin Proc* 2000; 75:1055–1063.
50. CrossRefPubMedWeb of ScienceGoogle Scholar
51. Molld JW, Reed LE, Davis AB, Allen ML, Decktor DL, Robinson M. Prevalence of gastroesophageal reflux in elderly patients in a primary care setting. *Am J Gastroenterol* 1991; 86:965–970.
52. PubMedWeb of ScienceGoogle Scholar