

# Diagnostic omics based molecular techniques in oral pathology: Current perspectives and future prospects

Bijo Alexander

Professor and Head, Dept. of Oral Pathology and Microbiology, Noorul Islam College of Dental Sciences, Trivandrum, Kerala, India

\*Corresponding Author: Bijo Alexander

Email: drbijo73@gmail.com

## Abstract

The completion of human genome project has been a landmark event in the development of diagnostic pathology and personalized medicine. Like other fields of medicine, oral pathology also tends to show a paradigm shift from conventional diagnostic techniques to a molecular based format. Omics based molecular techniques can be accurately used for diagnosis, prognosis, risk analysis and treatment of various oral lesions especially oral cancer. This will go a long way in providing a more precise diagnosis which in turn will give the patient a risk free treatment protocol tailored for his individual needs. The paper analyses the evolution of oral pathological diagnostic techniques with an impetus into the current omics based techniques and a peep into the future development prospects.

**Keywords:** Oral cancer, Omics based, Molecular.

## Introduction

Basic biomedical laboratory techniques are often the bed rock on which clinical practice is based. Compared to conventional diagnostic techniques newer methods exhibit more accuracy and depth which helps the clinician in justifying his decision making.<sup>1</sup> The Flexner report published in 1910 linked the scientific principles underlying human biology and disease to the clinical decision making that is central to effective medical practice.<sup>2</sup> There is a trend to apply omics based techniques to improve human health by accelerating drug discoveries and motivating life styles.<sup>3</sup> Even though omics based diagnostic techniques hold great promise several factors discourage their use over conventional diagnostic methods especially in third world countries. These include lack of skilled man power,<sup>4</sup> lack of infrastructure like clinics and laboratories,<sup>5</sup> poor computer connectivity, unfavorable health policies,<sup>6</sup> lack of data and even religious beliefs.<sup>7</sup> Hence it is imperative that newer techniques are made more cost effective so that better clinical care can be provided to the patient.

## Pre-omics Diagnostic Techniques in Oral Pathology

Oral cancer has always been an extremely challenging diagnostic dilemma for oral pathologists as its prognosis depends mainly on its staging. Almost half of the cases of oral cancer have been diagnosed in TNM stages III and IV.<sup>8</sup> Oral and oropharyngeal cancers usually have a survival rate of less than 50% after 5 years but show great improvement if diagnosed early.<sup>9</sup> Cancer mortality is affected by delay in diagnosis especially if it is done in advanced stages beyond the sequelae of treatment.<sup>10</sup> Hematoxylin & Eosin staining backed by use of special stains and good clinicopathologic acumen is often considered the gold standard in diagnosis of oral lesions. The next innovation was the use of immunohistochemistry which is rarely used for a primary diagnosis but is most significant when applied for poorly differentiated neoplasms and in categorization of lymphomas.<sup>11</sup> However there are severe limitations of immunohistochemistry like variable antibody reactivity,

staining, poor quantification and subjective interpretation.<sup>12</sup> Traditional biochemistry has used several approaches for diagnosis including ELISA, gene cloning, conventional and real time qualitative PCR, chromosomal cytogenetic analysis, comparative genomic hybridization, karyotyping, electrophoresis and fluorescence in situ hybridization (FISH).<sup>13</sup> These techniques have enabled identification of molecular markers of several diseases. The angiogenic marker cluster of CD34 serves as an important predicting tool for recurrent OSCC.<sup>14</sup> Saliva has been used as a potential agent in detection of oral diseases and several studies have identified small ncRNAs in them which is a biomarker for cancer.<sup>15</sup> Elevated expression of TGF and EGFR in premalignant lesions with dysplasia was used as an intermediate marker of malignancy.<sup>16</sup> Kang et al<sup>17</sup> showed an enhanced BMI (B lymphoma –MLV insertion region 1 homolog), an oncogene expression in premalignant lesions.

## Omics based Diagnostic Techniques Genomics

Cancer based genomics is a rapidly growing discipline in which the genetic molecular basis of malignancy is studied at the scale of whole genomes and successfully identified specific oncogenes and tumor suppressors.<sup>18</sup> Genomic alterations have been identified for leukoplakia as well as in the process in sequential cancer tumorigenesis.<sup>19</sup> Even though specific oncogenes and tumor suppressor genes have been identified, therapy for cancer management strategies' is still at its infancy.<sup>18</sup>

## Genome wide Association Studies (GWAS)

The genome wide association study attempts to discover novel genes affecting an outcome by testing a large number of genetic variants for association. GWAS is usually hypothesis generating rather than hypothesis testing. While GWAS discoveries can be directly used for clinical risk prediction on rare occasions, the more typical scientific path is for GWAS to generate gene that may be associated with a

disorder. GWAS studies in oral health including periodontal diseases, salivary flow rate and dental fear are currently under way. Notable association for oral health include PVT1/GSDMC locus on chromosome 8q24 associated with nonsyndromic CLL with or without CLP. IRF6 gene associated with cleft lip/palate, ACTN2 gene with regulatory ameloblasts during enamel formation and TFIP11, associated with extracellular enamel matrix has been identified by GWAS.<sup>20</sup>

### Whole Exome Sequencing (WES)

Sequencing the entire protein coding regions is called exome sequencing. It is a promising omics tool for rapid identification of functional variants.<sup>21</sup> Al-Hebshi et al.<sup>22</sup> used WES technique in addition to confirming known genes of oral squamous cell carcinoma identified several novel driver events and pathways providing further evidence for genetic heterogeneity of oral cancer. Stephens et al (2013)<sup>23</sup> and Nichols AC(2012)<sup>24</sup> used whole exome sequencing in study of adenoid cystic carcinoma and HPV positive an negative head and neck squamous cell carcinoma

### Next Generation Sequencing (NGS)

Next generation sequencing, massively parallel or deep sequencing are related terms that denote a DNA sequencing technique where an entire human genome can be sequenced within a day.<sup>25</sup> Progress in genome sequencing technologies post the genome project completion in 2013 has led to decreased cost per megabase and increase in number and diversion of sequence genomes. Probing genomes in greater depth can lead to enhanced understanding of how genome sequence variants underlie phenotype and disease.<sup>26</sup> Use of NGS has enabled researchers in identifying alterations in OSCC. Whole genome sequencing studies have identified alterations with TP53, CDKN2A, PIK3CA and HRAS genes. NGS was used to identify NOTCH 1 gene involved in squamous differentiation.<sup>27</sup> Sweeney et al<sup>28</sup> used NGF metod and found that mutations in SMO gene coded smooth protein was more common in maxillary ameloblastomas while BRAFV600E mutations were commoner in mandibular ameloblastomas.

### Microbiomics

Microbiotic cells amounting upto 10 to 100 trillion within a single human form a significant propotion of the total living cells in the body. The population of our microbial occupants is known as our microbiome. The greater uniqueness of individual microbiome can help us tackle health issues pharmacogenetically with greater ease.<sup>29</sup> The adoption of an ecological approach for maintenance of oral health with a shift to personalized strategies to refrubrich health of helpful oral microbiota subsequent to the dysbiosis after treatment of oral ailments like periodontitis.<sup>30</sup>

### Proteomics

Chi et al<sup>31</sup> used mass spectrometry based quantitative proteomic analysis by way of labeled mass spectrometry completed to high performance liquid chromatography

system (HPLC) to reveal signaling pathway for oral cancer cells in vitro. Their findings reveal that interferon signaling pathway is significantly altered in OSCC lesions. The role of Interferon in pathogenesis of OSCC as a potential biomarker and for targeted therapy should be explored further.

### Transcriptomics

Saliva is a noninvasive accessible and efficient diagnostic media. Salivary transcriptomic analysis can detect a large panel of human DNA which has a diagnostic value in evaluation of OSCC .Li et al<sup>32</sup> analysed potential salivary RNA biomarkers IL8, IL1B, DUSP1, MA3, OAZ1, S100P and SAT and yielded 91% specificity and sensitivity in detecting OSCC from controls proving that the novel clinical approach can be a very robust reproducible tool for early cancer detection. Studies analyzing the role of HPV virus in oropharyngeal squamous cell carcinoma using different transcriptomics profiling using RNA scope to stratify subtypes of head and neck cancer.<sup>33</sup>

### Epigenomics

Genetic information such as DNA sequences have been limited to fully explain mechanism of gene regulation and disease process. Epigenetic mechanisms which include DNA methylation, histone modification and non-coding RNAs can regulate gene expression and affect disease progression. Epigenetic mechanism play important role in gene expression during developmental and pathological processes of dental diseases. Epigenetic modification can lead to dental abnormalities during teeth development. Inflammatory reaction in dental pulp and periodontal tissues affect epigenetic modifications causing change in gene expression.<sup>34</sup> Oral cancer has also been subjected to epigenetic analysis with numerous studies revealing that the development and progression of this malignancy are partially induced by altered epigenetic substrate together with genetic alterations and prolonged exposure to environmental risk factors.<sup>35</sup>

### Lipidomics

Lipids play critical functions in cellular survival, proliferation, interaction and death since they are involved in chemical energy storage, cellular signaling and cell-cell interactions. These cellular processes are strongly related to carcinogenesis pathways particularly to transformation, progression and metastasis suggesting the mediator role of bioactive lipids in oncogenesis.<sup>36</sup> Rapid intraoperative mass spectrometry (REIMS) is an emerging technique that allows near real time characterization of human tissue in vivo by analysis of aerosol released during electrosurgical dissection. The coupling of REIMS technology with electro surgery for tissue diagnostics is called intelligent knife (iknife) The use of iknife which is a fast, real time mass spectrometry based identification of surgical margins of tissues introperatively by differentiating tumor and normal areas and hence waiting for frozen sections can be avoided.<sup>37</sup>

## Metabolomics

Metabolomics is a core descriptive science of system biology focusing on the study of low molecular weight compounds in biological system. Analysis of human metabolome which is composed of diverse group of metabolites can aid in diagnosis and prognosis of oral squamous cell carcinoma.<sup>38</sup> Breathomics, a branch of metabolomics measures total amount of volatile organic compounds (VOCs) in exhaled air. VOCs produce in the body and released into blood and transported into the lungs where they are exhaled. Measurement of unique VOCs may indicate the presence of chronic inflammation and oxidative stress that can be used as potential biomarkers for head and neck cancer detection.<sup>39</sup>

## Ancillary Omics based Molecular Techniques

### Molecular Imaging

Imaging techniques can be used as non invasive approach to detect molecular and cellular techniques in living cell and organisms. Techniques like CT, MRI and PET can be used to screen patients with oral tumors particularly OSCC in early stages.<sup>40</sup> Antibody coated gold nanoparticles target and illuminate cancer cells under a reflectance based optical imaging system. Gold nanoparticles provide an optical contrast to discriminate normal and cancerous cells.<sup>41</sup> Oral cavity when exposed to light in UV-visible region, fluorophores absorb a portion of photons and gets excited and the lower energy photons can be detected as fluorescence. Cancerogenesis changes the concentration of fluorescence emitting properties of native fluorophores which is helpful in detecting cancer sensitivity. Autofluorescence spectroscopy is also useful to separate normal, premalignant and malignant lesions.<sup>42</sup> By using Fourier transform (FT)-Raman spectroscopy changes in vibrational band of normal, dysplastic and OSCC seem to arise from the structural, conformational and compositional structure of proteins.<sup>43</sup> Principal component analysis (PCA) of the Raman spectral data can discriminate between normal, inflammatory, premalignant and malignant conditions of oral tissues.<sup>44</sup> Confocal Reflective microscopy is a non invasive optical tool where image contrast is determined by differences in refractive indices of organelles and other subcellular structures within the tissue. It differs from conventional microscope as it illuminates a small spot within tissue with laser.<sup>45</sup> It can be used noninvasively to image tissue without fixation, sectioning and staining and can be used for pre surgical determination of cancer margins and intrasurgical guidance of patients in real time thus being valuable diagnostic tool for precancer and cancer.<sup>46</sup>

### Microfluidic System

Microfluidic system, proposed by Manz and Wedmer in 1990 also called micrototal analysis systems ( $\mu$ TAS) or "lab on a chip" have shown tremendous potential in the field of single cell analysis as the microfluidic systems have the advantages of high analysis efficiency, easy operation, small sample and reagent composition and dimensional matching

between channel sizes and cell diameter.<sup>47</sup> Microfluidic devices utilize cellular biophysical properties and devices have been developed to measure single cell mechanical or electrical properties distinguishing normal cells and their malignant counterparts. Cancer cells can be identified from clinical samples by cell affinity chromatography, magnetic activated cell sorting microfiltration and On Chip DEP (Dielectrophoresis).<sup>48</sup>

### Stem Cell Therapies

In head and neck OSCC, cancer stem cells have an integral role in tumor initiation, disease progression and treatment. It has features like self-renewal, high migration capacity, drug resistance and high proliferation ability.<sup>49</sup> Various techniques for cancer stem cell detection proposed include marker expression and cell expression by flow cytometry, side population assay, ALDH activity test, spheres test, soft agar assay and cryopreservation of tumor cells.<sup>50</sup>

### Conclusion

Due to the exponential increase in genome based techniques with declining costs it is not far when genome related visits by patients to clinics becomes the norm rather than the exception. The individual ownership of a person's genome and its formation and its rights to informed consent and privacy over sequencing, testing and disclosing genomic information is yet to gain legal recognition.<sup>51</sup> It is imperative that these molecular based techniques should be used as in an adjunct role for in depth diagnosis while maintaining conventional pathologic techniques as a gold standard.

**Conflict of Interest:** Nil.

**Source of Funding:** Nil.

### References

1. Brass EP. Basic Biomedical Sciences and the Future of Medical Education: Implications for Internal Medicine. *J Gen Intern Med* 2009;24(11):1251-4.
2. Flexner A. Medical Education in the United States and Canada. New York 1910.
3. Kumar. D. From evidence based medicine to genomic medicine. *Genomic Med* 2007;1(3-4):95-100.
4. Naickar S, Planje-Rhule J, Tutt RC, Eastwood JB. Shortage of healthcare workers in developing countries-Africa. *Ethn Dis* 2009;19(1 Suppl 1):S1-60-4.
5. Alemenji GA, Zeh C, Yao K, Fonjongo PN. Strengthening national health laboratories in Sub-Saharan Africa: A decade of remarkable progress. *Trop Med Inf Health* 2014;19(4):450-8.
6. Hagopian A. Strengthening health systems in poor countries: A code of conduct for nongovernmental organizations. *Am J Public Health* 2008;98(12):2134-40.
7. Fanning BJ, Clayton EW. Introduction: Religious and Spiritual Issues in Medical Genetics. *Am J Med Genet C Semin Med Genet* 2009;151C(1):1-5.
8. Guneri P, Epstein JB. Late stage diagnosis of oral cancer: components and possible solutions. *Oral Oncol* 2014;50(12):1131-6.

9. Kowalski LP, Carvalho AL, Martins Priante AV, Magrin J. Predictive Factors for distant metastasis for oral and oropharyngeal squamous cell carcinoma. *Oral Oncol* 2005;41(5):534-41.
10. Biazevic MGH, Castellanos A, Antunes JLF, Michel-Crosato E. Trends in oral cancer mortality in the city of Sao Paulo, Brazil, 1980-2002. *Cad Saude Publica* 2006;20(10):2105-14.
11. Fox H. Is H&E morphology coming to an end? *J Pathol* 2000;53(1):38-40.
12. Hofman F. Immunohistochemistry. *Current Protoc. Immunol* 2002;21:21-4.
13. Adeola HA, Soyele OO, Adefuye AO, Jimoh SA, Butali A. Omics based molecular techniques in oral pathology centered cancer: prospect and challenges in Africa. *Cancer Cell Int* 2017;17:61:1-12.
14. Kademani D, Lewis JT, Lamb DH, Rallis RJ, Harrington JR. Angiogenesis and cd34 expression as a predictor of recurrent squamous cell carcinoma. *J Oral Maxillofac Surg* 2009;67:1800-5.
15. Majem B, Rigau M, Reventos J, Wong DT. Non-coding RNAs in saliva: Emerging biomarkers for molecular diagnostics. *Int J Mol Sci* 2015;16:8676-98.
16. Srinivasan M, Jevell SD. Evaluation of TGF- $\alpha$  and EGFR in oral leukoplakia and submucous fibrosis by quantitative immunohistochemistry. *Oncol* 2001;69(4):2284-92.
17. Kang MK, Kim RH, Kim SJ. Elevated bmi-1 expression is associated with dysplastic cell transformation during oral carcinogenesis and is required for cancer cell replication and survival. *Br J Cancer* 2007;96:126-33.
18. Patel L, Parker B, Yang D, Zhang W. Translation genomics in cancer research: Converting profiles into personalized cancer medicine. *Cancer Biol Med* 2013;10(4):214-20.
19. Cervigne NK, Machado J, Goswami RS. Recurrent genomic alterations in sequential progressive leukoplakia and oral cancer: drivers of oral tumorigenesis? *Human Mol Genet* 2014;23(10):2618-28.
20. Shaffer JR, Feingold E, Marazita ML. Genome wide Studies- Prospects and Challenges for Oral Health. *J Dent Res* 2012;91(7):637-41.
21. Adeola HA, Soyele DO, Adefuye AO, Jimoh SA, Butali A. Omics based molecular techniques in oral pathology centered cancer: prospects and challenges in Africa. *Cancer Cell Int* 2017;17(61):1-12.
22. Al-Hebeshi NN, Li S, Nasher AT, El-Setouhy M, Al Sanosi R, Blancato J, Loffredo C. Exome sequencing of oral squamous cell carcinoma in users of Arabian stuff reveals novel candidates for driver genes. *Int J Cancer* 2016;139(2):363-72.
23. Stephens PJ, Davies HR, Mitani Y. Whole exome sequencing of adenoid cystic carcinoma. *J Clin Invest* 2013;123(7):2965-8.
24. Nichols AC, Chen-Seng-Yue M, Yoo J, Xu W. A Pilot study Comparing HPV-Positive and HPV-Negative Head and Neck Squamous cell Carcinomas by Whole Exome Sequencing. *ISRN Oncol* 2012.
25. Behjati S, Tarpey PS. What is next generation sequencing? *Arch Dis Child Educ Pract Ed* 2013;98(6):236-8.
26. Goodwin S, Mepherson JD, McCombie WR. Coming of age: ten year of next generation sequencing technologies. *Nat Rev Genet* 2016;17(6):333-51.
27. Sapna G, Gokul S. Next generation sequencing in oral disease diagnostics. *World J Stomatol* 2018;6(2):6-10.
28. Sweeny RT, McClary AC, Myers BR. Identifications of recurrent SMO and BRAF mutation in ameloblastomas. *Nat Genet* 2014;46(7):722-5.
29. Saxena R, Saxena R, Raj A. Microbiomics in the Molecular Era: A birds Eye view into the Future of Personalised Medicine. *Acta Sci Microbiol* 2018;1(8):4-39.
30. Kilian M, Chapple ILC, Hannig M. The Oral Microbiome-an update for the Oral Health Care Professionals. *Br Dent J* 2016;221:10:657-66.
31. Chi LM, Lee CW, Chang KP. Enhanced Interferon Signalling Pathways in Oral Cancer Revealed by Quantitative Proteome Analysis of Microdissected Specimens Using O<sup>16</sup>/O<sup>18</sup> Labelling and Integrated Two Dimensional LC-ESI-MALDI Tandem MS. *Mol Cell Proteomics* 2009 8(7):1453-74.
32. Yang LI, Maie AR, St John, Xiaofeng Zhou. Salivary Transcriptome Diagnostics for Oral Cancer Detection. *Clin Cancer Res* 2004;10(24):844-50.
33. Mirghani H, Amen F, Moreau F, Guigay J. Human papilloma virus testing in oropharyngeal squamous cell carcinoma. what the clinicians should know. *Oral Oncol* 2014;50(1):1-9.
34. Seo JY, Park YJ, Yi YA, Hwang JY, Lee IB. Epigenetics: General characteristics and implications for oral health. *Restor Dent Endod* 2015;40(1):14-22.
35. Irimie AI, Ciocan C, Gulei D, Mehlerov N. Current Insights into Oral Cancer Epigenetics. *Int J Mol Sci* 2018;19(3):670.
36. Perotti F, Rosa C, Cicalini I. Advances in Lipidomics for Cancer Biomarkers Discovery. *Int J Mol Sci* 2016;17(12):1992.
37. Balog G, Sasi-Szabo L, Kinron L. Intraoperative tissue identification using rapid evaporative ionization mass spectrometry. *Sci Transl Med* 2013;5(194):194ra193.
38. Sridharan G, Ramani P, Patankar S. Serum metabolomics in oral leukoplakia and oral squamous cell carcinoma. *J Cancer Res Ther* 2017;13(3):558-61.
39. van Mastricht E, Reyes-Reyes A, Brand K. Exhaled breath profiling using broadband quantum cascade laser based spectroscopy in healthy children and children with asthma and cystic fibrosis. *J Breath Res* 2016;10(2):026003.
40. Keshavarzi M, Darijani M, Momenti F, Mirzaei H. Molecular Imaging and Oral Cancer Diagnosis and Therapy. *J Cell Biochem* 2017;118(10):3055-60.
41. Kah JC, Kho KW, Lee CG. Early diagnosis of oral cancer based on the surface plasma resonance of gold nanoparticles. *Int J Nanomed* 2007;2:785-98.
42. DeVeld DC, Witjes MJ, Sternberg HJ, Roodenberg JL. The status of in vivo antifluorescence spectroscopy and imaging for oral oncology. *Oral Oncol* 2005;41:117-31.
43. Oliveira AP, Bitar RA, Silveira I, Zangaro RA, Martin AA. Near infrared Raman spectroscopy for oral carcinoma diagnosis. *Photomed Laser Surg* 2006;24:348-53.
44. Malini R, Venkatesh K, Kurien J, Pai KM, Rao L, Kartha VB, Krishna CM. Discrimination of normal, inflammatory, premalignant and malignant oral tissue: A Raman Spectroscopy Study. *Biopolymers* 2006;81:17-193.
45. White WM, Baldassano M, Rajadhyaksa M. Confocal reflective imaging in head and neck surgical specimens. A comparison with histologic analysis. *Arch Otolaryngol Head Neck Surg* 2004;130:923-8.
46. Olivio M, Bhuvaneshwari R, Keogh I. Advances in Bio-Optical Imaging for the Diagnosis of Early Oral Cancer. *Pharm* 2011;3:354-78.
47. Wu H, Zhu J, Huang Y, Wu D and Sun J. Microfluidic – Based Single Cell Study: Current status and Future Perspective. *Molecules* 2018;23(9):2347.
48. Chen J, Li J, Sun Y. Microfluidic approach for cancer detection, characterization and separation. *Lab Chip* 2012;12:1753-67.
49. Mery B, Guy JB, Esperel S, Wonzy AP. Targeting head and neck tumoral stem cells: From biological aspects to therapeutic perspectives. *World J Stem Cells* 2016;8(1):13-21.
50. Tirino V, Desiderio V, Pano F, Pappacio G, DC Rosa M. Methods for Cancer Stem Cell Detection and Isolation. *Methods Mol Biol* 2012;1879:513-29.

51. Robertson JA. The \$1000 genome: Ethical and legal issues in whole genome sequencing of individuals. *Am J Bioeth*, *AJOB*,3(3):W-1F1.

**How to cite this article:** Bijo Alexander. Diagnostic omics based molecular techniques in oral pathology: Current perspectives and future prospects. *J Oral Med, Oral Surg, Oral Pathol, Oral Radiol* 2019;5(2):57-61.