



REVIEW ARTICLE

HIGH VALUE OF BIOACTIVE COMPOUNDS FROM *CORIANDRUM SATIVUM* L FOR MULTIPLE BIOTHERAPEUTIC APPLICATIONS

Norma M. De la Fuente-Salcido¹ , David Francisco Lafuente-Rincón²

¹Autonomous University of Coahuila, Center for Interdisciplinary Studies and Research, Arteaga, Coahuila, 25354, Mexico.

²Autonomous University of Coahuila, Faculty of Biological Sciences, Torreón, Coahuila, 27275, Mexico.

Article Info:



Article History:

Received: 6 December 2024

Reviewed: 9 January 2025

Accepted: 17 February 2025

Published: 15 March 2025

Cite this article:

De la Fuente-Salcido NM, Lafuente-Rincón DF. High value of bioactive compounds from *Coriandrum sativum* L for multiple biotherapeutic applications. Universal Journal of Pharmaceutical Research 2025; 10(1): 69-79. <http://doi.org/10.22270/ujpr.v10i1.1274>

*Address for Correspondence:

Dr. Norma M. De la Fuente-Salcido,
Autonomous University of Coahuila, Center for
Interdisciplinary Studies and Research, Arteaga,
Coahuila, 25354, Mexico. Tel:+52(844)689.1058
E-mail: normapbr322@gmail.com

Abstract

Currently, people demanding nutrient-rich food products that provide additional health benefits. In this sense, *Coriandrum sativum* L, the most consumed spice around the world increasing its popularity due to its high nutritional value and confirmed pharmacological effects. Bioactive compounds derived from coriander have a wide variety of medical and biotherapeutic applications that extend beyond their well-established gastronomic relevance. Here, the information currently available on the presence, diversity, and extraction methods of the main components of coriander are reviewed. The multiple benefits of include cilantro in the diet and its effect on the treatment of global impact chronic diseases of such as cancer, heart disease, Alzheimer's disease, diabetes, kidney disease, cerebrovascular diseases, among others, are also highlighted. Likewise, the primary biological activities of coriander are detailed, such as antioxidant capacity and antimicrobial activity derived from specific phytochemicals.

This update shows the composition and describe the mixture of bioactive constituents included in the herbaceous plant and/or extracts, emphasizing the biological functions mainly as antimicrobial agents and its confirmed antioxidant function. Coriander is a reservoir of bioactivities combined, with promising potential to use and/or increase health and promote well-being.

Keywords: Antimicrobial, antioxidant, coriander, phytochemicals.

INTRODUCTION

Coriander (*Coriandrum sativum* L.) is a potential source of high-value compounds and is globally recognized for its multiple beneficial health effects, such as natural antioxidant and antimicrobial capabilities that have been successfully studied and confirmed, primarily in the fields of medicine and the food industry¹.

In this review, a brief overview of the chemical composition of Coriander leaves and seeds as well as the biological importance of phytochemical components, is summarized. In addition, concepts about the oxidation and reduction processes and/or microbial control are contextualized to explain how the phytochemicals compounds of coriander act as natural antioxidant and/or antimicrobial agents. Likewise, several important investigations that successfully evidenced the effectiveness of highlighted several important investigations that have successfully demonstrated the effectiveness of coriander phytochemicals under various experimental conditions for food preservation and medical treatments.

Coriander is a widely cultivated crop. Generally known as a popular herb, it is characterized by being an erect plant with pronounced main root, slender stems with develop branching grow up between 20 to 70 cm in height. Crop maturity is reached approximately 2 to 3 months, after that is harvested including the thin roots¹. The taxonomic data of the PLANTS database from the United States Department of Agriculture², provides standardized information that describes coriander as a flowering plant of the *Magnoliophyta* division, *Magnoliopsida* class, *Rosidae* subclass and *Apiales* order. The genus *Coriandrum* L. and the species *Coriandrum sativum* L. belong to the *Apiaceae* family, also known as the Carrot family.

It is important to know the differences between both *C. sativum* species, *vulgare* and *microcarpum*. In the *vulgare* variety produces larger fruits than the *microcarpum*, while the *vulgare* variety has lower yields of essential oils (0.1-0.35% (v/w)) than the *microcarpum* (0.8-1.8% (v/w))^{3,4}. The uses of *C. sativum* are well known in global gastronomy, food preservation, traditional medicine, and natural therapies. The coriander as a therapeutic herb date back to prehistoric times, long before these properties were

attributed to its diverse range of bioactive phytochemical components. This review includes a comprehensive overview of the taxonomic classification of coriander, the nutritional and phytochemical profile of the species, and emphasizes the chemical compounds with important biological activities, such as antimicrobial and antioxidant properties. Likewise, some generalities are included about the pharmacological properties of the coriander plant that are involved in health care and the well-being of the population.

Phytochemicals brief overview

Plants in general, including coriander, synthesize primary and secondary metabolites. Plant growth is driven by primary metabolites; on the other hand, secondary metabolites, while not essential for the general development of the plant, are crucial for adaptation and chemical protection processes. Plant secondary metabolites are called phytochemical compounds, characterized by their broad biological activities.

The type and concentration of phytochemicals synthesized by plants are determined by the species, genotype, physiology, stage of development, and environmental conditions during growth⁵. Phytochemicals, or chemicals in plants, play an important role in their growth and development, but

these are also the best source of bioactive compounds when included in people's diets.

According to their chemical structure, two main groups of phytochemicals can be distinguished. The first is a very diverse group that includes non-phenolic compounds such as carotenoids, tocopherols, phytosterols, and organo-sulfur. The second group includes phenolic compounds. Phenolic compounds are mainly isolated from leaves and seeds and are the most studied and reported in plant research⁶. Specifically, polyphenolic compounds are the main components found in aqueous extracts of coriander leaves⁷.

Plant phenolics compounds, such as carotenoids, anthocyanins, and tocopherols, have been used in different medical studies confirming their beneficial impact on human health. For instance, some antioxidant properties found in plants are attributed to vitamins, anthocyanins, phenolics and tannins⁸. Regarding flavonoids, the main identified groups are flavonols, including quercetin, kaempferol, isorhamnetin, and cyanidin as the most representative. These flavonols differ significantly in qualitative and quantitative profiles between species. Flavonoids such as flavones, isoflavones, flavonoids, flavonols, flavanones, and anthocyanins also have significant antioxidant activity⁹.

Table 1: Nutritional values of Coriander seeds and fresh leaves (per 100 g).

<i>Coriandrum sativum</i> L				
Principle	Seed		Leaves	
	Nutrients value	RDA*(%)	Nutrients value	RDA*(%)
Energy	298 Kcal	15	23 Kcal	1
Carbohydrates	54.99 g	42	3.67 g	3
Protein	12.37 g	22	2.13 g	4
Total Fat	17.77 g	60	0.52 g	2
Cholesterol	0 mg	0	0 mg	0
Dietary Fiber	41.9 g	110	2.80 g	6.50
Vitamins				
Folates	1 µg	<1	62 µg	15.50
Niacin	2.130 mg	13	1.114 mg	7
Pantothenic acid	-	-	0.570 mg	11
Pyridoxine	-	-	0.149 mg	11
Riboflavin	0.290 mg	22	0.162 mg	12
Thiamin	0.239 mg	20	0.067 mg	5.50
Vitamin A	0 IU	0	6748 IU	225
Vitamin C	21 mg	35	27 mg	45
Vitamin E	-	-	2.50 mg	17
Vitamin K	-	-	310 mcg	258
Electrolytes				
Sodium	35 mg	2	46 mg	3
Potassium	1267 mg	27	521 mg	11
Minerals				
Calcium	709 mg	71	67 mg	7
Iron	0.975 mg	108	1.77 mg	22
Magnesium	16.32 mg	204	26 mg	6.50
Manganese	330 mg	83	0.426 mg	18.50
Phosphorus	1.900 mg	82	48 mg	7
Selenium	409 mg	39	0.9 mg	2
Zinc	4.70 mg	43	0.50 mg	4.50
Phyto-components				
Carotene-α	-	-	36 µg	-
Carotene-β	-	-	3930 µg	-
Crypto-xanthin-β	-	-	202 µg	-
Lutein-zeaxanthin	-	-	865 µg	-

*Recommended Dietary Allowance. (USDA: National Nutrient data base <https://ndb.nal.usda.gov>).

Table 2: Methods for extraction of essential oils⁴²⁻⁵⁵.

Fundamental bases		General process
Hydro-distillation by Clevenger's apparatus	The oldest oil extraction by distillation using water	Fresh samples immersed into water are hydro-distilled in a Clevenger's apparatus (3–6 h)
Hydro-diffusion	Developed for large quantities of dried and sensitive to boiling plants	Collecting hydrosol and essential oils separately
Soxhlet extraction (SE)	Plant samples with solvents are heated and condensed.	Samples with hexane, methanol or acetonitrile are heated and condensed several times
Extraction in solvents	Plant oils extracted with solvents acetone, ether, ethanol, hexane, methanol and using heating.	The process includes heating, filtration, and evaporation of the solvents.
Steam distillation	Widely used method to produce large quantities of EOs (8-9 h).	Volatile oils pass through condenser, and water and oil are mixed in the liquid.
Cold pressing	Extraction of essential oils mechanical means, cold press technique	Oil is pressed at low pressures and temperatures
Microwave-assisted extraction (MAE)	Heating by absorption of microwave energy disrupts plant cells increasing the coming out rate of EOs	Plant materials are ground and soaked in solvent and the mixture is placed into a microwave oven
Super-critical fluid extraction method	The EOs are extracted from fluids with carbon dioxide and water	Reusing fluids in repeated steps of compression and decompression
Solvent-free microwave extraction (SFME)	Combines heating with microwave oven and dry distillation by atmospheric pressure without solvents	Rapid, high yield and selective EOs separation from aromatic herbs, spices and dried seeds
Microwave hydro-diffusion and gravity (MHG)	Technique based on the microwave oven heating combined with earth gravity	Uses microwaves for hydro-diffusion of plant EOs, separate and collect them with atmospheric pressure
Solar hydro-distillation system	Thermal decomposition of leaves with solar energy (120-180 min)	Vacuum-type solar energy tube or panel with a temperature controller and EOs collection unit

Flavonoids and linalool are the main bioactive compounds in the water-soluble extracts of coriander and have a significant anxiolytic effect¹⁰. Non-flavonoids or phenolic acids (benzoic and cinnamic compounds) are found in very low concentrations, can be present bound, and especially in the free form. Other compounds such as stilbenes, tannins, lignins, and lignans, also exhibit beneficial effects on health, due to their antioxidant properties^{11,12}.

Occurrence and role of major constituents of coriander

In recent decades, scientific reports on the phytochemistry of coriander have increased and diversified. Therefore, this review highlights the importance of phytochemical composition of coriander, as well as the contribution of every component to the attractive nutritional content and its impact on human health¹³. The above includes a description of the specific biological activities of coriander's main components, as well as therapeutic applications. These characteristics have generated greater attraction among habitual consumers of this spice and have led to an increase in the number of scientific investigations in recent years. First, it is important to note that there is a lot of confusion in the common nomenclature of the herb, such as cilantro or coriander, in different places around the world. While the herb is recognized as coriander in the United Kingdom, in the United States, the fresh herb is called cilantro, but the seeds are commonly known as coriander. Cilantro and coriander are undoubtedly two coined names for the same herb, and both are correct. Here, the term *C. sativum* refers to both parts of the plant equally, and when a specific part

of the plant is discussed, the terms cilantro or coriander seed will be used. The actual differences considered here focus on the chemical or phytochemical composition of the leaves, stems and seeds, whether fresh or dehydrated¹⁴. Both the seeds and the fresh leaves of this small apiaceous herb contain a wide range of chemical components (nutrients) with unique properties that are directly associated with the nature and quantity of each constituent¹⁵.

Coriander not only provides nutrients but also a large amount of heterogeneous phytochemicals (non-nutrients) that have been beneficial for well-being and preserving health since ancient times (460–377 BC)¹⁶. Sometimes these phytochemicals are inappropriately recognized as phytonutrients, however they are not essential such as fats, proteins, minerals and vitamins¹⁷. Table 1 show the outstanding nutritional value of coriander seeds and fresh leaves (per 100 g) (USDA National Nutrient data base).

Since the 1990s, more than 5,000 phytochemicals have been detected and identified, however, many of them have not been adequately studied to take advantage of the many benefits they could bring to human health^{18,19}. Phytochemicals are recognized for providing plants with unique organoleptic properties, such as their color, taste, smell, and flavor²⁰. The nature of phytochemical compounds is diverse; therefore, their classification is also complex, generally defined according to their origin, their synthesis during the stages of plant growth, or the composition of each plant product (grains, legumes, beans, fruits, herbs, nuts, roots, leaves, seeds)²¹.

Table 3: Antioxidant activity in Coriander compounds⁶⁴⁻⁷⁴.

Phytochemical compound	Bioactivity	Assay	Mechanism of action effect
Coriander EOs (β -linalool 66.07%, camphor 8.34%, geranylacetate 6.91%, cymene 6.35%)	Antioxidant	Total antioxidant capacity of biomolecules 39.38 mg TEAC/L	Radical scavenging activity (AA%=51.05 %) mg/L of Trolox equivalents
Coriander EOs (1.2%)	Antioxidant scavenging assay	Scavenged DPPH	Inhibit oxidation and quench free radicals Inhibition DPPH (I %=66.2%)
Coriander APEOs mixture with <i>Apium graveolens</i> L, <i>Thymus vulgaris</i> L	Antioxidant activity potentiation	FRAP, ABTS adical scavenging methods	AA=16.7%
Coriander SEOs (polyphenols)	Enhanced anti oxidant activity	DPPH, FRAP and cellular antioxidant activity assays	% FRAP reduction 0.951 mg/mL DPPH radical scavenging=1.053 to 2.221 mg/mL
Coriander SEOs	Antioxidant activity	Scavenged DPPH %	% DPPH radical scavenging=72.05 %
Coriandrum SEOs (100 g/mL)	Antioxidant profiling	Scavenged DPPH %	% DPPH radical scavenging=78 %
Coriander FEOs (Heneicos-1-ene, 200 ppm)	Antioxidant activity	Scavenged DPPH %	% DPPH radical scavenging=89.6
Coriandrum EOs		Scavenged DPPH % IC ₅₀ , TPC (μ g GAE/mg)	% DPPH (IC ₅₀)=756.43 μ g/mL TPC=6.20 μ g/mg
Coriander SEOs (200 μ g/mL)	Antioxidant activity	Radical scavenging DPPH, Anti-lipid peroxidative activity (PV), Iron chelating activity, IC ₅₀	% Free radical scavenging =70. % iron chelating=0.73 PV=1.64I meq/kg oil IC ₅₀ =82.1 mg/mL
Coriander SEOs	Antioxidant activity	DPPH (IC ₅₀) activity, FRAP assay, NO activity	DPPH(IC ₅₀)=121.8 μ g/mL FRAP=662.92 μ mol Fe(II)/g NO (IC ₅₀)=14.06 μ g/mL
Coriandrum SEOs	Antioxidant activity	TPC, TF, % DPPH (IC ₅₀)	TPC=126-555 GAE/g TF=64-455 mg QE/g IC ₅₀ =3.10-7.57 mg/mL
Linalool, α -pinene, β -pinene, p-cymene, γ -terpinene (seeds)	Antioxidant activity	RSA High antioxidant activity	RSA=66.48% Inhibiting oxidation processes (57.3%)

EOs: Essential oils; SEOs: Seed Essential oils; APEOs: Aerial parts essential oils; TAC: Trolox equivalent antioxidant capacity; AA: Antioxidant activity; DPPH: Difenilpicrilhidrazilo; FRAP: Ferric Reducing Antioxidant Power; ABTS: (2,2'-azino-bis-(3-ethylbenzothiazoline-6-sulfonic) acid); IC₅₀: Half maximal inhibitory concentration; TPC: Total phenolic content; TFC: Total flavonoid content; PV: Peroxide values; GAE: Gallic acid equivalents; RSA: Radical scavenging activity.

In general, the thousands of phytochemicals identified are classified as primary metabolites (chlorophyll, amino acids, proteins, common sugars or simple carbohydrates, membrane lipids, pyrimidines and purines of nucleic acids) and secondary metabolites (alkaloids, terpenoids, phenolic compounds, lignins, plants steroids, curcumins, glucosides, flavonoids, saponins), according to their role in the plant's metabolism²². Plants synthesize a wide arsenal of secondary metabolites as part of their defense system and constitute a group of molecules with a broad diversity of functions in the plant²³. These metabolites also provide plants with many beneficial properties for health of people who regularly consume them in the diet^{24,25}.

Focusing on the essential nutrients of coriander, they are currently widely involved in the formulation and production of new foods, whether fortified with the vast number of nutritional components (vitamins, minerals, proteins, fibers, among others), as well as functional and nutraceutical foods. These essential coriander compounds are also very important ingredients as a commercial strategy to enhance color and flavor in food products²⁶. On the other hand, the beneficial effects of the consumption of phytochemicals considered non-nutritive compounds of coriander, have an important role in the generation of

new biotherapeutics for the prevention of chronic diseases, and many other conditions that compromise human health and well-being, including the perspective of coriander agrobiotechnology²⁷⁻³¹.

It is important to know that the bioavailability of essential dietary nutrients and non-nutrient phytochemicals related to coriander consumption can be influenced by several factors, such as maturity, part of the herb (root, leaves, flowers, seeds), different growth stages, cultivation conditions, phytochemical extraction method, food matrix incorporation, human microbiota, and physiological state of each consumer^{21,32}.

Diversity of phytochemicals in coriander

The plant essential oils (EOs) generally contain two chemical constituents: terpenes and phenylpropanoids³³. The first group is classified into terpenes with a hydrocarbon structure, including monoterpenes, sesquiterpenes, diterpenes, and their oxygenated by-products (acids, alcohols, aldehydes, ketones, phenols, esters, oxides and lactones)³⁴. The second group comprises phenylalanine-derived molecules (C6-C3) such as flavonoids (anthocyanins, flavans, flavonones, flavanols and flavanonols, condensed tannins, and lignans), as well as non-flavonoids (non-carboxylic phenols, phenolic acids), which possess useful biological activities for medical applications³⁵.

Table 4: Antimicrobial activity in Coriander compounds⁸¹⁻⁹⁰.

Phytochemical	Bioactivity	Effect	Assay	Target	Mechanism of action
Coriander EOs (1.2%)	Antifungal	<i>In-vitro</i> MFC	Broth microdilution	<i>C. albicans</i>	Inhibites fungal growth (0.02 mg/mL)
Coriander EOs + conventional antibiotics	Antibacterial	<i>In-vitro</i> Improve antibiotics action MIC detrmiantion	Microdilution broth susceptibility assay	<i>A. baumannii</i>	Synergistic interaction (EOs+CHL, CIP, GEN, TET); Additive interaction (EOs + PIP, CFP)
Coriander EOs (β -linalool 66.07%; camphor 8.34%, geranylacetate 6.91%, cymene 6.35%)	Antibacterial	<i>In-vitro</i> Inhibition of cellular growth	Agar disc diffusion method	<i>B. subtilis</i> (10.69 \pm 0.47 mm) <i>Stenotrophomonas maltophilia</i> (9.22 \pm 0.08 mm)	Inhibites bacterial growth (inhibition halos)
Coriander SEOs (5 μ L/mL)	Antibacterial	<i>In-vitro</i> and <i>In-situ</i> MIC, MBC, effectiveness in carrots	Broth micro dilution method	<i>S. enterica</i>	Reduce and contain the growth of <i>Salmonella</i>
Coriander EOs (β -linalool 66.07% camphor 8.34%,	Antifungal	<i>In-situ</i> Antifungal analysis on bread	Agar microdilution method (MIC)	<i>P. expansum</i> MK-SF 33	Reduces mycelial growth and sporulation
Coriander SEOs (500 μ g/mL)	Antiviral	Antiviral activity by plaque assay *IC ₅₀ , SI	Plaque reduction assay (Vero cell line)	Herpes simplex Virus (HSV-1) 50 PFU	HSV-1 Inhibition (99%) *IC ₅₀ =350-250 μ g/mL
Coriander SEOs (1.2%)	Antibacterial	<i>In-vitro</i> MIC against Clinical pathogens	Broth microdilution method	Clinical pathogens <i>E. coli</i> (0.64mg/mL),	Inhibits bacterial growth (turbidity detection)
Coriander ESOs (4%)	Antibacterial	<i>In-vitro</i> MIC determination	Agar diffusion method	Clinical pathogen <i>S. aureus</i>	Inhibition of cell growth and death (inhibition 14.3 mm)
Coriander EOs (Linalool 73.5%, α -pinene 5.3%), γ -terpinene 4.5%)	Antiparasitic activity	<i>In-vitro</i> Anti-promastigote screening	Cytotoxic activity MTT assay	<i>L. amazonensis</i>	Inhibition against <i>L. amazonensis</i> IC ₅₀ =19.1 \pm 0.7 μ g/mL,
Coriander LEOs (40 mg/mL)	Anthelmintic activity	<i>In-vitro</i> Immersion assay	Paralysis/death elapsed time	<i>Pheretima posthuma</i>	paralysis (31 min) and death (78 min)
Coriander SEOs	Anthelmintic activity	<i>In-vitro</i> and <i>in-vivo</i> anthelmintic activity	White Balb/c mice	<i>Hymenolepis nana</i>	Anthelmintic agent (500-750 mg/kg B.W.)
Coriander HES	Antimicrobial Antibiofilm	Serial agar dilution (MIC); Biofilm	<i>Galleriamellone</i> lla L	<i>E. coli</i> , <i>S. enterica</i> , <i>P. aeruginosa</i> ,	Antimicrobial effect (MIC=1 mg/mL); <i>S. aureus</i> , <i>E. coli</i>
Coriander EOs (β -linalool 66.07% camphor 8.34%,	Antibiofilm formation	<i>In-vitro</i> Glass and wood surfaces	Agar microdilution method	<i>Bacillus subtilis</i> 7.42 μ L/mL <i>S. ltophilia</i>	Reduces biofilm formation (50% planktonic cells; 50% reduction in the cell
Coriander LEs and seeds extract	Antibacterial	<i>In-vitro</i> antibacterial screening	Disc-diffusion method	Human pathogenic <i>E. coli</i> ,	Inhibitory growth effect except against <i>P. aeruginosa</i>
Coriander EOs (linalool 70.93%, monoterpenes 77.34%)	Antibacterial	<i>In-vitro</i> Antimicrobial activity and viable count cells	Agar-well diffusion method	<i>E. coli</i> ATCC 8739 <i>Micrococcus luteus</i> ATCC 9341	Viable count cfu/mL (99.99 % killing)

EOs: Essential oils; SEOs: Seed Essential oils; LEs: Leaves extracts; HES: Hexane extract; MFC: Minimum Fungicidal Concentration; MIC: Minimum Inhibitory Concentration; MBC: Minimum bactericidal concentration; CHL: Chloramphenicol; CIP: Ciprofloxacin; CFP: Cefoperazone; GEN: Gentamicin; TET: Tetracycline; PIP: Piperacillin; PFU: Plaque-forming unit; MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; CLSM: Confocal laser scanning microscopy; SEM: Scanning electron microscopy; QS: Quorum sensing.

The phytochemical composition of coriander herb is similar to essential oil (EO) constituents, and additionally contains tannins, terpenoids, reducing sugars, alkaloids, phenols, flavonoids, fatty acids, sterols, and glycoside. The main components responsible for the characteristic odor of the herb are characterized by the presence of chemical groups such as various types of monoterpenes (oxide, alcohol, carbonyl, hydrocarbon, ester), aliphatic compounds (alcohol, aldehyde, hydrocarbon), phenolic compounds (phenols), sesquiterpenes, as well as some minor compounds such as acetic acid and α -p dimethyl styrene^{36,37}.

The essential oil of coriander is rich in phytochemicals, and the most recent quantitative analysis indicates the following compounds in mg/g: linalool (256.65) as main compound, α -pinene (19.85), geranyl acetate (15.76), γ -terpinene (12.08), camphor (8.51), p-cymene (6.10), geraniol (5.78), myrcene (3.25), limonene (2.90), sabinene (1.15), citronellal (0.77), borneol (0.76), geranial (0.60), neryl acetate (0.31), neral (0.28), eucalyptol (0.15), caryophyllene (0.06), thymol

(< 0.01), carvacrol (< 0.01), α -terpineol (< 0.01), nerol (< 0.01).

In recent years, advanced methods have been applied to extract new phytochemicals in plants such as coriander and its extracts. The heneicos-1-enes are new molecules isolated through chromatography³⁸. Furthermore, sophisticated nuclear magnetic resonance (NMR) techniques have enabled structural resolution with the support of high-resolution mass spectrometry (HRMS). Likewise, the increase in the extraction yield of the heneicos was increased using techniques like ionic liquid-based microwave-assisted extraction (IL-MAE) and the application of the response surface methodology³⁹. Microwave-assisted extraction based on ionic liquid is a fast and environmentally friendly extraction method applied to secondary metabolites in medicinal plants, in this case, coriander. Additionally, it is one of the unconventional extraction methods that has been developed and rapidly applied in recent years⁴⁰. In general, heneicos-1-ene belongs to the class of organic compounds known as unsaturated hydrocarbons; they are aliphatic hydrocarbons that

contain one or more unsaturated carbon atoms. These compounds, with a molecular formula of $C_{21}H_{42}$, contain one or more double or triple bonds and a molecular weight of 294.567 g/mol⁴¹.

The value of the bioactive compounds of *Coriandrum sativum* L treated in this manuscript are increased by molecules of heneicos-1-ene due to the outstanding potential to eradicate free radicals and proven antibacterial activity, as same as molecules isolated from coriander foliage, as confirmed by modern scientific studies⁴².

Extraction of bioactive compounds from plant extracts

Coriander oil is a natural compound, and its availability is extensive. Methods for oil extraction are very diverse and include both, traditional and modern techniques. Traditional and modern methods commonly applied for oil extraction, as well as the fundamental and general processes are briefly described in Table 2⁴²⁻⁵⁵.

Methods for the extraction of phytochemicals from coriander

The most popular methods for EOs extraction are still solvent extraction and steam distillation because they are relatively inexpensive, safe, and can provide a high-quality end product⁴⁶. After extraction, chemical identification and accurate quantification of the molecules contained in EOs are performed using Gas Chromatography coupled to Mass Spectrometry (GC-MS), High Resolution Gas Chromatography (GC-FID), and High-Performance Liquid Chromatography (HPLC) techniques⁵¹.

It is important to note that the success of EOs extraction depend on the type of method(s) and operating conditions selected for the process. Modern Green Extraction (MGE) methods have been reported to provide advantages in terms of performance compared to traditional methods (hydrodistillation) and improved selectivity compared to Soxhlet extraction⁵⁶.

To achieve the extraction of EOs from Coriander, traditional methods (hydrodistillation, solid-liquid extraction) were compared with modern methods (supercritical fluid extraction -SCFE- and subcritical water extraction -SCWE-). The comparison show yields of primary compounds (γ -(+)-limonene, terpinene, geraniol, camphor and linalool). Soxhlet extraction provides high proportions of oxygenated monoterpenes (camphor, geraniol, and linalool) in seeds, while the extraction yields of γ - terpinene and (+)-limonene were higher with MGE SFE, which contained high levels of camphor, geraniol and linalool⁴⁵.

The variations in EOs of coriander fruits depend not only on the extraction methods but also on the concentration of phytochemicals, which is directly related to the stage of growth or maturity, the season of herb harvest, and the prevailing climate and environmental conditions in the region of growth¹.

Regarding the growth of coriander, the quantity of Eos varies with the maturity stage of the herb, especially in the seeds⁴³. The linalool, the main component in coriander, increases as the herb grows, ranging from 36% during the first (flowering) stage to 78% during the last stage (maturity). These variations have also

been confirmed in crop yields in different regions of the world^{2,57}.

Outstanding biological activities of coriander

Antioxidant capacity of Coriander

During the metabolic processes of cells, biochemical reactions can generate free radicals, which are considered an oxidative process that needs to be controlled to avoid endangering cell function and integrity. Reactive oxygen species (ROS) can be produced, generating oxygen ions, free radicals and peroxides, reactive species derived from oxygen that cause oxidative stress. Reactive nitrogen species (RNS) can also be produced, such as nitric oxide, nitrogen dioxide, among others, causing nitrosative stress^{58,59}.

Free radicals can accumulate in cells and damage DNA, certain lipids and proteins, causing significant harmful effects on human health, and promoting the development of chronic degenerative diseases⁶⁰. This has generated enormous interest in developing scientific research, including epidemiological research, to counteract the effects of oxidative and nitrosative stress. Naturally, cells react by counteracting ROS through enzymatic mechanisms (catalase -CAT-, superoxide dismutase -SOD- glutathione peroxidase -GPX-) and non-enzymatic mechanisms (vitamins E and C, flavonoids, among others).

However, to help counteract the body's natural oxidation process, antioxidants can be acquired through diet. Vegetable consumption with being an important source of antioxidants such as polyphenols to neutralize free radicals^{58,61}. Polyphenols are capable of transferring electrons from the hydroxyl groups of their molecule, exerting an antioxidant action by reducing the free radicals in the environment⁶². Medicinal plants such as cilantro, in particular the extract of the oil from the seeds, are a source of potent antioxidant properties that have been investigated for their valuable biotherapeutic potential⁶³.

The antioxidant properties of cilantro are associated with or derived from polyphenolic and flavonoid compounds, and some others bioactive compounds. Especially the essential oil of coriander, linalool, has enormous antioxidant properties that neutralize the effect of lipid oxidation and/or lipid peroxidation⁶⁴.

Table 3 summarizes several representative studies of the potential and confirmed antioxidant capacity of coriander and/or some of its constituents⁶⁴⁻⁷⁴.

The high natural antioxidant properties of coriander constituents in EOs extract are widely used in the food industry protection, as substitutes for dangerous synthetic/chemical antioxidants. These chemical antioxidants traditionally including butylhydroxy-anisole, butylhydroxytoluene, tertbutyl hydroxyl-quinone and propyl gallate⁶⁵.

Likewise, the table shows the activity exerted and the assessments carried out to determine its effect by any of the action mechanisms (Radical scavenging activity, Ferric reducing ability of plasma, Nitric oxide radical scavenging, among others).

Antimicrobial activity derived from Coriander

Currently, pharmaceuticals, mostly synthetic drugs, are the dominant therapeutic agents in modern medicine. However, the phytomedicinal drugs derived from

plants, are becoming more popular as a potential alternative strategy against antimicrobial resistance. In this regard, the WHO reports clearly show that more and more countries are recognizing the role of traditional and complementary medicine in their national health systems. The WHO strives to implement and follow up the WHO Traditional Medicine Strategy 2014-2023 worldwide.

Coriander is a globally important crop used as a vegetable, spice, fragrance, and traditional medicine⁵⁷. Also, coriander has been used since ancient times for its ability to control various organisms due to its recognized antibacterial activity. In general, antimicrobial activity refers to the process of killing or inhibiting disease-causing micro-organisms, mostly bacteria. Many antimicrobial agents are used to reduce or eliminate microbial growth by different modes of action and mechanisms. Anti-microbials can be antibacterial, antifungal, or antiviral, and they all have different modes of action by which they act to suppress the infection. The antimicrobials act on specific targets identified as nucleic acids synthesis, cell wall synthesis (peptidoglycan), and structural and/or functional damage to the cell membrane by depolarization, disruption or permeability⁷⁵.

Human mycosis is one of the most important health problems worldwide. However, abundant research in the field of antifungal drug fungal treatment research slowly than bacterial infections control. The common antifungal activity may include the inhibition of ergosterol synthesis by lanosterol 14- α demethylase activity in fungal endoplasmic reticulum (azoles). Also, antifungals binding to ergosterol causing disruption of the fungal cell membrane (polyenes), and the inhibition of (1,3) β -D-glucan synthase stops glucan synthesis (echinocandins). The fungistatic nature of many popular antifungals and the emergence of resistance limit their success, and new approaches to fungal treatment are being sought.

These new approaches include phytochemical screening in medicinal plants to isolate bioactive compounds as therapeutic tools against fungal diseases. In this regard, linalool has shown *in vitro* antifungal activity against *Candida albicans*, and inhibits activity in 1, 3 beta-glucan synthase, lanosterol 14 alfa-demethylase, and delta 14-esterol reductase, damaging the cell wall and plasmatic membrane in fungi⁷⁵⁻⁷⁷. Also, both clinical investigations and *in vivo* experiments have confirmed the antifungal activity of flavonoids^{78,79}. The efficacy of linalool against *L. monocytogenes* in the planktonic stage and biofilm formation has been explored, as well as its putative mechanism of action. The evidence shown strong anti-listeria activity in the planktonic stage by linalool⁸⁰.

Table 4 summarizes several representative studies of the potential and confirmed antimicrobial capacity of coriander and/or some of its constituents^{81,82-90}.

Coriander's multiple health benefits

In recent years, coriander and its phytochemical components have been studied from a medical perspective primarily for their great potential to promote health as curative agents for therapeutic management of many diseases. As mentioned above,

phytochemical content of coriander is linked to its biological functions, so high expectations for the use of coriander in biotherapeutic treatments against chronic and degenerative diseases¹⁰. Chronic diseases, both in developing and developed countries, encompass a wide range of medical conditions, including cancer, heart disease, Alzheimer's disease, diabetes, kidney disease, and cerebrovascular diseases, among others.

In this context, the World Health Organization (WHO) reports cancer as the second leading cause of death worldwide, with an estimated 9.6 million deaths, or one in six deaths, in 2018⁶⁵. Additionally, statistically, the American Cancer Association estimated 1.8 million new cancer cases in 2020, unfortunately resulting in approximately 606,520 cancer-related deaths in the United States. In 2024, 2,001,140 new cancer cases and 611,720 cancer deaths are projected to occur in this country⁹¹. The *anticancer* effect of coriander has been confirmed, specifically by affecting the survival of the cells and triggering apoptosis, thereby impacting the invasive and colony formation properties of different cell lines, such as prostate cancer⁹², breast cancer⁹³, human colon cancer, and ovarian carcinoma⁹⁴.

Heart and cardiovascular diseases (HCVDs) are the leading cause of death globally according to the WHO, with an estimated 17.9 million deaths each year⁹⁵. The HCVDs include heart disorders of the heart and blood vessels and encompass coronary heart disease, cerebrovascular disease, rheumatic heart disease and other conditions. In the treatment of heart disease, coriander extract can lower blood pressure as well as reduce other heart disease risk factors such as bad cholesterol levels⁹⁶. Also, coriander seeds can create significant decrease low-density lipoproteins (LDL) and increase high-density lipoprotein (HDL). The above indicates a previously reported hypolipidemic effect and decreases in cholesterol synthesis. Cholesterol significantly increases the risk of developing cardiovascular heart diseases and mortality⁹⁷.

The treatment of chronic diseases such as *Alzheimer's* is very varied and, at the same time, very exhausting for patients and their families. In recent years, physicians have sought natural alternatives for biotherapeutic treatment, and coriander leads a very promising application. For example, the inhalation of coriander volatile oil reduces oxidative stress in a beta-amyloid (1-42) rat model of Alzheimer's disease⁹⁸.

Nowadays, diabetes is a chronic (long-lasting) disease that affects the pathways used by the human body to metabolize food into energy. There are three main types of diabetes: type 1, type 2, and gestational diabetes (diabetes while pregnant). More than 122 million Americans are living with diabetes (34.2 million) or prediabetes (88 million)⁷⁵. Many patients with diabetes mellitus consider medicinal plants such as coriander as complementary treatment for this common chronic and metabolic disease and its derivative complications such as impaired wound healing⁹⁹.

Coriander may stimulate endocrine glands and pancreatic insulin secretion to increase the level of insulin in the blood. In that way, coriander regulates

the absorption and assimilation of sugar, resulting in a drop in blood sugar levels¹⁰⁰. Coriander helps to reduce the risk of risky fluctuations in blood sugar limits, and also promotes metabolic functions. The management of diabetes by effect of coriander can attenuate lipids, glucose and creatinine levels and decrease lipid peroxidation. In addition, it also increases the activity of antioxidant enzymes in patients^{101,102}.

Neurodegeneration leading to cognitive deficits and a lack of cognitive capability in humans, for which there is no cure or preventive treatment currently, can potentially be addressed through traditional. The essential oils of coriander contain a great amount of linalool, a phytochemical with bioactivity against the neurotoxic effects of beta amyloid 1-42 (A β 1-42), which is mainly responsible for the functional loss in neuronal cells such as cell viability, synaptic function and neuronal network activity¹⁰³. Regarding older adults and the promotion of the quality of life to reach independence for people as they age, the coriander has significant value¹⁰⁴.

CONCLUSIONS

Coriander is phytochemical-rich herbaceous, and maybe the most popular and economically valuable spice worldwide. The recent rise in popularity is due to their multiple health benefits linked to the phytochemicals/bioactive compounds with a broad spectrum regarding to phytochemical and biological activities. These activities include antimicrobial and antioxidant properties to develop anticancer, anti-inflammatory, hypoglycemic, hypolipidemic, hypotensive among others. Undoubtedly, the coriander is excellent source of pharmaceuticals, and could be efficiently extracted and studied *in vitro* or *in vivo* to be applied according to their phytochemical diversity, as well as, by the wide variety of beneficial effects exerted on nutrition and human health to well-being promotion and enhance the healthy aging conditions. Multiple scientific researches about plethora of bioactivities from coriander, their medical applications to promote the well-being and enhance the healthy aging, undoubtedly could be getting closer to benefiting the world's population health.

ACKNOWLEDGEMENTS

None to declare.

AUTHOR'S CONTRIBUTION

De la Fuente-Salcido NM: writing original draft, investigation, analysis of data. **Lafuente-Rincón DF:** design and conceptualization of tables. Final manuscript was approved by both authors.

DATA AVAILABILITY

Upon request, the accompanying author can furnish the empirical data used to bolster the findings of the study.

CONFLICT OF INTEREST

None to declare

REFERENCES

1. Iqbal M, Butt MS, Suleria HAR. Coriander (*Coriandrum sativum* L.): Bioactive molecules and health effects. In Mérillon JM, Ramawat K editors. Bioactive molecules in food. Reference Series in Phytochemistry. Springer, Champ; 2019; 14-21.
2. National Nutrient data base USDA. <https://data.nal.usda.gov/dataset/usda-national-nutrient-database-standard-reference-legacy-release> Accessed May 2024.
3. Mandal S, Mandal M. Coriander (*Coriandrum sativum* L.) Essential oil: Chemistry and biological activity. 2015; Asian Pac J Trop Biomed 2015; 5(6):421-428. <https://doi.org/10.1016/j.apjtb.2015.04.001>
4. Chahal K, Singh R, Kumar A, Bhardwaj U. Chemical composition and biological activity of *Coriandrum sativum* L: A review. J Nat Prod Resour 2017; 8(3):193-203. <https://10.56042/ijnpr.v8i3.1313>
5. Isah T. Stress and defense responses in plant secondary metabolites production. Biol Res 2019; 52 (39):1-25. <https://doi.org/10.1186/s40659-019-0246-3>
6. Jiménez-Gómez A, García-Estévez I, García-Fraile P, Escribano-Bailón MT, Rivas R. Increase in phenolic compounds of *Coriandrum sativum* L. after the application of a *Bacillus halotolerans* biofertilizer. Sci Food Agric 2020; 100(6):2742-2749. <https://doi.org/10.1002/jsfa.10306>
7. Nguyen DTP, Lu N, Kagawa N, Takagaki M. Optimization of photosynthetic photon flux density and root-zone temperature for enhancing secondary metabolite accumulation and production of Coriander in plant factory. Agronomy 2019; 9(5):224. <https://doi.org/10.3390/agronomy9050224>
8. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: extraction, isolation, and identification of bioactive compounds from plant extracts. Plants (Basel) 2017; 6(4):42. <https://doi.org/10.3390/plants6040042>
9. Li Z, Lee HW, Liang X, Liang D, Wang Q, Huang D, Ong CN. Profiling of phenolic compounds and antioxidant activity of 12 Cruciferous vegetables. Molecules 2018; 23(5):1139. <https://doi.org/10.1104/pp.20.01185>
10. Prachayasittikul V, Prachayasittikul S, Ruchirawat S, Prachayasittikul V. Coriander (*Coriandrum sativum*): A promising functional food toward the well-being. Food Res Int 2018; 105:305-323. <https://doi.org/10.1016/j.foodres.2017.11.019>
11. Papuc C, Goran GV, Predescu CN, Nicorescu V, Stefan G. Plant polyphenols as antioxidant and antibacterial agents for shelf-life extension of meat and meat products: Classification, structures, sources, and action mechanisms. Compr Rev Food Sci 2017; 6(6):1243-1268. <https://doi.org/10.1111/1541-4337.12298>
12. Ismail F, Ismail WU, Muhammad G, Hussain MA, Zulfiqar Z. Phytochemistry, pharmacological attributes, and clinical evaluations of *Coriandrum sativum*: A comprehensive review. Natural Prod Commun 2025; 20(1). <https://doi.org/10.1177/1934578X241312791>
13. Asfaw MD. A Review on the chemical constituent, pharmacological and medicinal properties of *Coriandrum sativum*. Nat Prod Chem Res 2021; 9(8):419. <http://10.9790/3013-067031742>
14. Manville RW, Abbott GW. Cilantro leaf harbors a potent potassium channel-activating anticonvulsant. FASEB J 2019; 33(10):11349-11363. <https://doi.org/10.1096/fj.201900485R>

15. Önder A. Coriander and its phytoconstituents for the beneficial effects. In: El-Shemy HA editors. Intech Open 2018. <http://doi.org/10.5772/intechopen.78656>
16. Laribi B, Kouki K, M'Hamdi M, Bettaieb T. Coriander (*Coriandrum sativum* L.) and its bioactive constituents. Fitoterapia 2015; 103: 9-26. <https://doi.org/10.1016/j.fitote.2015.03.012>
17. Mahleyuddin NN, Moshawih S, Ming LC, *et al.* *Coriandrum sativum* L.: A review on ethnopharmacology, phytochemistry, and cardiovascular benefits. Molecules 2021; 27(1):209. <https://doi.org/10.3390/molecules27010209>
18. Tang Y, Huang Y, Zhang B, Luo T and Zhong W. Food rich in phenolic compounds and their potential to fight obesity. Front Nutr 2023; 10: 1204981. <https://doi.org/10.3389/fnut.2023.1204981>
19. Abbas A, Anwar F, Ahmad N, *et al.* Characterization of bioactives and nutra-pharmaceutical potential of supercritical fluid and hydro-distilled extracted Coriander leaves essential oil. Dose Response 2022; 20(4):15593258221130749. <https://doi.org/doi:10.1177/15593258221130749>
20. Ćimović MG, Kostadinović LM, Puvača NM, Popović SJ, Urošević MI. Phytochemical constituents of selected plants from Apiaceae family and their biological effects in poultry. Food Feed Res 2019; 43(1):35-41. <https://doi.org/doi:10.5937/FFR1601035A>
21. Gómez-Gómez ME, Zapico SC. Frailty, Cognitive Decline, Neurodegenerative Diseases and Nutrition Interventions. Int J Mol Sci 2019; 20(11):2842. <https://doi.org/doi:10.3390/ijms20112842>
22. Mc Ausland L, Lim M-T, Morris DE, Smith-Herman HL, Mohammed U, Hayes-Gill BR, Crowe JA, Fisk ID and Murchie EH. Growth spectrum complexity dictates aromatic intensity in Coriander (*Coriandrum sativum* L.). Front Plant Sci 2020; 11:462. <https://doi.org/doi:10.3389/fpls.2020.00462>
23. Ahmad NI, Rahman SA, Leong Y-H, Azizul NH. A Review on the phytochemicals of Parkia Speciosa, stinky beans as potential phytomedicine. J Food Sci Nutr Res 2019; 2 (3): 151-173. <https://doi.org/doi:10.26502/jfsnr.2642-11000017>
24. Park K. The Role of Dietary Phytochemicals: Evidence from epidemiological studies. Nutrients 2023; 15 (6): 1371. <https://doi.org/10.3390/nu15061371>
25. Wei JN, Liu ZH, Zhao YP, Zhao LL, Xue TK, Lan QK. Phytochemical and bioactive profile of *Coriandrum sativum* L. Food Chem 2019; 286: 260-267. <https://doi.org/10.1016/j.foodchem.2019.01.171>
26. Azemi AK, Nordin ML, Hambali KA, *et al.* Phytochemical contents and pharmacological potential of *Parkia speciosa* Hassk for diabetic vasculopathy: A Review. Antioxidants 2022; 11(2):431. <http://doi:10.3390/antiox11020431>
27. Al-Khayri JM, Banadka A, Nandhini M, *et al.* Essential Oil from *Coriandrum sativum*: A review on its phytochemistry and biological activity. Molecules 2023; 28(2):692. <http://doi:10.3390/molecules28020696>
28. Tylewicz U, Nowacka M, Martín-García B, Wiktor A, Gómez Caravaca AM. 5-Target sources of polyphenols in different food products and their processing by-products. In: C.M. Galankis editors. Polyphenols: Properties, Recovery, and Applications. Elsevier Inc 2018; 135-175.
29. Nathaniel S, Fatima A, Fatima R, *et al.* Phytochemical study of acetone solvent extract of Coriander sativum. J Pharmacogn Phytochem 2019; 8(6): 186-140.
30. Güneş Bayır A, Güney M. Nutritional assessment and use of complementary and alternative medicine in cancer patients treated with radiotherapy and chemotherapy. Altern Ther Health Med 2019; 25(6):28-33.
31. Gantait S, Sharangi AB, Mahanta M, Meena NK. Agri-biotechnology of coriander (*Coriandrum sativum* L.): an inclusive appraisal. Appl Microbiol Biotechnol 2022; 106(3): 951-969. <https://doi.org/10.1007/s00253-022-11787-4>
32. El-Zaedi H, Calin-Sánchez A, Nowicka Pet *et al.* Preharvest treatments with malic, oxalic, and acetylsalicylic acids affect the phenolic composition and antioxidant capacity of coriander, dill and parsley. Food Chem 2017; 226:179-186. <https://doi.org/10.1016/j.foodchem.2017.01.067>
33. Bhardwaj K, Islam MT, Jayasena V, *et al.* Review on essential oils, chemical composition, extraction, and utilization of some conifers in Northwestern Himalayas. Phytother Res 2020; 34(11):2889-2910. <https://doi.org/10.1002/ptr.6736>
34. Satyal P, Setzer WN. Chemical compositions of commercial essential oils from *Coriandrum sativum* fruits and aerial parts. Nat Prod Commun 2020; 15(7):1-12. <https://doi.org/10.1177/1934578X20933067>
35. Rodriguez-Garcia C, Sánchez-Quesada CJ, Gaforio J. Dietary Flavonoids as Cancer Chemopreventive agents: An updated review of human studies. Antioxidants 2019; 8(5):137. <https://doi.org/doi:10.3390/antiox8050137>
36. Al-Snafi AE. A Review on chemical constituents and pharmacological activities of *Coriandrum sativum*. IOSR J Pharm 2016; 6(7):17-42. <https://doi.org/10.9790/3013-067031742>
37. Micić D, Ostojić S, Pezo L, Blago S, Pavličić B, Zeković Z, Đurović S. Essential oils of coriander and sage: Investigation of chemical profile, thermal properties and QSRR analysis. Ind Crops Prod 2019; 138:111438. <https://doi.org/10.1016/j.indcrop.2019.06.001>
38. Priyadarshi S, Harohally NV, Roopavathi C, Naidu MM. Isolation, identification, structural elucidation and bioactivity of Heneicos-1-ene from *Coriandrum sativum* L. Foliage Sci Rep 2018; 8:17414. <https://doi.org/10.1038/s41598-018-35836-z>
39. Priyadarshi S, Balaraman M, Naidu MM. Ionic liquid-based microwave-assisted extraction of Heneicos-1-ene from coriander foliage and optimizing yield parameters by response surface methodology. Prep Biochem Biotechnol 2020; 50(3):246-251. <https://doi.org/10.1080/10826068.2019.1687519>
40. Ahmad I, Yanuar A, Mulia K, Mun'im A. Ionic liquid-based microwave-assisted extraction: Fast and green extraction method of secondary metabolites on medicinal plant. Pharmacogn Rev 2018; 12(23): 20-26. https://doi.org/10.4103/phrev.phrev_40_17
41. Hosen SMZ, Junaid M, Alam MS, *et al.* GreenMolBD: Nature Derived Bioactive Molecules Database. Med Chem 2022; 18 (6):724-733. <https://doi.org/10.2174/1573406418666211129103458>
42. Goncales R de Melo AC, dos Santos M DV, Carvalho Neto MF, *et al.* Phytochemical trial and bioactivity of the essential oil from coriander leaves (*Coriandrum sativum*) on pathogenic microorganisms. Chem Eng Trans 2019; 75:403-408. <https://doi.org/10.3303/CET1975068>
43. Msaada K, Taarit MB, Hosni K, *et al.* Comparison of different extraction methods for the determination of essential oils and related compounds from Coriander (*Coriandrum sativum* L.). Acta Chim Slov 2012; 59 (4):803-813.
44. Abbassi A, Mahmoudi H, Zaouali W, *et al.* Enzyme-aided release of bioactive compounds from coriander (*Coriandrum sativum* L.) seeds and their residue by-products and evaluation of their antioxidant activity. J Food Sci Technol 2018; 55(8):3065-3076. <https://doi.org/10.1007/s13197-018-3229-4>
45. Pavlic B, Vidovic S, Vladoic J, Radosavljevic R, Zekovic Z. Isolation of coriander (*Coriandrum sativum* L.) essential oil by green extractions versus traditional techniques. J Supercrit Fluids 2015;99: 23-28. <http://dx.doi.org/10.1016/j.supflu.2015.01.029>
46. Gonzalez-Marrugo LB, Granados-Llamas EA, Granados-Conde C, Tejada-Tovar CN, Ortega-Toro R. Extraction and evaluation of the antioxidant properties of Coriander (*Coriandrum sativum*) seed essential oil. Contemp Eng Sci 2018; 11:3841-3848.

- <https://doi.org/10.12988/ces.2018.87356>
47. Anita P, Sivasamy S, Madan Kumar PD, Balan IN, Ethiraj S. *In vitro* antibacterial activity of *Camellia sinensis* extract against cariogenic microorganisms. *J Basic Clin Pharm* 2014; 6 (1): 35-39.
<https://doi.org/10.4103/0976-0105.145777>
 48. Zeković Z, Adamović D, Četković G, Radojković M, Vidović S. Essential oil and extract of coriander (*Coriandrum sativum* L.). *Acta Period Technol* 2011; 42:281-288. <https://doi.org/10.2298/APT1142281Z>
 49. Üstün Argon Z, Gökkyer A, Gümüş ZP, Büyükhelvacıgil M. Evaluation of some medicinal herbs cold pressed oils according their physicochemical properties with chemometry. *Int J Sec Metabolite* 2017;(3, Special Issue 2), 473-481. <https://doi.org/10.21448/ijsm.37731950>
 50. Malysa-Pasko M, Lukaszewicz M, Jakubowski P. Microwave-assisted extraction of bioactive compounds from seeds of milk thistle, black cumin and coriander. The 19th International Electronic Conference on Synthetic Organic Chemistry 2015;
<https://doi.org/10.3390/ecsoc-19-b005>
 51. Shrirame BS, Geed SR, Raj A, *et al.* Optimization of supercritical extraction of Coriander (*Coriandrum sativum* L.) seed and characterization of essential ingredients. *J Essent Oil-Bear Plants* 2018; 2:330-344.
<https://doi.org/10.1080/0972060X.2018.1470943>
 52. Mejri J, Aydi A, Abderrabba M, Mejri M. Emerging extraction processes of essential oils: a review. *Asian J Green Chem* 2018; 2 (3): 246-267.
<https://doi.org/10.22034/ajgc.2018.61443>
 53. Sourmaghi MH, Kiaee G, Golfakhrabadi F, *et al.* Comparison of essential oil composition and antimicrobial activity of *Coriandrum sativum* L. extracted by hydrodistillation and microwave-assisted hydrodistillation. *J Food Sci Technol* 2015; 52(4): 2452-2457.
<https://doi.org/10.1007/s13197-014-1286-x>
 54. Asofiei I, Călinescu I, Gavrilă AI, Ighigeanu D, Martin D, Matei C. Microwave hydrodiffusion and gravity, a green method for the essential oil extraction from ginger - energy considerations. *U.P.B. Sci Bull* 2017; 79:81-92. ISSN 1454-2331
 55. Hii CL, Ong SP, Chiang CL, Menon AS. A review of quality characteristics of solar dried food crop product. *IOP Conf Ser: Earth Environ Sci* 2019; 292.
<https://doi.org/10.1088/1755-1315/292/1/012054>
 56. Pavlić B, Bera O, Teslić N, Vidović S, Parpinello G, Zeković Z. Chemical profile and antioxidant activity of sage herbal dust extracts obtained by supercritical fluid extraction. *Ind Crops Prod* 2018; 120:305-312.
<https://doi.org/10.1016/j.indcrop.2018.04.044>
 57. Assefa M, Yaya EE, Chandravanshi BS, Assefa M. Fatty acid and essential oil compositions of seeds of coriander (*Coriandrum sativum* L.) cultivated in different regions of Ethiopia. *Bull Chem Soc Ethiop* 2024; 38 (4): 863-876.
<https://dx.doi.org/10.4314/bcse.v38i4.4>
 58. Pizzino G, Irrera N, Cucinotta M, *et al.* Oxidative stress: harms and benefits for human health. *Oxid Med Cell Longev* 2017; 8416763.
<https://doi.org/10.1155/2017/8416763>
 59. García-Sánchez A, Miranda-Díaz AG, Cardona-Muñoz EG. The role of oxidative stress in physiopathology and pharmacological treatment with pro- and antioxidant properties in chronic diseases. *Oxid Med Cell Longev* 2020; 2082145.
<https://doi.org/10.1155/2020/2082145>
 60. WHO World Health Organization 2020. Noncommunicable diseases https://www.who.int/health-topics/noncommunicable-diseases#tab=tab_1 Accessed 15 Jan 2024.
 61. Xu DP, Li Y, Meng X, Zhou T, Zhou Y, Zheng J, Zhang JJ, Li HB. Natural antioxidants in foods and medicinal plants: extraction, assessment and resources. *Int J Mol Sci* 2017; 18 (1): 96. <https://doi.org/10.3390/ijms18010096>
 62. Mohotti S, Rajendran S, Muhammad T, *et al.* Screening for bioactive secondary metabolites in Sri Lankan medicinal plants by microfractionation and targeted isolation of antimicrobial flavonoids from *Derris scandens*. *J Ethnopharmacol* 2019; 246:112148.
<https://doi.org/10.1016/j.jep.2019.112158>
 63. Vasconcelos Dos Santos MD, De Carvalho Neto MF, Gonçalves Reis De Melo AC, *et al.* Chemical Composition of Essential Oil of Coriander Seeds (*Coriandrum sativum*) Cultivated in the Amazon Savannah, Brazil. *Chem Eng Trans* 2019; 75:409-414.
<https://doi.org/10.3303/CET1975069>
 64. Kačaniová M, Galovičová L, Ivanišová E, *et al.* Antioxidant, antimicrobial and antibiofilm activity of coriander (*Coriandrum sativum* L.) essential oil for its application in foods. *Foods* 2020; 9 (3): 282.
<https://doi.org/10.3390/foods9030282>
 65. Kebede M, Admassu S. Application of antioxidants in food processing industry: options to improve the extraction yields and market value of natural products. *Adv Food Technol Nutr Sci Open J* 2019.
<https://api.semanticscholar.org/CorpusID:78088380>
 66. Jeya KR, Veerapagu M, Sangeetha V. Antimicrobial and antioxidant properties of *Coriandrum sativum* L. seed essential oil. *Am J Essent Oil Nat Prod* 2019; 7 (2): 06-10.
 67. Crespo YA, Bravo Sánchez LR, Quintana YG, *et al.* Evaluation of the synergistic effects of antioxidant activity on mixtures of the essential oil from *Apium graveolens* L., *Thymus vulgaris* L. and *Coriandrum sativum* L. using simplex-lattice design. *Heliyon* 2019; 15(6):e01942.
<https://doi.org/10.1016/j.heliyon.2019.e01942>
 68. Senrayan J, Venkatachalam S. Solvent-assisted extraction of oil from papaya (*Carica papaya* L.) seeds: Evaluation of its physicochemical properties and fatty-acid composition. *Sep Sci Technol* 2018; 53 (3):1-8.
<https://doi.org/10.1080/01496395.2018.1480632>
 69. Baliyan S, Mukherjee R, Priyadarshini A, *et al.* Determination of antioxidants by DPPH radical scavenging activity and quantitative phytochemical analysis of *Ficus religiosa*. *Molecules* 2022; 27(4):1326.
<https://doi.org/10.3390/molecules27041326>
 70. Afsheen N, Ur-Rehman K, Jahan N, *et al.* Cardioprotective and metabolomic profiling of selected medicinal plants against oxidative stress. *Oxid Med Cell Longev* 2018; 9819360. <https://doi.org/10.1155/2018/9819360>
 71. Miri YB, Djenane D. Evaluation of protective impact of Algerian *Cuminum cyminum* L. and *Coriandrum sativum* L. essential oils on *Aspergillus flavus* growth and aflatoxin b1 production. *Pak J Biol Sci* 2018; 21 (2): 67-77.
<https://doi.org/10.3923/pjbs.2018.67.77>
 72. Bag A, Chattopadhyay RR. Evaluation of antioxidant potential of essential oils of some commonly used Indian spices in *in vitro* models and in food supplements enriched with omega-6 and omega-3 fatty acids. *Research in Environ Sci Pollut* 2018; 25(1):388-398.
<https://doi.org/10.1007/s11356-017-0420-5>
 73. Anusha MB, Shivanna N, Kumar GP, Anilakumar KR. Efficiency of selected food ingredients on protein efficiency ratio, glycemic index, and *in vitro* digestive properties. *J Food Sci Tech* 2018; 55(5):1913-1921.
<https://doi.org/10.1007/s13197-018-3109-y>
 74. Khan IU, Verma P, Rathore BS, Verma JR, Sharma LK. Genetic variability for essential oil, polyphenols and antioxidant activity of coriander (*Coriandrum sativum* L.) genotypes grown in humid south eastern plain zone V of Rajasthan. *Int J Chem Stud* 2017; 5(5):2292-2297.
 75. Medeiros CIS, Sousa MNA, Filho GGA, *et al.* Antifungal activity of linalool against fluconazole-resistant clinical strains of vulvovaginal *Candida albicans* and its predictive mechanism of action. *Braz J Med Biol Res* 2022; 15(55):e11831.
<http://doi:10.1590/1414-431X2022e11831>
 76. de Andrade Monteiro C, de Santos JRA. Phytochemicals and their antifungal potential against pathogenic yeasts. In:

- V Rao, D Mans, L Rao editors. Phytochemicals in Human Health. Intech Open 2019.
<https://doi.org/10.5772/intechopen.87302>
77. Aboody MSA, Mickymaray S. Anti-fungal efficacy and mechanisms of flavonoids. *Antibiotics* 2020;9 (2):45.
<https://doi.org/10.3390/antibiotics9020045>
78. Mickymaray S. Efficacy and mechanism of traditional medicinal plants and bioactive compounds against clinically important pathogens. *Antibiotics* 2019; 8(4):257.
<https://doi.org/10.3390/antibiotics8040257>
79. Sulaiman M, Nissapatorn V, Rahmatullah M, *et al.* Antimicrobial secondary metabolites from the Mangrove Plants of Asia and the Pacific. *Mar Drugs* 2022; 20 (10): 643. <https://doi.org/10.3390/md20100643>
80. Gao Z, Van Nostrand JD, Zhou J, Zhong W, Chen K, Guo J. Anti-listeria activities of linalool and its mechanism revealed by comparative transcriptome analysis. *Front Microbiol* 2019; 10:2947.
<https://doi.org/10.3389/fmicb.2019.02947>
81. Duarte A, Ferreira S, Silva F, Domingues FC. Synergistic activity of coriander oil and conventional antibiotics against *Acinetobacter baumannii*. *Phytomedicine* 2012; 19(3-4): 236-238.
<https://doi.org/10.1016/j.phymed.2011.11.010>
82. Pellegrini M, Rossi C, Palmieri S, *et al.* Salmonella enterica control in stick carrots through incorporation of coriander seeds essential oil in sustainable washing treatments. *Front sustain Food Syst* 2020; 4: 14.
<https://doi.org/10.3389/fsufs.2020.00014>
83. Fayyad AG, Nazlina I, Wan AY. Evaluation of biological activities of seeds of *Coriandrum sativum*. *Int J Eng Res* 2017; 8(7):1058-1063.
84. Hamudeng AM, Serliawati S. Effectiveness of antibacterial extract of coriander seeds (*Coriandrum sativum* L.) against *Staphylococcus aureus*. *J Dentomaxillofac Sci* 2019; 4(2):71.
<https://doi.org/10.15562/jdmfs.v4i2.840>
85. Monzote L, Herrera I, Satyal P, Setzer WN. *In-vitro* evaluation of 52 commercially-available essential oils against *Leishmania amazonensis*. *Molecules* 2019; 24(7):1248. <https://doi.org/10.3390/molecules24071248>
86. Nair SN, Neeraja ED, Rejimon G, Meenu B, Varghese A. Comparative evaluation of the anthelmintic activity of *Coriandrum sativum* Linn. and *Apium graveolens* Linn. *Res J Pharm Tech* 2017; 10(11): 3857-3859.
<https://doi.org/10.5958/0974-360X.2017.00699.0>
87. Hosseinzadeh S, Ghalesefidi MJ, Azami M, *et al.* *In vitro* and *in vivo* anthelmintic activity of seed extract of *Coriandrum sativum* compared to Niclosamid against *Hymenolepis nana*. *J Parasit Dis* 2016; 40(4):1307-1310.
<https://doi.org/10.1007/s12639-015-0676>
88. Molina RDI, Campos-Silva R, Macedo AJ, Blázquez AM, Rosa MEA. Antibiofilm activity of coriander (*Coriander sativum* L.) grown in Argentina against food contaminants and human pathogenic bacteria. *Ind Crops Prod* 2020; 151:1-9. <https://doi.org/10.1016/j.indcrop.2020.112380>
89. Verma A, Agarwal D, Dhanik J, Arya N, Nand V. Antibacterial efficacy and cluster analysis of genotypic extracts of Coriander leaves and seeds against human pathogenic bacterial strains. *Int J Curr Microbiol Appl Sci* 2018; 7(5):2727-2736.
<https://doi.org/10.20546/ijcmas.2018.705.316>
90. Kahil, N, Ashour, Fikry S, Naser A. Chemical composition and antimicrobial activity of the essential oils of selected Apiaceous fruits. *Future J Pharm Sci* 2018; 4(1):88-92.
<https://doi.org/10.1016/j.fjps.2017.10.004>
91. American Cancer Society.
<https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/2024> Accessed 20 June 2024.
92. Elmas L, Secme M, Mammadov R, Fahrioglu U, Dodurga Y. The determination of the potential anticancer effects of *Coriandrum sativum* in PC-3 and LNCaP prostate cancer cell lines. *J Cell Biochem* 2019; 120(3):3506-3513.
<http://doi.org/10.1002/jcb.27625>
93. Tang EL, Rajarajeswaran J, Fung SY, Kanthimathi MS. Antioxidant activity of *Coriandrum sativum* and protection against DNA damage and cancer cell migration. *BMC Complement Altern Med* 2013; 13:347.
<https://doi.org/10.1186/1472-6882-13-347>
94. Iwasaki K, Zheng YW, Murata Set *al.* Anticancer effect of linalool via cancer-specific hydroxyl radical generation in human colon cancer. *World J Gastroenterol* 2016; 22 (44): 9765-9774. <https://doi.org/10.3748/wjg.v22.i44.9765>
95. World Health Organization 2019. WHO global report on traditional and complementary medicine 2019.
<https://apps.who.int/iris/handle/10665/312342>
96. Jain S, Buttar HS, Chintameneni M, Kaur G. Prevention of cardiovascular diseases with anti-inflammatory and antioxidant nutraceuticals and herbal products: An overview of pre-clinical and clinical studies. *Recent Pat Inflamm Allergy Drug Discov* 2018; 12(2):145-157.
<http://10.2174/1872213X12666180815144803>
97. Das S, Rajadnya V, Kothari R, Tilak AV, Raveendran S, Deshpande T. Hypolipidemic activity of *Coriandrum sativum* in diabetic dyslipidemic rats. *Int J Basic Clin Pharmacol* 2019; 8(6):1393-1397.
<https://doi.org/10.18203/2319-2003.ijbcp20192208>
98. Hosseini M, Boskabady MH, Khazdair MR. Neuroprotective effects of *Coriandrum sativum* and its constituent, linalool: A review. *Avicenna J Phytomed* 2021; 11(5):436-450. <http://doi.org/10.22038/AJP.2021>
99. Mechchate H, Es-Safi I, Amaghounje A, *et al.* Antioxidant, anti-inflammatory and antidiabetic proprieties of LC-MS/MS identified polyphenols from Coriander seeds. *Molecules* 2021; 26(2):487.
<http://doi.org/10.3390/molecules26020487>
100. Aligita W, Susilawati E, Septiani H, Atsil R. Antidiabetic activity of Coriander (*Coriandrum sativum* L) leaves' ethanolic extract. *J Pharm Phytopharmacological Res* 2018; 8(2):59-63. <https://ejppr.com/g0YW8P3>
101. Zamany S, Malek A, Tabriz M, Pirouzpanah S, Barzegar. The effects of coriander seed supplementation on serum glycemic indices, lipid profile and parameters of oxidative stress in patients with type 2 diabetes mellitus: A randomized double-blind placebo-controlled clinical trial. *Res Square* 2022.
<https://doi.org/10.21203/rs.3.rs-262149/v2>
102. Kajal A, Singh R. *Coriandrum sativum* seeds extract mitigate progression of diabetic nephropathy in experimental rats via AGEs inhibition. *PLoS ONE* 2019; 14(3): e0213147.
<http://doi.org/10.1371/journal.pone.0213147>
103. Caputo L, Piccialli I, Ciccone R, *et al.* Lavender and coriander essential oils and their main component linalool exert a protective effect against amyloid- β neurotoxicity. *Phytother Res* 2020; 35(1):486-493.
<https://doi.org/10.1002/ptr.6827104>
104. Rana A, Samtiya M, Dhewa T, Mishra V, Aluko RE. Health benefits of polyphenols: A concise review. *J Food Biochem* 2022; 46(10): e14264.
<https://doi.org/10.1111/jfbc.14264>