

Research Article

DOI: 10.15740/HAS/AJSS/12.1/135-142

Effect of different growth regulator combinations on the per cent media browning in walnut *in vitro* studies using MS medium

■ I. A. LONE, F. A. MISGER AND F. A. BANDAY

Received : 10.03.2017; Revised : 28.04.2017; Accepted : 11.05.2017

MEMBERS OF RESEARCH FORUM:

Corresponding author :

I. A. LONE, Regional Research
Station (SKUAST-K), WADURA (J&K)
INDIA**Summary**

The present investigation on effect of different growth regulator combinations on the per cent media browning in walnut *in vitro* studies using MS medium was carried out in order to document the available genetic variability in walnut germplasm and to select elite walnut genotypes possessing superior attributes and quality traits. During the survey, data was recorded on one hundred fifty two (152) walnut trees growing in different areas of Kashmir valley. The study also involved establishment of response of elite walnut selections to different plant growth regulators in shoot morphogenesis. Woody species have been found to be far more difficult to clone *in vitro* than herbaceous plants. Poor response of the explants from mature woody species to *in vitro* manipulation is usually associated with the problem of browning and explant necrosis. The present studies were conducted on forced explants from three walnut selections (SKUAST 002, SKUAST 008, SKUAST 010). Murashiage and Skoog's basal medium supplemented with 0.3 mg/l⁻¹ Benzylamino purine and 0.1 mg/l⁻¹ indole-3-butyric acid gave best response in the establishment of initiating cultures, minimum media browning (80.44 %), minimum explant browning (78.22 %) and minimum mean browning score per explant (9.17 %). The survival (23.45%) and growth of the cultures (21.77%) was also found to be maximum in MS medium supplemented with BAP 0.3 mg l⁻¹ and IBA 0.1 mg l⁻¹.

Key words : Wlanut, *Juglans regia* L., Variability, Yield component, Quality traits, Shoot morphogenetic response

Co-authors :

F. A. MISGER AND F. A. BANDAY,
Regional Research Station (SKUAST-
K), WADURA (J&K) INDIA

How to cite this article : Lone, I.A., Misger, F.A. and Banday, F.A. (2017). Effect of different growth regulator combinations on the per cent media browning in walnut *in vitro* studies using MS medium. *Asian J. Soil Sci.*, 12 (1) : 135-142 : DOI : 10.15740/HAS/AJSS/12.1/135-142.

Introduction

The persian walnut (*Juglans regia* L.), known as the English walnut, belongs to the family Juglandaceae. English walnut has its origin in the eastern Europe, Asia minor and points eastward to Himalayan mountains. The native habitat of walnut extends from the Carpathian mountains to Europe across Turkey, Iraq, Afghanistan, South Russia and further eastward into the foot hills of the Himalayas. In India walnuts are usually grown in the

mid hill areas of Jammu and Kashmir, Himachal Pradesh and upper hills of Uttarakhand and Arunachal Pradesh. The soil most suitable for its cultivation should be well-drained and deep silt loamy containing organic matter in abundance. It should not have a fluctuating water level, hard pan and/or sandy sub-soil with alkaline reaction. A soil 2.5 to 3.0 m deep gives best results because the roots can penetrate deep and utilize residual soil moisture during dry spell and also make available sufficient nutrients. Furthermore, availability of sufficient moisture

in the leaves can reduce the damage due to sun burning of leaves, shoots and young fruits. Walnut is grown commercially in about 48 countries with an area of 66, 58, 966 hectares. The world walnut production is about 16, 70, 109 MT. The chief walnut producing countries are China (22%), USA (20%), Iran (12%) and Turkey (10%) (Anonymous, 2007 and 1984). India accounts for about 2.0 per cent of the world production. In India, Jammu and Kashmir is leading both in area as well as in production with an area of 82.04 thousand ha and production of 146.78 thousand tonnes. However, the productivity level of 1.79 t ha⁻¹ is far below than other countries. Himachal Pradesh has an area of 6.54 thousand ha with a production of 1.24 thousand tonnes and productivity level of 0.19 t ha⁻¹; while Uttarakhand has an area of 19.26 thousand ha with a production of 8.73 thousand tonnes and productivity level of 0.45 t/ha and Arunachal Pradesh has an area of 2285 ha with a production of about 51 tonnes and productivity level of 0.022 t/ha. In the state of Jammu and Kashmir, Anantnag is the leading district both in area as well as production corresponding to an area of 13647 ha and production of 41180 tonnes with a productivity level of 3.01 t ha⁻¹, followed by the Kupwara district that covers an area of 8175 ha with 22103 tonnes production and a productivity level of a 2.70 t ha⁻¹. Kulgam ranks 6th in area and 3rd in production in the J&K state and has the highest productivity of 3.52 t ha⁻¹, which is even higher than that of USA. This indicates that the state has the right type of agro-climatic conditions and vast potential to produce export quality walnut and kernels. Micro propagation studies in walnut are not so well established nor any fool proof protocol is yet developed for efficient and faster multiplication of superior plants. The presence of phenolic compounds and entophytic bacteria are still the main limiting factors for establishing plant micro propagation in walnuts. The use of young vegetal material is the usual technique for *in vitro* set up of walnut (Driver and Kuniyuki, 1984 and Jay-Allemand *et al.*, 1993). Quality in regeneration of *in vitro* plant material is correlated with maintenance of mother plants in the controlled environments, with regular hormone application and proper choice of physiological stage for collecting materials. The correct temperature in growth chambers is essential for a proper regeneration as well for subsequent multiplication (Dolcet-Sanjuan *et al.*, 1993). The addition of PVP to the culture medium as well as the substitution of agar by gelrite are the main factors

reported for the control of phenolic compounds.

The current methodology of woody crop rooting by a biotopic process is well documented in walnut (Driver *et al.*, 1984) with the use of IBA. Walnut is hard to propagate through micro propagation. Various attempts have been made using different types of explants, media, culture condition and rooting techniques (Driver and Kuniyuki, 1984). Poor proliferation and rooting rate is one of the main obstacles that limit the micro propagation efficiency in walnut. Intensive and well planned research is needed to develop a perfect protocol for micro propagation for this crop. Genotype plays a major role in vegetative propagation, in particular for micro propagation.

In many cases the propagation ratio can be improved by using a stronger cytokinin or increasing its concentration. However, this can sometimes have detrimental effects in the later stages of micro propagation. Micro propagation studies have also been carried out in some other species of nuts and similar trees like hazelnut (Radojevic *et al.*, 1975; Mele and Messeguer, 1983 and Perez *et al.*, 1983), chestnut (Vieter and Vieter, 1980) and almond (Mehra and Mehra, 1974). But reports on *in vitro* walnut culture are scarce.

Resource and Research Methods

Shoot morphogenetic studies :

The investigation was carried out at the Tissue Culture Laboratory of Regional Research Station Wadura, SKUAST (K), during the year 2014-2015.

The investigation involved shoot morphogenetic response in walnut to different treatment combinations of growth hormones.

Plant materials used:

The mature and bearing elite genotypes of walnut selections *viz.*, SKUAST-002, SKUAST-008 and SKUAST-010 growing at the Experimental Farm of the Division of Polmology at the main campus Shalimar were used as stock plants in the present study. The plant materials were subjected to uniform cultural practices.

Chemicals used:

The major and minor elements required for the preparation of media were obtained from the Hi Media Pvt. Ltd. The amino acids, vitamins and plant growth regulators used were obtained from the Sigma Chemicals.

Glass wares:

The glass ware used for the experiment was obtained from the Borosil Glass-ware. Before use the glassware was soaked in potassium dichromate- nitric acid solution for six hours followed by thorough washing in a jet of tap water so as to completely remove all traces of dichromate solution. They were then soaked in detergent solution (Teepol 1%) overnight and were thoroughly washed in tap water and rinsed twice with double distilled water. The glass-ware was then dried in hot air oven at 100°C for 24 h and later stored under aseptic conditions till use.

Culture media :

Selection of culture media:

Murashige and Skoog (1962) medium was used for the present investigation.

Inositol (100mg), sucrose (30g) and agar (7.0g) required for one litre culture medium were added directly at the time of preparation of culture media.

Adjuvants to the basal medium :

The MS basal medium was appropriately supplemented with vitamins, aminoacids and growth

regulators. Stock solution of organics (vitamins and amino acids) were prepared in double distilled water.

Preparation of the culture media :

The required quantity of sucrose was dissolved in double distilled water and to this stock solution macronutrients, micronutrients, vitamins and growth substance were added as per treatment required. The media of various formulations were prepared.

The pH of the medium was adjusted to 5.5 with HCl or 0.1 N NaOH prior to the addition of agar. Agar (7.0 g⁻¹) was dissolved in boiling distilled water and added to the medium and the volume was made up with distilled water. The medium was boiled and the hot medium was then immediately dispensed in the test tubes or conical flasks which were plugged tightly with non-absorbent cotton plugs and covered with aluminum foil. The medium was autoclaved at 15 psi (121°C) for 15 minutes (Dodds and Robert, 1982).

Transfer area and aseptic manipulations :

All the aseptic manipulations like surface sterilization, preparation and inoculation of explants and subsequent subculturing were carried out under hood of clean laminar air flow chamber. The working table of

Table A : Composition and preparation of stock solutions for Murashige and Skoog (1962) medium

Stock solution	Components	Chemical formula	Quantity (mg l ⁻¹)	Quantity for 20 litres	Quantity of stock solution prepared	Conc. of stock solution
A	Ammonium nitrate	NH ₄ NO ₃	1650	33.00 g	200 ml	100 x
B	Potassium nitrate	KNO ₃	1900	38.00 g	200 ml	100 x
C	Potassium iodide	KI	0.83	16.6 mg	200 ml	100 x
	Boric acid	H ₃ BO ₃	6.2	124 mg		
	Potassium phosphate, monobasic	KH ₂ PO ₄	170	3.4 g		
	Molybdic acid (Sodium salt) 2H ₂ O	Na ₂ MoO ₄ .2 H ₂ O	0.25	5 mg		
	Cobalt chloride hexahydrate	CoCl ₂ . 6H ₂ O	0.025	0.5 mg		
D	Calcium chloride, dihydrate	CaCl ₂ . 2H ₂ O	440	8.8 g	200 ml	100 x
E	Magnesium sulphate, heptahydrate	MgSO ₄ . 7H ₂ O	370	7.4 g	200 ml	100 x
	Zinc sulphate, heptahydrate	ZnSO ₄ . 7H ₂ O	8.6	172 mg		
	Managanese sulfatem tetrahydrate	MnSO ₄ . 4H ₂ O	22.3	446 mg		
	Cupric sulfate pentahydrate	CuSO ₄ . 5H ₂ O	0.025	0.5 mg		
F	Ethylenediaminetera acetic acid disodium salt	Na ₂ EDTA.2H ₂ O	37.3	746	200 ml	100 x
	Ferrous sulfate heptahydrate	FeSO ₄ .7H ₂ O	27.8	556 mg		
G	Glycine	-	2.00	40 mg	200 ml	100 x
	Nicotinic acid	-	0.5	10		
	Pyridoxine HCl	-	0.5	10		
	Thiamine HCl	-	0.1	2		

the laminar air flow chamber was first surface sterilized with absolute alcohol followed by ultra violet light for 30 minutes. The forceps and scalpel used for the inoculation as well as the Petri dishes were first steam sterilized in an autoclave at 121°C for 20 minutes and later flame sterilized before each inoculation. The hands were cleaned and wiped with alcohol before working. Further, the standard general procedure for sterile technique suggested by Street (1977) was followed.

Incubation chamber :

The cultures were incubated at 24±1°C in an air condition culture room with a 16 h photoperiod (3500 lux).

Collection and preparation of explant :

Forcing of stock plant:

This experiment consisted of field grown adult shoots and artificially forced walnut cuttings of the genotypes SKUAST-002, SKUAST-008 and SKUAST-010 of walnut. For forcing the dormant cuttings of 15-20 cm length were collected in January. Dormant cuttings were treated with 0.2 per cent captan and stored in cold store at 4±3°C in polythene bags for two months. Cuttings were withdrawn from cold store as and when required and placed in jars containing distilled water and kept in incubation chamber. The cuttings were incubated at 24±1°C in an air condition room with 16 hours of Photoperiod (3.5 lux). Sprouting of dormant buds occurred after one month of transferring to incubation chamber which served as explant source. Shoot tips from all the three genotypes were excised and cultured on semi-solid and stationery liquid medium for establishment. Each treatment combination consisted of 25 explants and was replicated 3 times for each genotype.

Surface sterilization and inoculation :

The explants were surface sterilized with 0.1 per cent (w/v) mercuric chloride solution for 10 minutes and then washed thoroughly with double distilled sterile water (5-6 times) under aseptic conditions in the laminar flow chamber. The explants were then placed on the medium in such a manner that conformed to the original polarity and were exposed above the surface of medium.

Details of *in vitro* studies :

Details of growth regulators:

The explants were cultured in Murashige and Skoogs (1962) basal medium (MS) supplemented with different concentrations of growth regulators. The details of the combinations of different concentration of growth regulators are presented Table B.

Table B: Details of growth regulators				
Growth regulators	Concentration used			
BAP	0.3 mg l ⁻¹	0.6 mg l ⁻¹	0.9 mg l ⁻¹	1.2 mg l ⁻¹
IBA	0.1 mg l ⁻¹	-	-	-

Treatment combinations used :

$$T_1 = \text{BAP } 0.3\text{mg l}^{-1} + \text{IBA } 0.1\text{mg l}^{-1}$$

$$T_2 = \text{BAP } 0.6 \text{ mg l}^{-1} + \text{IBA } 0.1 \text{ mg l}^{-1}$$

$$T_3 = \text{BAP } 0.9 \text{ mg l}^{-1} + \text{IBA } 0.1 \text{ mg l}^{-1}$$

$$T_4 = \text{BAP } 1.2 \text{ mg l}^{-1} + \text{IBA } 0.1 \text{ mg l}^{-1}$$

The data recorded on various parameters of tissue culture studies were put to statistical analyses. Values in per cent were transformed to arc sin (angular) or square root transformation before analysis. The mean performance of the cultivars and growth regulator combinations was presented for experimental findings alongwith their interaction (cultivars x growth regulator combination).

Survival of shoot apices :

Survival of explants was recorded within 8 days of inoculation in each treatment combination for each cultivar and expressed as per cent as follows :

$$\text{Per cent survival of explants} = \frac{\text{Number of explants survived}}{\text{Total number of explants culture}} \times 100$$

Per cent growing cultures :

Number of explants that showed growth were also recorded within 3 weeks of inoculation and expressed in per cent.

$$\text{Per cent growing cultures} = \frac{\text{Number of explants showing growth}}{\text{Total number of explants cultured}} \times 100$$

Explants showing browning :

Explants that showed browning within 8 days were recorded and expressed as per cent for each treatment.

$$\text{Per cent explants browning} = \frac{\text{Number of explants browned}}{\text{Total number of explants cultured}} \times 100$$

Mean browning score per explant :

Explants that showed browning were scored for browning intensity as under :

- 0 = No browning
- 1 = Browning on excised wound tissue.
- 2 = Partial browning on outer tissue of explant
- 3 = Complete browning of the outer tissue of explant
- 4 = Browning of the entire explant tissue.

Explants showing substrates browning :

Substrate browning (irrespective of explant browning) was recorded for each culture treatment in a genotype and expressed as per cent.

$$\text{Per cent substrate browning} = \frac{\text{Number of culture tubes showing media browning}}{\text{Total number of explants cultured}} \times 100$$

Volume of browned media per culture :

Volume of the browned media was calculated on the basis of diameter of browned media surrounding the explant and depth of the browned media by the following relationship.

$$V = f \times r \times d$$

where,

V = Volume in cubic millimetres.

r = Radius of the browned media in millimetres

d = Depth of the browned media in millimetres.

Browning score of the explant and volume of browning media in each treatment was worked out per explant basis. The values were expressed finally as volume of browned media (mm)³ x 1/100 per explant. Observation on growth rating of explant and percentage of callused cultures was recorded after 30 days of culture initiation and growing culture were again transferred to fresh medium of the same composition.

Growth rating of explants :

Explants that revived and showed growth were scored for growth vigour as under :

- 1 = Low growth rate
- 2 = Medium growth rate

3 = High growth rate.

Percentage of callused cultures :

Explants that survived and revealed growth were scored for callusing and expressed as per cent.

$$\text{Per cent cultivars showing callusing} = \frac{\text{Number of explants expressing callus}}{\text{Number of surviving and growing explants}} \times 100$$

Statistical analysis of the data :

The data recorded for phenotypic variability of trait were put to statistical analyses and different parameters estimated.

Genetic variability studies :

Data for each trait of each earmarked *in-situ* tree was recorded over two crop seasons (*Kharif* 2008 and *Kharif* 2009) and statically analyzed.

Estimation of variability parameters :

Different variability parameters estimated were as under :

Mean and range :

The population mean and range of each trait was expressed in the same units as that of the trait.

Phenotypic co-efficient of variation :

Phenotypic co-efficient of variation (PCV) was expressed as:

$$\text{PCV} = \frac{\sqrt{2p}}{X} \times 100$$

Genotypic co-efficient of variation :

Genotypic co-efficient of variation (GCV) was expressed as:

$$\text{GCV} = \frac{\sqrt{2g}}{X} \times 100$$

Components of variance :

The variance for each trait comprised variance

Table C : Analysis of variance				
Source of variation	Df	Expected mean	Sum of squares	Variance ratio
Replications (years)	y ⁻¹	MS ₁	² e+g ² y	MS ₁ /MS ₃
Cultivars (genotypes)	g ⁻¹	MS ₂	² e+y ² g	MS ₂ /MS ₃
Replications x cultivars	(y ⁻¹)(g ⁻¹)	MS ₃	² e	

arising from the cultivars difference (genotypic variance) and that arising from the environmental factor (in this case years). The expected mean sum of squares would be as under:

$$MS \text{ for cultivars} = \uparrow^2e + y \uparrow^2g = MS_2$$

$$MS \text{ for error} = \uparrow^2e = MS_3$$

$$\text{Genotypic variance, } ^2g = \frac{MS_2 - MS_3}{Y}$$

$$\text{Phenotypic variance} = \uparrow^2p = \uparrow^2g + \uparrow^2e$$

Heritability (broad sense) :

It was measured as the ratio of genotypic variance to phenotypic variance.

$$h^2(\text{bs}) = \uparrow^2g / \uparrow^2p$$

Genetic gain :

Genetic gain (% of mean) was expressed as :

$$\text{Genetic gain} = \frac{h^2(\text{b.s}) \times p \times k}{\bar{X}}$$

where,

$h^2(\text{b.s})$ = Heritability in broad sense,

σp = Phenotypic standard deviation

K = Selection intensity (the value of k at 5% was taken as 2.06) and

\bar{X} = Population mean of the trait.

Research Findings and Discussion

Explants of three elite walnut cultivars viz., SKUA-

002, SKUA-008 and SKUA-010 were studied for their response to shoot morphogenesis under micro propagation (tissue culture) techniques. The standard basal medium of Murashige and Skoog (1962) was supplemented with BAP (0.3, 0.6, 0.9 and 1.2 mg l⁻¹) together with a uniform concentration of 0.1 mg l⁻¹ of IBA. The results pertaining to different parameters of *in vitro* studies are briefly presented in the following.

Media browning :

Effect of growth regulators and cultivars on the per cent media browning revealed that 80.00 to 85.33 per cent culture tubes showed browning in different treatments. The per cent values were transformed to square root and analyzed for significance of treatment effects. Cultivars exhibited no effect on the media browning development where the transformed values (mean values between 82.00 to 82.33%) were non-significant. Among growth regulators also only T₂ (BAP 0.6 mg l⁻¹ + IBA 0.1 mg l⁻¹) had significantly higher per cent media browning than T₁ (BAP 0.3 mg l⁻¹ + IBA 0.1 mg l⁻¹). Rest of the comparisons had non-significant influences on the per cent media browning.

Higher proportion of explant browning was observed in the *in-vitro* studies in all the three accessions and the effect of growth regulators also did not have sizeable impact. The explant browning in all the treatment combinations ranged from 77.33 to 85.33 per cent. Walnut accessions had almost similar and non-significant impact on explant browning. However, application of BAP 0.3

Walnut accessions	Growth regulators				Mean
	T ₁	T ₂	T ₃	T ₄	
SKUA-002	8.97* (80.00)	9.19 (84.00)	9.11 (82.66)	9.04 (81.33)	9.08 (82.00)
SKUA-008	8.97 (80.00)	9.26 (85.33)	9.11 (82.66)	9.04 (81.33)	9.10 (82.33)
SKUA-010	9.04 (81.33)	9.19 (84.00)	9.04 (81.33)	9.04 (81.33)	9.08 (82.00)
Mean	8.99 (80.44)	9.21 (84.44)	9.09 (82.22)	9.04 (81.33)	9.08 (82.11)

*Square root transformed value of the mean = ($\sqrt{0.5 + \text{mean value}}$)

Critical difference at 5%	
Accessions (AC)	0.164
Growth regulators (GR)	0.189
Interaction (AC x GR)	NS

T₁ = BAP 0.3 mg l⁻¹ + IBA 0.1 mg l⁻¹

T₂ = BAP 0.6 mg l⁻¹ + IBA 0.1 mg l⁻¹

T₃ = BAP 0.9 mg l⁻¹ + IBA 0.1 mg l⁻¹

T₄ = BAP 1.2 mg l⁻¹ + IBA 0.1 mg l⁻¹

NS = Non-significant

mg l⁻¹ + IBA 0.1 mg l⁻¹ significantly reduced the explant browning as compared to higher concentration of BAP (0.6, 0.9 and 1.2 mg l⁻¹) + IBA 0.1 mg l⁻¹.

Application of BAP 0.3 mg l⁻¹ + IBA 0.1 mg l⁻¹ revealed significantly lower explant browning in the cultivars SKUA-002 and SKUA-008 as compared to most of other interaction (cultivars x growth regulator) effects.

Ichihashi and Kako (1977) reported that the browning of cattleya shoot tips was most effectively controlled by the addition of antioxidants (potassium cyanide, ascorbic acid, cysteine, indolebutyric acid) into stationary liquid medium. Monaco *et al.* (1977) suggested that coffee explants should be pretreated with an antioxidant (e.g. cysteine, HCl, ascorbic acid) to reduce sbrowning.

Tissues are injured during explant excision from stock plants. This often causes the release of various compounds that are oxidised (Robinson, 1983) by peroxidases (Loomis and Battaile, 1966 and Vaugh and Duke, 1984) or polyphenol oxidases (Hu and Wang, 1983 and Mayer and Harel, 1979) and turn brown or black resulting in darkening of both tissue and culture medium. Polyphenol oxidases, including catechol oxidases, are confined to particles in healthy and non-senescent tissues (Vaugh and Duke, 1984) when explant is injured in course of explant preparation. Phenolic compounds that are largely localized in vacuole are mixed with the contents of plastids or other organelles resulting in formation of dark or brown pigmentation associated with oxidised polyphenolic compounds (Compton and Preece, 1986). The oxidative browning caused by the oxidation products of phenols inhibit enzyme activity (Hu and Wang, 1983) and may result in the decline or lethal darkening of the explants. The build up of toxic brown or black substance on wounded surface of cultured tissues was listed as one of the basic problems of micropropagation by Pierik (1988). Interaction arising among accessions and growth regulators revealed non-significant influence. *In vitro* studies revealed very low survival, growth and callusing of explants. The development of browning on explants, media and its volume was significantly high. Accessions played no role in the *in vitro* studies, but growth regulator BAP at lower concentration significantly increased survival and growth of explants and reduced browning. Callusing increased in the growing cultures with increase in BAP concentration from 0.6 to 1.2 mg l⁻¹.

Literature Cited

- Anonymous (1984). *Descriptor for walnut (Juglans regia spp.)*. International Plant Genetic Resource Institute, Rome Italy, pp. 28-40.
- Compton, M.E. and Preece, J.E. (1986). Exudation and explant establishment. *Newsletter Internat. Assoc. Plant Tissue Cult.*, **50** : 9-18.
- Dodds, J.H. and Robert, L.W. (1982). *Experiments in plant tissue culture*. Cambridge University Press, London pp. 178.
- Dolcet-Sanjuan, R., Calveria, E. and Alled, J. (1993). La micropropagation en clones de *Juglans regia* e hybrids de *J. regia* x *J. Nigra*. *Una Soluacion Parcial a la Propagation Conal del nogal*, **7** : 114-119.
- Driver, J.A. and Kuniyuki (1984). *In vitro* propagation of paradise walnut rootstock. *Hort. Sci.*, **19** : 507-509.
- Driver, J.A., Kuniyuki, A.H. and Hamber, P.C. (1984). Studies on *in vitro* culture of paradise walnut (*Juglans regia* L.). *Hort. Sci.*, **19** : 510-516.
- Hu, C.Y. and Wang, P.J. (1983). Meristem shoot tip and bud cultures. In : *Handbook of Plant Cell Cultures* [Eds. D.A. Evans; W.R. Sharp, P.V. Ammirata and Y. Yamada]. Mac Millan Press, New York pp. 177-217.
- Ichihashi, S. and Kako, S. (1977). Studies on clonal propagation of *Cattleya* through tissue culture method II, Browning of *Cattleya*. *J. Japanese Society of Hort. Sci.*, **46** : 325-330.
- Jay-Allemand, C.H., Peng, S., Capelro, P. and Cornu, D. (1993). Micropropagation of walnut hybrid tree : Some factors involved in rooting. *Acta Horticulturae*, **311** : 117-124.
- Loomis, W.D. and Battaile, J. (1966). Plant phenolic compounds and the isolation of plant enzymes. *Phytochemistry*, **51** : 423-438.
- Mayer, A.M. and Harel, E. (1979). Polyphenol oxidases in plants. *Phytochemistry*, **18** : 193-215.
- Mehra, A.S. and Mehra, P.N. (1974). Organogenesis and plantlet formation *in vitro* in almond. *Botanical Gazette*, **135** : 61-73.
- Mele, E. and Messeguer, J. (1983). *Clonal propagation of Corylus avellana* L. *in vitro*. Proceedings of the International Plant Propagators Society (Italy).
- Monaco, L.C., Sondahl, M.R., Carvalho, A., Chrocomo, O.J. and Sharp, W.R. (1977). Application of tissue culture in the improvement of coffee. In : *Plant Cell, Tissue and Organ Culture* [Eds. J. Reinert and Y.P.S. Bajaj]. Springer-Verlag Berlin, pp. 109-129.
- Murashige, T. and Skoog, F. (1962). A revised medium for

rapid growth and bioassays with tobacco tissue cultures. *Plant Physiol.*, **15** : 473-497.

Perez, C., Fernander, B. and Rodriguez, R. (1983). *In vitro* plantlet regeneration through asexual embryogenesis in cotyledonary segments of *Corylus avellana* L. *Plant Cell Reproduction*, **2** : 226-228.

Pierik, R.L.M. (1988). Handicaps for the large scale commercial application of micropropagation. *Acta Horticulturae*, **230** : 63-71.

Radojevic, L.R., Vujicic, R., Neskoncm, M. (1975). Embryogenesis in tissue culture of *Corylus avellana* L. *Pflanzezenphysiologia*, **77** : 33-41.

Robinson, T. (1983). *The organic constituents of higher plants.* (5th Ed.). Cordus Press, North America (U.S.A) **60**: 161-

165.

Street, H.E. (1977). Laboratory organization. In : *Plant tissue and cell culture* (Ed. H.E. Street). Blackwell Scientific Publications, Oxford. pp. 11-30.

Vaugh, K.C. and Duke, S.O. (1984). Function of polyphenol oxidase in higher plants. *Plant Physiol.*, **60** : 106-110.

Vieter and Vieter (1980). The significance of allelopathy chestnut cultural systems. *North America Growers Nut Association. Annual Report*, **72** : 117-134.

Webliography

Anonymous (2007). *Corylus colorna*. Hazel, Turkish filbert, Turkish Hazelnut, Turkish tree hazel. [*c.f.http://: zipcodezoo.com/ plants/c/ corylus- colornaasp.*].

★ ★ ★ ★ ★ of Excellence ★ ★ ★ ★ ★
12th Year