

RESEARCH ARTICLE

Incidence of seed mycoflora in different cultivars of safflower

■ D. Amrutha Gayathri and V. Krishna Rao

SUMMARY

An investigation was conducted to detect the associated seed mycoflora in safflower and its control. A total of 19 safflower seed samples were collected from major growing areas of safflower. Blotter method and agar plate methods were used for detection of seed mycoflora of safflower seeds. Across the two methods adopted, a total of seven fungal genera including *Alternaria*, *Aspergillus*, *Chaetomium*, *Rhizopus*, *Curvularia* and *Fusarium* were detected. The fungi detected were identified based on their cultural and morphological characteristics. The fungal species namely *Alternaria carthami*, *Aspergillus niger* were found associated with all the tested hybrids/varieties/germplasms while species of *Chaetomium* sp., *Rhizopus* sp., *Curvularia* sp. and *Fusarium* sp. were not detected in some of the hybrids/varieties/germplasms. Among the seven fungal species detected the occurrence of *Alternaria carthami* was high 47.94 per cent followed by *Aspergillus niger*. The cultivar, Nira showed higher per cent incidence of seed mycoflora. Per cent incidence of seed mycoflora varied across the methods adopted and cultivars tested the highest per cent incidence of 46% was observed with the fungus *Alternaria carthami* on Nari NH1 in blotter method. Out of the two methods tested blotter method was found superior over agar plate method in which higher number of fungi were recorded.

Key Words : Safflower, Mycoflora, *Alternaria carthami*, Seed borne

How to cite this article : Gayathri, D. Amrutha and Rao, V. Krishna (2018). Incidence of seed mycoflora in different cultivars of safflower. *Internat. J. Plant Sci.*, 13 (1): 114-117, DOI: 10.15740/HAS/IJPS/13.1/114-117.

Article chronicle : Received : 29.07.2017; Revised : 25.11.2017; Accepted : 10.12.2017

Safflower (*Carthamus tinctorius* L.) is one of the major Rabi oilseed crops in India, it belonging to the family *Asteraceae* is an important source of oil and proteins. The crop is cultivated over an area of

MEMBERS OF THE RESEARCH FORUM

Author to be contacted :

D. Amrutha Gayathri, Department of Plant Pathology, College of Agriculture, Professor Jayasankar Telangana State Agricultural University, Rajendranagar, Hyderabad (Telangana) India

Address of the Co-authors:

V. Krishna Rao, Department of Plant Pathology, College of Agriculture, Professor Jayasankar Telangana State Agricultural University, Rajendranagar, Hyderabad (Telangana) India

287 thousand hectares with the production of 658 kg ha⁻¹ (CMIE, 2009) and production of 179 thousand t (CMIE, 2010) in India. One of the major constrain in safflower production is the lack of quality seed at the time of planting. The health of safflower plant is affected by number of fungal diseases. The diseases are Fusarial wilt (*Fusarium oxysporum* f.sp. *carthami*), *Alternaria* leaf spot (*Alternaria carthami*), Rust (*Puccinia carthami*), Bud blight (*Phytophthora drechslera*), Ramularia leaf spot (*Ramularia carthami*), *Cercospora* leaf spot (*Cercospora carthami*), Bacterial spot (*Pseudomonas syringae*) and Cucumber mosaic virus.

Seed borne fungi are carried over by infested seeds. They cause deterioration in soil, before seed germination causing seedling mortality and infection of foliage at adult stages. Fungi including *Alternaria carthami*, *Aspergillus flavus*, *Aspergillus niger*, *Cladosporium*, *Macrophomina*, *Chaetomium* sp., *Rhizopus* sp., *Curvularia* sp. and *Fusarium* sp. were found associated with safflower, *Alternaria carthami* was the most destructive pathogen of safflower, as it cause leaf spot and blight.

MATERIAL AND METHODS

The seed samples of 19 safflower cultivars were collected from major growing areas of Telangana. The seeds were collected in polythene bags and stored at room temperature of 25±2°C.

The collected seed samples were analyzed for the presence of seed mycoflora by employing standard blotter method and agar plate method (ISTA, 1976). In all the methods 400 seeds taken randomly from each sample were subjected to analysis without sterilization.

For conducting incubation tests sterile glass Petri plates of 9 cm diameter were used. In blotter method, the seeds were placed on three layers of moistened blotter papers in Petri plates. In agar plate method, the seeds were placed over the surface of solidified potato dextrose agar medium in Petri plates. In both the cases, the seeds were plated in Petri plates at the rate of 10 seeds/ plate at equidistance and incubated in an incubator set to 25±2°C temperature for seven days. The incubated seeds was observed on eighth day by using steriobinocular and compound microscope.

The mycoflora associated with seed were identified using key given by Barnett and Hunter (2003), Booth (1971) and Subramanian (1971).

RESULTS AND DISCUSSION

The analysis of seed mycoflora of 19 safflower cultivars using standard blotter and agar plate methods showed the association of seven fungal species. The fungi detected were identified based on their cultural and morphological characteristics. The fungal species

Table 1 : Incidence of seed mycoflora in different cultivars of safflower in different locations of Telangana*

Sr. No.	Name of the district	Sr. No.	Hybrid/variety / germplasm	<i>Alternaria carthami</i>	<i>Aspergillus flavus</i>	<i>Aspergillus niger</i>	<i>Chaetomium</i> sp.	<i>Curvularia</i> sp.	<i>Fusarium</i> sp.	<i>Rhizopus</i> sp.	Total mycoflora
Per cent incidence of seed mycoflora											
1.	RangaReddy (DOR)	1.	A-1	32	11	24	-	-	-	3	70
		2.	JSF414	37	10	22	-	23	-	-	92
		3.	Nari 38	13	-	29	-	35	-	-	77
		4.	Bhima	38	9	21	12	-	-	-	80
		5.	Nari 6	36	-	20	4	-	-	4	64
		6.	SSF 658	8	10	11	-	-	-	5	34
		7.	Nari NH 1	46	12	18	2	-	-	-	78
		8.	Nari NH 15	15	10	21	-	-	-	-	46
2.	Khammam	9.	Manjeera	39	4	4	-	3	-	4	54
3.	Adilabad	10.	Bhima	25	7	10	10	1	-	6	59
4.	Nalgonda	11.	Manjeera	32	3	5	2	5	0	4	51
5.	Mahaboobnagar	12.	Sagara mutyalu	28	4	3	6	3	0	5	49
		13.	Sagara mutyalu	45	-	20	4	22	--	-	91
Ranga Reddy (Tandur)	14.	Manjeera	16	5	3	6	19	1	-	-	50
	15.	Nira	32	15	10	4	20	-	-	6	95
	16.	TSF-1	30	-	26	-	-	2	-	-	58
	17.	GMU-5536	38	-	13	-	-	-	-	-	51
	18.	GMU-5653	36	11	5	-	5	4	-	-	61
	19.	IVT-1002	22	13	10	4	10	-	-	3	62
	Total			560	92	255	54	146	7	40	1168
Per cent				47.94	7.87	21.8	4.62	12.5	0.59	3.42	

* Standard blotter method

Table 2 : Incidence of seed mycoflora in different cultivars of safflower in different locations of Telangana*

Sr. No.	Name of the district	Sr. No.	Hybrid/variety/germplasm	<i>Alternaria carthami</i>	<i>Aspergillus flavus</i>	<i>Aspergillus niger</i>	<i>Chaetomium</i> sp	<i>Fusarium</i> sp	<i>Rhizopus</i> sp	Total mycoflora
Per cent incidence of seed mycoflora										
1.	Ranga Reddy (DOR)	1.	A-1	20	-	13	-	-	1	34
		2.	JSF414	26	6	11	-	4	2	49
		3.	Nari 38	2	3	18	-	-	-	23
		4.	Bhima	27	-	19	-	-	3	49
		5.	Nari 6	27	-	10	-	-	1	38
		6.	SSF 658	2	-	3	-	-	2	7
		7.	Nari NH 1	32	5	12	-	-	-	49
		8.	Nari NH 15	5	-	15	-	-	1	21
2.	Khammam	9.	Manjeera	34	1	12	7	7	1	62
3.	Adilabad	10.	Bhima	22	2	15	6	4	3	52
4.	Nalgonda	11.	Manjeera	30	4	16	5	-	3	58
5.	Mahaboobnagar	12.	Sagara mutyalu	32	2	14	-	4	-	52
6.	Ranga Reddy (Tandur)	13.	Sagara mutyalu	21	2	15	4	-	-	42
	RangaReddy (DOR)	14.	Manjeera	21	1	20	7	1	1	51
		15.	Nira	18	-	6	9	-	-	33
		16.	TSF-1	15	-	4	-	-	2	21
		17.	GMU-5536	14	-	8	-	5	-	27
		18.	GMU-5653	3	6	-	-	-	-	9
		19.	IVT-1002	17	-	5	-	4	2	28
			Total	368	32	216	38	29	23	705
			Per cent	52.2	4.5	30.6	5.4	4.1	3.3	

*Agar Plate method

detected through standard blotter method include *Alternaria carthami*, *Aspergillus flavus*, *Aspergillus niger*, *Chaetomium* sp., *Rhizopus* sp., *Curvularia* sp. and *Fusarium* sp. (Table 1) while, the agar plate method yielded six fungal species except *Curvularia* sp. (Table 2)

Per cent incidence of seed mycoflora varied across the methods adopted and cultivars tested. The highest per cent incidence of 46% was observed with the fungus *Alternaria carthami* on Nari NH1 in blotter method. Among the mycoflora detected *Alternaria carthami* (2-46 %) was recovered from all the cultivars in both the methods. Among the seven fungal species detected the occurrence of *Alternaria carthami* was high 46 per cent followed by *Aspergillus niger* (29 %). The cultivar, Nira showed higher per cent incidence of seed mycoflora. The fungal species viz., *Alternaria carthami*, *Aspergillus niger* was found associated with all the tested hybrids/varieties/germplasms while species of *Chaetomium* sp., *Rhizopus* sp., *Curvularia* sp. and *Fusarium* sp. were not detected in some of the hybrids/varieties/germplasms. The results also indicated that, the per cent incidence of *Rhizopus* sp. (0-6%), *Chaetomium*

sp. (0-12%) and *Fusarium* sp. (0-7%) was found less (Table 1 and 2).

Appearance of the colonies of *Alternaria carthami*, *Aspergillus flavus*, *Aspergillus niger*, *Chaetomium* sp., *Rhizopus* sp., *Curvularia* sp. and *Fusarium* sp. on the safflower seeds indicates their seed borne nature. Similar findings were reported by Padaganur and Anil Kumar (1976); Singh *et al.* (1987); Awadhiya (1992); Rajeswari *et al.* (2012) and Pushpavathi *et al.* (2012).

The predominance nature of *Alternaria carthami* is in agreement with the findings of Raghuwanshi and Deokar (2002) and Singh *et al.* (1987) who reported the highest percentage of association of *Alternaria carthami* with safflower seed. It was also proved as an externally seed borne fungus causing discoloration of seed in several safflower cultivars (Borkar and Shinde, 1989).

REFERENCES

- Awadhiya, G.K. (1992). Seed borne pathogenic mycoflora of safflower. *Crop Res.*, 5(2): 344-347.
- Barnett, H.L. and Hunter, B.B. (2003). *Illustrated genera of imperfect fungi*. Fourth edition APS Press, St. Paul, Minnestova. 92, 94.

- Booth, C. (1971). *The Genus Fusarium*. Common Wealth Mycological Institute, Kew Surrey, England. 271.
- Borkar, S.G. and Shinde, D. (1989). Detection of externally seed borne *Alternaria carthami* on safflower seeds. *Agric. Sci. Digest*, **9** (3):120-122.
- CMIE Agriculture (2009). Centre for Monitoring Indian Economy Pvt. Ltd. Andheri (East), Mumbai.
- CMIE Agriculture (2010). Centre for Monitoring Indian Economy Pvt. Ltd. Andheri (East), Mumbai.
- ISTA (1976). International rules for seed testing. *Seed Science & Technol.*, **4** : 51–177.
- Padaganur, G.M. and Anil Kumar, J.B. (1976). Seed mycoflora at safflower and its control. *Pesticides*, **10** (7) : 39-41.
- Pushpavathi, B, Rajender, J. and Narayan Reddy, P. (2012). Detection of seed mycoflora of safflower. *J. Oil Seeds Res.*, **29**: 364-367.
- Raghuwanshi, K.S. and Deokar, C.D. (2002). Studies seed borne mycoflora of safflower. *Sesame & Safflower Newslett. No. 17*.
- Rajeswari, B., Keshavulu, K. and Krishna Rao, V. (2012). Management of seed mycoflora of safflower. *J. Oil Seeds Res.*, **29** : 332-335.
- Singh, S.N., Agarwal, S.C. and Khare, M.N. (1987). Seed borne pathogenic mycoflora of safflower, their significance and control. *Seed Res.*, **15**:190-191.
- Subramanian, G.V. (1971). *Hyphomycetes*, Indian council of Agricultural Research, New Delhi, 930.

13th
Year
★★★★★ of Excellence ★★★★★